

### DarkSUSY darksusy-6.1.0

### Manual and description of routines

Documentation harvested from documentation and source directories with harvestdoc.pl Wed Jun 13 14:18:52 2018

http://www.darksusy.org

### Joakim Edsjö<sup>a\*</sup>, Torsten Bringmann<sup> $b\dagger$ </sup>, Paolo Gondolo<sup> $c\ddagger$ </sup>, Piero Ullio<sup>d</sup>, Lars Bergström<sup>a</sup>, Mia Schelke<sup> $\parallel$ </sup>, Edward A. Baltz<sup>\*\*</sup> and Gintaras Duda<sup>††</sup>

<sup>a</sup> The Oskar Klein Centre for Cosmoparticle Physics, Department of Physics, Stockholm University, AlbaNova, SE-106 91 Stockholm, Sweden.

<sup>b</sup> Department of Physics, University of Oslo, Box 1048, NO-0371 Oslo, Norway

<sup>c</sup> Department of Physics. University of Utah. 115 South 1400 East, Suite 201, Salt Lake City, UT 84112-0830, USA.

<sup>d</sup> SISSA and INFN, Sezione di Trieste, via Bonomea 265, 34136 Trieste, Italy.

<sup>\*</sup>E-mail address: edsjo@fysik.su.se

 $<sup>^{\</sup>dagger}\text{E-mail}$  address: torsten.bringmann@fys.uio.no

 $<sup>^{\</sup>ddagger}\mathrm{E}\text{-}\mathrm{mail}$ address: paolo.gondolo@utah.edu

<sup>&</sup>lt;sup>§</sup>E-mail address: ullio@he.sissa.it ¶E-mail address: lbe@fysik.su.se

<sup>&</sup>lt;sup>||</sup>E-mail address: schelke@...

<sup>\*\*</sup>E-mail address: eabaltz@physics.columbia.edu

<sup>&</sup>lt;sup>††</sup>E-mail address: gkduda@creighton.edu

### Abstract

DarkSUSY is a package for numerical dark matter calculations including, but not limited to, supersymmetric dark matter. This manual describes the theoretical background as well as details about the actual routines. Everything is not covered, but it should hopefully prove useful if you need more information than in our published articles.

**Disclaimer.** This manual is work in progress.

We try to keep it clear and up-to-date with the code, but there will be cases where changes/improvements in the code, for one reason or another, will not have propagated into the manual. Hence, check the actual code if you want to be certain about how a given process/feature is implemented.

### Contents

Ι	Prelude	7
1	Introduction	8
2	Quick start guide         2.1       Installation         2.1.1       Options for install         2.1.2       System requirements         2.12       Example programs         2.2       Example programs         2.2.1       Auxiliary example programs         2.2.2       Making your own example programs         2.3       Modifying individual subroutines or functions	<ol> <li>9</li> <li>10</li> <li>10</li> <li>10</li> <li>11</li> <li>12</li> </ol>
3	Guiding principles of the code         3.1 The main DarkSUSY library ds_core         3.2 Particle physics modules         3.2.1 Using a particle physics module         3.2.2 Adding a new particle physics module         3.3 Halo models         3.4 Interface functions         3.5 Commonly used functions         3.6 Replaceable functions	<ul> <li>13</li> <li>13</li> <li>14</li> <li>15</li> <li>16</li> <li>17</li> <li>17</li> <li>18</li> <li>19</li> </ul>
4	Comparison to previous DarkSUSY versions	<b>21</b>
5	Original articles	23
II	Main DarkSUSY routines in $src/$	<b>24</b>
6	an_yield: Annihilation yields in the halo – yields from simulations         6.1       Annihilation in the halo, yields – theory         6.1.1       Monte Carlo simulations	<b>25</b> 25 25
7	aux: General routines         7.1 General routines	<b>27</b> 27
8	aux_xdiag: Diagonalization routines 8.1 XDIAG	<b>28</b> 28
9	cr_aux: Cosmic rays – general 9.1 Cosmic Rays – auxiliary routines	<b>29</b> 29

#### CONTENTS

10	cr_axi: Cosmic rays – diffusion routines for axisymmetric distributions 10.1 Cosmic ray propagation in axially symmetric halos	<b>30</b> 30
11	cr_gamma: Cosmic rays – Gamma fluxes         11.1 Gamma rays from the halo – theory         11.2 Continuous gamma yields and line signals         11.3 Fluxes         11.4 Gamma rays from the halo – routines	<b>33</b> 33 34 34 35
12	cr_nu: Cosmic rays – Neutrino fluxes 12.1 Neutrino fluxes from the halo – theory	<b>36</b> 36
13	cr_ps: Cosmic rays – point sources 13.1 Cosmic Ray propagation for point sources	<b>37</b> 37
14	dd: Direct detection         14.1 Direct detection – theory	<b>38</b> 38
15	dmd_astro: Astrophysical source functions	39
16	dmd_mod: Dark matter distributions         16.1 Dark matter distributions – theory         16.1.1 Rescaling of the WIMP density         16.2 Implementation in DarkSUSY         16.2.1 Dark matter distributions – routines	<b>40</b> 40 40 41 42
17	dmd_vel: Dark matter velocity distributions 17.1 Dark matter phase-space distributions – theory	<b>43</b> 43
18	ini: Initialization routines         18.1 Initialisation routines	<b>44</b> 44
19	kd: Kinetic decoupling         19.1 Kinetic decoupling and microhalos (kd) – theory         19.1.1 Kinetic decoupling         19.1.2 The smallest protohalos         19.2 Kinetic decoupling – routines	<b>45</b> 45 45 46 47
20	rd: Relic density	48
	20.1 Relic density – theoretical background	$48 \\ 48 \\ 48 \\ 50 \\ 52$
	20.2 Relic density – numerical integration of the density equation	55 56 56 57
<b>21</b>	se_aux: Auxiliary routines for WIMP annihilation in the Sun/Earth 21.1 Sun and Earth models – auxiliary routines	<b>59</b> 59

#### CONTENTS

	00
22.1 Sun and Earth models – theory	60
22.2 Sun and Earth models – routines $\ldots$	60
	61
23 se_nu: Capture and annihilation in the Sun/Earth $\frac{1}{100}$	61
23.1 Neutrinos from the Sun and Earth – theory	61
23.1.1 Neutrino yield from annihilations	61
23.1.2 Evolution of the number density in the Earth/Sun $\ldots$ .	62
23.1.3 Approximate capture rate expressions	63
23.1.4 Earth and Sun composition	64
23.1.5 More accurate capture rate expressions	65
23.1.6 Accurate capture rates in the Earth for general velocity distributions	65
23.1.7 Accurate capture rates for the Earth for a Maxwell-Boltzmann veloci	ity dis-
tribution	67
23.1.8 Effects of WIMP diffusion in the solar system	68
23.2 Neutrinos from Sun and Earth – routines	60
	00
24 se vield: Yields from annihilation in the Sun/Earth	70
241 Muon vields from annihilation in the Earth/Sun – theory	70
24.1.1 Monte Carlo simulations with WimpSim	
25 si: Self-interactions	75
25.1 Dark matter self-interactions (si) – theory	75
25.2 Solf interactions – routines	· · · · · · 70
20.2 Sen-interactions – fournes	
III Particle physics modules in src models	78
III I afficie physics modules in sie_models	
26 Basic principles and common routines	79
26 Basic principles and common routines	<b>7</b> 9 70
<b>26</b> Basic principles and common routines 26.1 common/aux: auxiliary routines	<b>79</b> 
26 Basic principles and common routines         26.1 common/aux: auxiliary routines         26.2 common/sm: standard model	<b>79</b> 
<ul> <li>26 Basic principles and common routines</li> <li>26.1 common/aux: auxiliary routines</li></ul>	79 
<ul> <li>26 Basic principles and common routines</li> <li>26.1 common/aux: auxiliary routines</li></ul>	79 79 79 81
<ul> <li>26 Basic principles and common routines</li> <li>26.1 common/aux: auxiliary routines</li></ul>	79 79 79 
<ul> <li>26 Basic principles and common routines <ul> <li>26.1 common/aux: auxiliary routines</li> <li>26.2 common/sm: standard model</li> </ul> </li> <li>27 The empty model </li> <li>28 Generic decaying dark matter</li> </ul>	79 
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 79 79 81 82 83 84 85 85
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 79 79 81 82 83 84 85 85 
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 79 79 79 79 81 81 85 
<ul> <li>26 Basic principles and common routines 26.1 common/aux: auxiliary routines</li></ul>	79 
<ul> <li>26 Basic principles and common routines <ul> <li>26.1 common/aux: auxiliary routines</li> <li>26.2 common/sm: standard model</li> <li>27 The empty model</li> </ul> </li> <li>28 Generic decaying dark matter</li> <li>29 Generic WIMPs</li> <li>30 The Minimal Supersymmetric Standard Model <ul> <li>30.1 mssm/ac: Accelerator constraints</li> <li>30.1.1 Accelerator bounds</li> <li>30.2 mssm/an: (Co-)annihilation cross sections</li> <li>30.2.1.1 Annihilation cross sections</li> <li>30.2.1.1 Annihilation cross sections</li> <li>30.2.1.2 Coannihilation diagrams</li> </ul> </li> </ul>	79 79 79 79 79 81 82 82 84 85 86 
<ul> <li>26 Basic principles and common routines <ul> <li>26.1 common/aux: auxiliary routines</li> <li>26.2 common/sm: standard model</li> <li>27 The empty model</li> </ul> </li> <li>28 Generic decaying dark matter</li> <li>29 Generic WIMPs</li> <li>30 The Minimal Supersymmetric Standard Model <ul> <li>30.1 mssm/ac: Accelerator constraints</li> <li>30.1.1 Accelerator bounds</li> <li>30.2 mssm/an: (Co-)annihilation cross sections</li> <li>30.2.1.1 Annihilation cross sections</li> <li>30.2.1.2 Coannihilation diagrams</li> <li>30.2.1.3 Neutralino and chargino annihilation</li> </ul> </li> </ul>	79 
<ul> <li>26 Basic principles and common routines <ul> <li>26.1 common/aux: auxiliary routines</li> <li>26.2 common/sm: standard model</li> </ul> </li> <li>27 The empty model </li> <li>28 Generic decaying dark matter </li> <li>29 Generic WIMPs </li> <li>30 The Minimal Supersymmetric Standard Model <ul> <li>30.1 mssm/ac: Accelerator constraints</li> <li>30.1.1 Accelerator bounds</li> <li>30.2 mssm/an: (Co-)annihilation cross sections</li> <li>30.2.1 Annihilation cross sections</li> <li>30.2.1.1 Annihilation cross sections</li> <li>30.2.1.1 Annihilation diagrams</li> <li>30.2.1.3 Neutralino and chargino annihilation</li> <li>30.2.1.4 Squark-squark annihilation</li> </ul> </li> </ul>	79 
<ul> <li>26 Basic principles and common routines <ul> <li>26.1 common/aux: auxiliary routines</li> <li>26.2 common/sm: standard model</li> </ul> </li> <li>27 The empty model </li> <li>28 Generic decaying dark matter </li> <li>29 Generic WIMPs </li> <li>30 The Minimal Supersymmetric Standard Model <ul> <li>30.1 mssm/ac: Accelerator constraints</li> <li>30.1.1 Accelerator bounds</li> <li>30.2 mssm/an: (Co-)annihilation cross sections</li> <li>30.2.1 Annihilation cross sections</li> <li>30.2.1.1 Annihilation cross sections</li> <li>30.2.1.2 Coannihilation diagrams</li> <li>30.2.1.3 Neutralino and chargino annihilation</li> <li>30.2.1.4 Squark-squark annihilation</li> </ul> </li> </ul>	79 
<ul> <li>26 Basic principles and common routines <ul> <li>26.1 common/aux: auxiliary routines</li> <li>26.2 common/sm: standard model</li> </ul> </li> <li>27 The empty model </li> <li>28 Generic decaying dark matter </li> <li>29 Generic WIMPs </li> <li>30 The Minimal Supersymmetric Standard Model <ul> <li>30.1 mssm/ac: Accelerator constraints</li> <li>30.1.1 Accelerator bounds</li> <li>30.2 mssm/an: (Co-)annihilation cross sections</li> <li>30.2.1 Annihilation cross sections</li> <li>30.2.1.1 Annihilation cross sections</li> <li>30.2.1.2 Coannihilation diagrams</li> <li>30.2.1.3 Neutralino and chargino annihilation</li> <li>30.2.1.5 Squark-neutralino annihilation</li> </ul></li></ul>	79 
26 Basic principles and common routines         26.1 common/aux: auxiliary routines         26.2 common/sm: standard model         27 The empty model         28 Generic decaying dark matter         29 Generic WIMPs         30 The Minimal Supersymmetric Standard Model         30.1 mssm/ac: Accelerator constraints         30.2 mssm/an: (Co-)annihilation cross sections         30.2.1 Annihilation cross sections         30.2.1.1 Annihilation cross sections         30.2.1.2 Coannihilation diagrams         30.2.1.3 Neutralino and chargino annihilation         30.2.1.4 Squark-squark annihilation         30.2.1.5 Squark-neutralino annihilation         30.2.1.6 Squark-chargino annihilation	79 
26 Basic principles and common routines         26.1 common/aux: auxiliary routines	79 79 79 79 79 79 79 79 79 80 80 80 80 80 80 80 
26 Basic principles and common routines         26.1 common/aux: auxiliary routines	79 
26 Basic principles and common routines         26.1 common/aux: auxiliary routines	79 79 79 79 79 79 79 81 82 82 82 86 86 86 86 87 91 91 92 92 92

#### CONTENTS

30.3 mssm/an_11: Annihilation cross sections (1-loop)	. 93
30.3.1 Annihilation cross sections at 1-loop – general	. 93
30.4 mssm/an_ib: Internal bremsstrahlung	. 93
30.4.1 Internal Bremsstrahlung (IB) – theory	. 93
30.4.1.1 General considerations	. 94
30.4.1.2 IB from neutralino annihilations	. 94
30.4.1.3 The implementation in $DarkSUSY$	. 95
30.5 mssm/an_sf: Annihilation cross sections (with sfermions) $\ldots \ldots \ldots \ldots \ldots$	. 96
30.5.1 Annihilation cross sections with sfermions – general $\ldots$	. 96
30.6 mssm/an_stu: $t, u$ and $s$ diagrams for $ff$ -annihilation	. 96
30.6.1 Annihilation amplitudes for fermion-fermion annihilation $\ldots \ldots \ldots \ldots$	. 96
30.7 mssm/dd: Direct detection $\ldots \ldots \ldots$	. 97
$30.7.1$ Direct detection – theory $\ldots$	. 97
30.8 mssm/ge: General SUSY model setup: masses, vertices etc	. 99
30.8.1 Supersymmetric model	. 99
30.8.1.1 Parameters	. 99
30.8.1.2 Mass spectrum	. 99
30.8.1.3 Three-particle vertices	. 102
30.8.2 General supersymmetry – routines	. 104
30.9 mssm/ge cmssm: cMSSM interface (Isasugra)	. 105
30.9.1 mSUGRA (ISASUGRA) interface to DarkSUSY	. 105
30.10mssm/ge slha: SUSY Les Houches Accord interface	. 105
30.10.1 SUSY Les Houches Accord	. 105
30.11mssm/ini: Initialization routines	. 105
30.11.1 Initialization routines	. 105
30.12mssm/rd: Relic density	. 106
30.12.1 Relic density of neutralinos	. 106
30.12.2 Internal degrees of freedom	. 106
30.12.2.1 Neutralino-chargino annihilation	. 106
30.12.2.2 Chargino-chargino annihilation	. 107
30.12.2.3 Neutralino-sfermion annihilation	. 108
30.12.2.4 Chargino-sfermion annihilation	. 109
30.12.2.5 Sfermion-sfermion annihilation	. 109
30.12.2.6 Squark-squark annihilation	. 110
30.12.2.7 Sfermion-squark annihilation	. 111
30.12.2.8 Summary of degrees of freedom	. 111
30.13Relic density – more details on routines	. 111
30.13.1 Global parameters	. 111
-	
31 Silveira-Zee (Scalar Singlet)	113
22 Solf Internating Dayle Matter	11/
32 Sen-Interacting Dark Matter	11 <i>1</i>
32.1 vdSIDM/all. Allminiation fournes	. 114
32.2  vu SIDM/ti, $U$ Similar $1$ and $1$ an	. 110 115
32.5 vdSIDM/m. Model Setup	. 110 115
32.4 valiDM/rd. Relic density	. 110 116
32.6 vdSIDM/it. Relic tensity	. 110 116
$52.5$ vasi $D_{\rm M}$ si. Dark matter sen-interactions	. 110
Bibliography	116

Part I Prelude

## Chapter 1 Introduction

DarkSUSY is a set of Fortran routines to allow advanced numerical calculations connected to dark matter physics. In part I of this Manual we explain the basic structure of the code, and provide an introduction on how to get quickly started and use it. Part II is devoted to a description of the DarkSUSY *master library*, which contains all routines that are completely independent of the underlying DM particle physics. In part III, we describe the additional libraries, or *particle modules*, for specific DM models that are currently implemented (including supersymmetric dark matter in the Minimal Supersymmetric Standard Model, the MSSM). The modular structure of the code allows to easily link the master library to any of those particle modules, as well as to add user-defined new modules in a straight-forward way.

The physics involved is covered in the DarkSUSY papers [1, 2]. In this manual we will mainly cover the more techincal aspects of DarkSUSY, i.e. how to call different subroutines, both particle-physics dependent and not, and how to change various switches and options in more advanced routines. We will only briefly review the necessary physics involved when needed and refer the reader to [1, 2] and the original papers behind DarkSUSY (see Section 5) for more details. If you use DarkSUSY please consider the original physics work behind and give proper credit to [2] and the relevant references in Section 5. If you use non-standard options, e.g. a different propagation model for antiprotons, please remember to give proper credit to that model.

### Quick start guide

### 2.1 Installation

To get started, first download DarkSUSY from www.darksusy.org and unpack the tar file. To compile, run the following in the folder where you unpacked it

./configure make

You will then have compiled the main DarkSUSY library, as well as all supplied particle physics modules. To test whether the installation was successful, type

```
cd examples/test ./dstest
```

The program will take up to a minute to run and reports if there are any problems.<sup>\*</sup> At the end, you should get an output of this form

The DarkSUSY test program ran successfully Total time needed (in seconds): 54.85805200000001 Total number of errors in dstest: 0

If you get something else then 0 errors, you should check the output more carefully. The way the test program runs is that for each observable it compares the result with a pre-calculated value. If the difference is larger than 0.3% an error is issued. You normally would not expect larger differences than this due to just numerical errors.

Even if you now have DarkSUSY running, it is more fun to start doing some calculations on your own. Possible next steps are explained in the next Sections.

#### Happy running!

<sup>\*</sup> Strictly speaking, this is only a test of the default particle physics module, mssm. To see what happens behind the scenes, you can run in verbose mode by replacing 'testlevel/2/' with 'testlevel/1/' at the beginning of dstest.f. Then type make and run dstest again. The program is also extensively commented, and can be used to learn about which DarkSUSY routines to call.

#### 2.1.1 Options for install

Even if the above install usually works, sometimes you want to use special options, like a specific compiler. Most options are specified at the time of configure and then propagated through to the DarkSUSY makefiles. Examples of options are

• To specify that you want to compile with gfortran, you can e.g. run configure with the following options (on one line)

```
./configure CC=gcc CFLAGS=-g CXX=g++ CXXFLAGS=-g FC=gfortran
FCFLAGS="-0 -ffixed-line-length-none -fopenmp"
```

For your convenience, this particular choice is a available as a script **conf.gfortran** that can be invoked instead of the string above.

#### 2.1.2 System requirements

DarkSUSY should run on most Linux/Unix systems including Mac OS X. You need to have a Fortran, C and C++ compiler available and the typical developer tools (like make and ranlib). You also need to have perl installed. If you are creating new particle physics modules and want to use the tools available to automatically create makefiles for you, you also need to have autoconf installed.

#### 2.2 Example programs

DarkSUSY is primarily a library that is intended to be used with your own main programs. However, to get you started, we supply a few sample programs in the /examples directory. These can be used as they are, but they are also extensively commented to help you understand which routines you are supposed to call for the most typical calculations. The most instructive general-purpose programs in /examples, apart from the already mentioned /test/dstest, are

dsmain\_wimp.F An example main program to calculate various DM observables. It can be the starting point if you want to write your own programs (see below). Note that you can run *the same code* with different particle modules that provide a WIMP DM candidate (the default is the mssm module); simply do

./> make -B dsmain\_wimp DS\_MODULE=(MY\_MODULE)

- and then ./dsmain\_wimp in the /examples directory. Choosing  $\langle MY\_MODULE \rangle$ =generic\_wimp, e.g., will produce results for a generic WIMP DM candidate (see Chapter 29 for details on the implementation).
- dsmain\_decay.f A main program that calculates the same observables, where relevant, as dsmain\_wimp.F – but for a generic decaying DM candidate (see Chapter 28 for details on the implementation) rather than a WIMP.

#### 2.2.1 Auxiliary example programs

In the folder examples/aux/ we list a set of auxiliary example programs that illustrate more specific usage of DarkSUSY. These example programs are typically set up for specific particle physics modules (often generic\_wimp) but, as explained below in more detail, it is straightforward to use them for other particle physics modules as well. Some of the currently available auxiliary example programs are

**flxconv.f** This program can be used to convert between different fluxes from the Sun and the Earth. It can be use to convert a limit on a muon flux to a limit on the scattering cross section, or to a limit on the annihilation rate (or vice versa).

- **DMhalo\_predef.f** This program illustrates how to use the default version of dsdmsdriver (provided with the DarkSUSY release) to load additional profiles into the currently active halo database by using its pre-defined halo parameterizations.
- **DMhalo\_table.f** This program demonstrates how to load a halo profile from a table in an accompanying data file.
- **DMhalo\_new.f** This program demonstrates how to correctly extend dsdmsdriver when adding a new profile parameterization in order to consistently make it available to all DarkSUSY routines that rely on the DM density (in this concrete example, we add the spherical Zhao profile [3], aka  $\alpha\beta\gamma$  profile).
- DMhalo\_bypass.f Here, we demonstrate a work-around of completely bypassing the default dsdmsdriver setup when switching to a user-provided new DM density profile. While easier to implement than DMhalo\_new.f, such an approach has the significant drawback that the advanced DarkSUSY system of automatic tabulation of quantities related to DM rates cannot easily be exploited.
- **generic\_wimp\_oh2.f** This program calculates the annihilation cross section needed to produce the correct relic density (as measured by Planck within errors). It does this for a range of WIMP masses so that you can plot e.g. the required annihilation cross section versus mass. It also let's you incorporate threshold effects (either as a hard cut, default), or with a more sophisticated sub-threshold treatment, that also illustrates the use of replaceable functions.
- **ScalarSinglet\_RD.f** This program calculates the couplings required to get the correct relic density in the Silveira-Zee (scalar singlet) model.
- ucmh test.f This program is an example of how the ultra-compact mini halo routines can be used.
- **wimpyields.f** This is a simple program that sets up a generic WIMP with a given mass that annihilates to a given channel and then calculates the yields of different particles from the hadronization/decay of the annihilation products.
- **caprates.f** This is a simple program that scans a range of WIMP masses and calculates the capture rates in the Sun via spin-independent and spin-dependent scattering.
- **caprates\_ff.f** This program is similar to **caprates.f**, but a bit more advanced as it performs the capture rate calculation with the most complete numerical setup and also calculates the capture rate on individual elements in the Sun.

#### 2.2.2 Making your own example programs

The simplest way to create your own example program is to copy one of the Fortran example files in examples/aux/ to a directory of your choice, then change the name of that file and modify it. You also need to copy the makefile from examples/aux/ to the same directory, and make sure that you update the name of the file that you just changed.<sup>†</sup> Running 'make' then compiles your own new example program without touching the release version of the DarkSUSY code. Always keeping your own code, or your modifications of DarkSUSY, separate like this is the recommended way to proceed because it facilitates debugging and minimizes the risk of introducing errors. Don't modify any of the DarkSUSY makefiles directly as they are overwritten every time you run configure.

If you want to *compile with a different particle physics modules*, you will in general need to do go through three simple steps:

 $<sup>^{\</sup>dagger}$ If you want, you can delete all the blocks for the other (not needed) example programs, in order to have the makefile look clearer and easier to understand.

- 1. change the corresponding block in the makefile, i.e. simply set the variable DS\_MODULE to a different value (you may also have to add additional libraries, e.g. lisajet for the mssm module)
- 2. update the model setup routines to the new particle module (see the various example programs; for more details, have a look at the respective Section of this manual and the header of setup routines starting with dsgivemodel).
- 3. make sure that your main program only calls routines that are supported for the new particle module. (If it doesn't, it will not compile, stating the functions that are not supported)

An explicit demonstration of how all these three steps are taken care of in one single example is provided in dsmain\_wimp.F. Note that the pre-compiler directives in that example are only necessary if you want to be able to compile the *same* code with two (or more) different particle modules.

### 2.3 Modifying individual subroutines or functions

If you want to modify some existing DarkSUSY function or subroutine, *don't do it*, instead add your own routine as a replaceable function (see also Section 3.6), by running the script scr/make\_replaceable.pl on the routine you wish to have a user-replaceable version of. You will then find that version in the corresponding user\_replaceables folder, where you can edit it to your liking. Following these steps, it is guaranteed that the newly created user-replaceable function is properly included in the library where the original DS function used to be, with all makefiles being automagically updated.

An alternative – and often even simpler – way of using user-replaceable functions is to leave the DS libraries untouched, and to instead link to the user-supplied function only when making the main program (this option is indicated in the left-most part of Fig. 3.1). For an explicit example, see (the makefile for) generic\_wimp\_oh2.f.

# Chapter 3 Guiding principles of the code

DarkSUSY (as of version 6) has a new structure compared to earlier versions of the code. The most striking difference is that we have split the particle physics model dependent parts from the rest of the code. This means in practice that we have separated DarkSUSY into one set of routines, ds\_core, which contains no reference to any specific particle model, as well as distinct sets of routines for each implemented model of particle physics. For supersymmetry, for example, all routines that require model-dependent information now reside in the mssm module. The advantage with this setup is that ds\_core and the particle physics modules may be put in separate FORTRAN libraries, which implies that DarkSUSY can now have several particle physics modules side by side. The user then simply decides at the linking stage, i.e. when making the main program, which particle physics module to include.

For this to work, the main library communicates with the particle physics modules via so-called *interface* functions (or subroutines), with pre-defined signatures and functionalities. Note that a particle physics module does not have to provide all of these predefined functions: which of them are required is ultimately determined only when the user links the main program to the (ds\_core and particle module) libraries. Assume for example that the main program wants to calculate the gamma-ray flux from DM (a functionality provided by ds\_core). This is only possible if the particle module provides an interface function for the local cosmic ray source function; if it does not, the main program will not compile and a warning is issued that points to the missing interface function. If, on the other hand, interface functions required by direct detection routines would be missing in this example, this would not create any problems at either runtime or the compile stage.

In addition, we have added the concept of *replaceable functions*, which allows users to replace essentially any function in DarkSUSY with a user-supplied version. DarkSUSY ships with dedicated tools to assist you setting up both replaceable functions and new particle physics modules. Fig. 3.1 illustrates these concepts by showing how a typical program would use DarkSUSY; below, we describe each of them in more detail.

### 3.1 The main DarkSUSY library ds core

As introduced above, the main library is in some sense the heart of the new DarkSUSY, offering all the functionality that a user typically would be interested in without having explicitly to refer to specific characteristics of a given particle physics model (after initialization of such a model). The main library thus contains routines for, e.g., cosmic ray propagation, solar models, capture rates for the Sun/Earth, a Boltzmann solver for the relic density calculation, yield tables from annihilation/decay etc. None of the routines in the main library contains any information about the particle physics module. Instead any information needed is obtained by calling a required function that resides in the particle physics module which is linked to (see below).



Figure 3.1: Conceptual illustration of how to use DarkSUSY. The main program links to both the main library, ds\_core, and to *one* of the available particle physics modules. User-replaceable functions are optional and may be linked to directly from the main program, or indirectly by including them in the various libraries. See examples/dsmain\_wimp.F for an example of a main program that demonstrates typical usage of DarkSUSY for different particle physics modules.

The source code for all functions and subroutines in the main library can be found in the src/directory of the DarkSUSY installation folder, with subdirectory names indicating subject areas as summarized in Table 3.1.

### 3.2 Particle physics modules

The particle physics modules contain the parts of the code that depend on the respective particle physics model. Examples are cross section calculations, yield calculations etc. The routines in the particle physics module have access to all routines in ds\_core, whereas the reverse is in general not true (with the exception of a very limited set of interface functions that each particle module provides).

DarkSUSY 5 and earlier was primarily used for supersymmetric, and neutralino DM, and those parts of the code now reside in the MSSM module mssm. However, many people used DarkSUSY even before for e.g. a generic WIMP setup, which was doable for parts of the code, but not all of it. We now provide a generic WIMP module generic\_wimp that can be used for these kinds of calculations in a much more general way. In a similar spirit, we also provide a module generic\_decayingDM for phenomenological studies of decaying DM scenarios. As an example for an actual particle physics model other than supersymmetry, DarkSUSY furthermore includes a module silveira\_zee which implements the DM candidate originally proposed by Silveira and Zee [4] and which now is often referred to as Scalar Singlet DM [5, 6, 7]. We also include an empty model, empty, which is of course not doing any real calculations, but contains (empty versions of) all interface functions that the core library is aware of – which is very useful for debugging and testing purposes. Designing new particle modules is a straight-forward exercise, see below, and it is generally a good idea to start with the most similar module that is already available.

Subdirectory		
name	Description	
an_yield	Annihilation yields in the halo – yields from simulations	
aux	General routines	
$aux\_xcernlib$	CERN routines needed by DarkSUSY	
aux_xdiag	Diagonalization routines	
aux_xquadpack	CMLIB routines needed by DarkSUSY	
cr_aux	Cosmic rays – general	
cr_axi	Cosmic rays – diffusion routines for axisymmetric distributions	
cr_gamma	Cosmic rays – Gamma fluxes	
cr_nu	Cosmic rays – Neutrino fluxes	
cr_ps	Cosmic rays – point sources	
dd	Direct detection	
dmd_astro	Astrophysical source functions	
dmd_aux	Auxiliary functions for dark matter distribution routines	
$dmd_mod$	Dark matter distributions	
dmd_vel	Dark matter velocity distributions	
ini	Initialization routines	
kd	Kinetic decoupling	
rd	Relic density	
se_aux	Auxiliary routines for WIMP annihilation in the Sun/Earth	
se_mod	Sun and Earth models	
se_nu	Capture and annihilation in the Sun/Earth	
se_yield	Yields from annihilation in the Sun/Earth	
si	Self-interactions	
ucmh	Ultra-compact mini-halos	

Table 3.1: Organization of the main library  $ds\_core$ : all functions and subroutines reside in the src/ folder of the DarkSUSY installation, with the names of the subdirectories indicating the subject area. (Note: this table is automatically generated from the actual directory structure in src/.)

A list of the currently available particle physics modules is given in table 3.2, and the source code for the particle modules can be found in the src\_models/ directory of the DarkSUSY installation folder, each of the subdirectories (e.g. src\_models/mssm/, src\_models/generic\_wimp/) typically reflecting a (sub)subdirectory structure analogous to what is shown in Table 3.1 for the core library. Many dark matter models, furthermore, constitute only relatively simple extensions to the standard model, inheriting most of its structure. For convenience, we therefore also provide various auxiliary routines, in src\_models/common/sm, that each particle module automatically has access to, and which return basic standard model quantities like, e.g., the masses of standard model particles and their running (so additional BSM effects have to be implemented in the respective particle module).

#### 3.2.1 Using a particle physics module

How to use a particle physics module obviously depends on how it is implemented, and in principle there are no formal requirements on how this should be done – as long as the provided interface functions are correctly set up, see Section 3.4 further down. The one subroutine that *is* required to exist, however, is

Module	Short	
No.	name	Description
1	common	Basic principles and common routines
2	empty	The empty model
3	generic_decayingDM	Generic decaying dark matter
4	generic_wimp	Generic WIMPs
5	mssm	The Minimal Supersymmetric Standard Model
6	silveira_zee	Silveira-Zee (Scalar Singlet)
7	vdSIDM	Self-Interacting Dark Matter

<b>D D</b>	1		
Particia	nnveice	mod	1100
I al uluic	DILVSIUS	mou	uico
	1 0		

Table 3.2: List of particle physics modules currently available in src\_models. (Note: this table is automatically generated from the actual content of src models.)

#### dsinit\_module

This is called directly from dsinit, and must be the first call to the particle module as it initializes all general settings and relevant common block variables.

While there is no required structure otherwise, there is a typical workflow associated to using a particle physics module in a main program. First, one needs to initialize a given model by specifying its model parameters. This is done by routines like

#### dsgivemodel...

A call to dsgivemodel\_decayingDM, e.g., allows to enter the defining parameters for a DM model in the generic\_decayingDM module, while dsgivemodel\_25 allows to enter the parameters for pMSSM model with 25 parameters in the mssm module. The next step is then typically a call to

#### dsmodelsetup

in order to transfer the model parameters to common blocks and calculate basic quantities like masses and couplings. Once a model is set up like this, a main program can use the full functionality of DarkSUSY supported by the respective module.

#### 3.2.2 Adding a new particle physics module

To create a new particle physics module, the easiest way is to start from an existing one as a template and create a new one from that one. To help you in this process we provide a script scr/make\_module.pl that takes two arguments, the module you want to start from and the new one you want to create (for further instructions, just call the script without arguments). It will then copy the module to a new one, change its name throughout the module and make sure that it is compiled by the makefiles and included properly when requested by the main programs. If you specify the option -i only interface functions will be copied (which creates a cleaner starting point, but also will most likely not compile without modifications). When creating a new module this way, the best is to copy from a module that is as similar as possible to your new model. If you want a clean setup, you can always copy from the empty module. If you want something more phenomenological that has a basic framework for calculating observables, starting from the generic\_decayingDM might be a good idea. A general advice is to view the modules we provide as a starting point as inspiration for your new modules.

Even though a particle physics module does not need to include all interface functions (which ones are needed only depends on the observables you try to calculate in your main programs), it needs to provide an initialization routine dsinit\_module.f. This routine should set a global variable

moduletag to the name of the module so that routines that need to check if the correct module is loaded can do so. When using the script scr/make\_module.pl this routine is always created and moduletag set as it should.

Note that the script scr/make\_module.pl only provides the framework in the configure script and makefiles (or rather makefile.in's) to make sure your module is compiled. It does *not* create a main program that uses your module, that is up to you. A good starting point for that can either be e.g. dsmain\_wimp.F in examples that already contain different blocks for different modules (controlled by pre-compiler directives, see the code for more details). Another option is to use any of the example programs in examples/aux as a starting point.

To make upgrading to new DarkSUSY versions as smooth as possible, we advice to create your own folder with your main programs, using e.g. the makefile in examples/aux as a starting point. Don't modify any of the DarkSUSY makefiles directly as they are overwritten every time you run configure, see Section 2.2.2 for more details.

Note that for the scr/make\_module.pl script to work you need to have autoconf installed as the script adds your new module to configure.ac and runs autoconf to create a new configure script.

#### 3.3 Halo models

Several routines in the core library need to know which DM halo should be adopted for the calculations. With this DarkSUSY version, we introduce a new and flexible scheme that avoids pre-defined hardcoded functions to describe the DM density profiles, and allows to consistently define different DM targets at the same time. For convenience, we still provide several pre-defined options for such halo parameterizations, and the user can choose between, e.g., the Einasto [8] and the Navarrow-Frenk-White profile [9], or read in any tabulated axi-symmetric (or spherically symmetric) profile. On a *technical* level, the halo models are handled by the dsdmsdriver routine which contains a database of which halo profiles the user has set up, and consistently passes this information to all parts in the code where it is needed.\*

We provide detailed hands-on examples on how to use this system by a set of example programs DMhalo\_\*.f, see Section 2.2.1. For further details, we refer to Section 16.1 of this manual.

### 3.4 Interface functions

Interface functions are functions that routines in ds\_core might need to call, and which therefore every particle physics module should contain if that particular observable is requested. Examples of interface functions include dsddsigma that returns the equivalent DM nucleus cross section, dscr-source that returns the source term for DM-induced cosmic rays (relevant for indirect detection), dsanxw that returns the invariant annihilation rate (in the case of WIMPs), etc. All interface functions are found by looking in the empty module, and always contain the keyword 'interface' in the function or subroutine header.

If a particle physics module contains the full set of interface functions, all routines in the main DarkSUSY code should work. However, this is not needed to set up a paticle physics module. E.g. a generic WIMP model does not need to have decay rates set up. If a routine in the main DarkSUSY routines is called that rely on this interface being present, an error will be thrown when trying to compile your code. This of course applies to all interface functions, not just unphysical ones. E.g.

<sup>\*</sup> From a technical point of view, it is actually not dsdmsdriver itself which acts as an interface to the rest of the code, but the set of wrapper routines collected in src/dmd\_astro (which all call dsdmsdriver in a specific way). Only those routines are called by cosmic-ray flux routines and other functions in src, never dsdmsdriver directly. The routines in src/dmd\_astro therefore provide examples of functions that *cannot be replaced* individually in a consistent way, but only as a whole set (along with dsdmsdriver, in case the user wants to change the structure of the driver itself).

if you are only interested in the relic density for a new particle physics module, you do not need to set up scattering rates, cosmic ray source functions, etc.

In Tab. 3.3, we provide the complete list of interface functions currently implemented in Dark-SUSY, along with a brief description. For more details, consult the headers of these files.

	Appear in		
Routine	modules	Used by	Description
dsacbnd	2, 5		Check collider bounds
dsanwx	2,  4,  5,  6,  7	rd	Self-annihilation invariant rate
dscrsource_line	2, 3, 4, 5, 6, 7	cr_axi	Source term for monochromatic contributions from dark matter
dscrsource	2, 3, 4, 5, 6, 7	cr_axi, cr_gamma, cr_nu, ucmh	Source term for dark matter induced cosmic rays
dsddgpgn	2, 4, 5, 6, 7	se_nu	Four fermion couplings (the use of this function will be phased out in favour of dsddsigma).
dsddsigma	2, 4, 5, 6, 7	dd	UNpolarized *equivalent* WIMP nucleus cross section including form factors
dsinit_module	2,  3,  4,  5,  6,  7	ini	Intialzation of module
dskdm2simp	2, 4, 5, 6, 7	kd	Scattering amplitude squared for zero momentum transfer
dskdm2	2, 4, 5, 6, 7	kd	Full scattering amplitude squared at zero momentum transfer
dskdparticles	2, 4, 5, 6, 7	kd	Initalization of kinetic decoupling for module
dsmodelsetup	2, 3, 4, 5, 6, 7		Sets up a new particle physics model in module
dsmwimp	2,  3,  4,  5,  6,  7	cr_axi, kd	WIMP mass
dsrdparticles	2, 4, 5, 6, 7	rd	Particles included in relic density calcu- lation (coannihilations, resonances and thresholds)
dsseyield	2, 4, 5, 6, 7	se_nu	Yields from annihilation in the ${\rm Sun}/{\rm Earth}$
dssigmav0tot	2,4,5,6,7	cr_axi, se_nu	Total annihilation cross section at $v = 0$
dssisigtm	2, 7	si	momentum-transfer cross section
dswimpspin	2,  3,  4,  5,  6	se_nu	WIMP spin

Table 3.3: Table of interface functions, which modules that contain them (with numbering from table 3.2, where they are used and a short description of them. (Note: this table is automatically generated from scanning through the particle physics modules in src\_models. The description is taken from the empty module routine headers.)

### 3.5 Commonly used functions

Routines that we believe are particularly useful for most users to call, we denote as *commonly used functions*. A list of the commonly used functions in the main DarkSUSY library in src is shown in

Table 3.4. The routines are described in more detail in their corresponding chapters in part II of this manual, as indicated in the table 3.4.

Routine	Description	Chapter
dsanyield_sim	Simulated particle yields	6
dsdbdphidtaxi	Local galactic differential antideuteron flux from dark matter	10
dsepdphidpaxi	Local galactic differential positron flux from dark matter	10
dspbdphidtaxi	Local galactic differential antiproton flux from dark matter	10
dscrgaflux_line_v0ann	Flux of monoenergetic gamma-rays from annihilation in the halo, in the limit of zero relative velocity	11
dscrgaflux_v0ann	Flux of gamma-rays from annihilation in the halo, in the limit of zero relative velocity	11
dsgafluxsph	gamma-rays from decay/annihilation from halo specified by halo label	11
dscrmuflux_v0ann	Flux of neutrino-induced muons from WIMP annihila- tion in the halo	12
dsdddrde	Differential WIMP-nucleus recoil rates	14
dsddhelp	help with options for scattering cross sections	14
dsddset	Set parameters for scattering cross sections	14
dsddsigmanucleon	Calculate nuclear cross sections	14
dsdfactor	D-factor (l.o.s. integral for decaying DM)	??
dsjfactor	J-factor (l.o.s. integral for annihilating DM)	??
dsdmdselect halomodel	select between halo models in the halo repository	16
dsdmdset_halomodel	add an existing halo model to halo repository	16
dsinit	Initialize DarkSUSY (should always be called)	18
dskdmcut	Cutoff mass in linear power spectrum	19
dskdtkd	Kinetic decoupling temperature	19
dsrdomega	Calculate the relic density $\Omega h^2$ of a dark matter particle	20
dssenu_rates	Rates of neutrinos and neutrino-induced leptons and	23
_	hadronic showers from WIMP annihilations in the $Sun/Earth$	
dssisigtmav	velocity-averaged momentum-transfer cross section	25

Table 3.4: List of commonly used functions and subroutines. (Note: this table is automatically generated from the routines in **src** that have the commonly used tag in their header.)

### 3.6 Replaceable functions

The concept of replaceable functions introduces the possibility to replace a DarkSUSY routine with one of your own. The way it works is that the user-provided function will be linked when you make your main program instead of the DarkSUSY default one. If, for example, you want to replace the yields from a typical final state of DM annihilation or decay (like  $\bar{b}b$ ) with a new function of your own – e.g. because you are interested in comparing the tabulated Pythia [10] yields with those provided by PPPC [11] – you create your own version of the routine dsanyield\_sim and let DarkSUSY use this one instead. To help you with this setup, we provide tools that can create (or delete) a replaceable function from any DarkSUSY function and set up the makefiles to use this user-supplied function instead. We also provide a simply way of managing large 'libraries' of user-supplied functions, via a list imported by the makefiles, where the user can determine on the fly which user-replaceable

functions should be included and which ones should not. Note that both routines in ds\_core and in any of the particle physics modules can be replaced in this way, including interface functions. There is also a possibility for the particle physics modules to replace a function in the core library. This is not used often, but is used e.g. for the function dsrdxi.f (which gives a possibility to have different dark matter and background temperatures in the early Universe).

There is also a simple way to bypass this system – indicated in the left-most part of Fig. 3.1 – by directly linking to the user-supplied function only when making the main program. For an explicit example, see (the makefile for) generic\_wimp\_oh2.f.

# Comparison to previous DarkSUSY versions

For those DarkSUSY users who are familiar with previous version of the code, the main structural difference introduced with DarkSUSY 6.0 is a highly modular setup that allows to fully disentangle the astrophysics-related parts from those that rely on a specific particle physics model (see Chapter 3 for a detailed description). There are also many new or significantly improved physics capabilities introduced after version 5 and 4 [1] – including electroweak and strong corrections to DM annihilation, improved routines to solve the Boltzmann equation for chemical and kinetic decoupling, and a new framework to handle the propagation of cosmic rays. For a detailed list of those new features, we refer to the publication describing this release of the code [2].

One technical aspect that has changed are the particle codes (k-variables), which are now treated as (module-)internal codes. They can be used by the particle physics module if the module so wishes, but the interface functions and routines in ds\_core instead use PDG codes when referring to particles.

In the course of re-organizing the code, it also became necessary to re-name some of the basic functions and subroutines that existed in earlier versions and which users may have become familiar with. For convenience, we therefore list below, in Table 4.1, the most commonly used functionalities in version 4 and 5 that have changed in name or usage with the present version of the code.

routine name up to $DarkSUSY\ 5$	new routine name	comments
dshmset	dsdmdset_halomodel	Setting up and refering to DM ha- los has fundamentally changed. See Section 3.3 for an overview and all Chapters with names containing src_dmd in Part II of the manual for more details.
dshmj	dsjfactor	Line-of-sight integrals now take a halo label as input, and can be computed for various objects at the same time.
dssusy	[dsmodelsetup]	Setting up a model (calculating the mass spectrum, relevant 3-particle vertices etc.) now depends on the particle module implementation.

dsmhtkd dsmhmcut	dskdtkd dskdmcut	The 'microhalo' routines are now more properly referred to as 'kinetic decoupling' routines.
dshmrescale_rho		A mismatch between local halo den- sity and DM abundance for a given DarkSUSY module is no longer hid- den as a rescaling factor in a com- mon block. Instead, such factors typically enter as explicit parame- ters in direct and indirect detection routines.
dshaloyield	[dsanyield]	total cosmic ray yield from $neutralino$ annihilation
dshayield	dsanyield_sim	simulated cosmic ray yields from individual annihilation/decay chan- nels to SM particles. Note that internal channel codes are now re- placed with PDG codes of the final state particles as input.
dshrgacontdiff	dsgafluxsph	Gamma-ray flux routines now work seamlessly together with both the new halo setup and the modular par- ticle physics structure.
dshrgaline	dscrgaflux_line_v0ann dscrgaflux_dec	Gamma-ray line routines now return both number, energies and widths of all such signals that are present in the current particle setup.
dshrpbardiff dshrdbardiff dsepdiff	dspbdphidtaxi dsdbdphidtaxi dsepdphidpaxi	Cosmic-ray propagation routines have been re-written from scratch. They are now much more flexible and can be used for any axisymmet- ric halo/diffusion model.
dsntrates	dssenu_rates	Neutrino rates from inside the sun or earth
dshrmuhalo	dscrmuflux_v0ann dscrmuflux_v0ann	Neutrino-induced muon flux from the halo (for annihilating and decay- ing DM, respectively).

Table 4.1: 'Translation table' for how the most commonly used functionalities in DarkSUSY version 5 and earlier have changed with the present version of the code. Routines in parentheses, e.g. [dsanyield], are no longer part of the DarkSUSY main library and only provided by specific particle physics modules. For more detailed descriptions of the new routines and functions, see the headers of the respective files.

### **Original articles**

A large part of the routines in the present DarkSUSY version have been implemented in the context of original research work. Therefore, when using those routines, please give proper credit not only to the main DarkSUSY paper [1, 2] but also to the relevant articles in the following list:

- Relic density P. Gondolo and G. Gelmini, Nucl. Phys. B360 (1991) 145 [12]; J. Edsjö and P. Gondolo, Phys. Rev. D56 (1997) 1879 [13]; J. Edsjö, M. Schelke, P. Ullio and P. Gondolo, JCAP 04 (2003) 001 [14].
- Kinetic decoupling and microhalos T. Bringmann, NJP 11 (2009) 10527 [15].
- Continuous gamma-rays L. Bergström, J. Edsjö and P. Ullio, Phys. Rev. D58 (1998) 083507 [16].
- Neutrino telescopes L. Bergström, J. Edsjö and P. Gondolo, Phys. Rev. D58 (1998) 103519 [17].
- Positrons E.A. Baltz and J. Edsjö, Phys. Rev. D59 (1999) 023511 [18].
- Antiprotons L. Bergström, J. Edsjö and P. Ullio, ApJ 526 (1999) 215 [19].
- QCD corrections to DM annihilation T. Bringmann, A. J. Galea, P. Walia, Phys.Rev. D93 (2016) 043529 [20].
- General MSSM, direct detection L. Bergström and P. Gondolo, Astrop. Phys. 5 (1996) 263 [21].
- Gamma lines from supersymmetric DM L. Bergström and P. Ullio, Nucl. Phys. B504 (1997) 27 [22]; P. Ullio and L. Bergström, Phys. Rev. D57 (1998) 1962 [23].
- Internal bremsstrahlung in the MSSM ( $\gamma$  and  $e^+$ ) T. Bringmann, L. Bergström and J. Edsjö, JHEP 0801 (2008) 049 [24]; L. Bergström, T. Bringmann and J. Edsjö, Phys.Rev. D78 (2008) 103520 [25].
- MSSM Electroweak corrections to indirect detection yields T. Bringmann and F. Calore, Phys.Rev.Lett. 112 (2014) 071301 [26]; T. Bringmann, F. Calore, A. J. Galea and M. Garny, JHEP 1709 (2016) 041 [27].

### Part II

# Main DarkSUSY routines in src/ $\,$

### an\_yield: Annihilation yields in the halo – yields from simulations

### 6.1 Annihilation in the halo, yields – theory

Here we calculate yields of different particles from annihilation of dark matter particles in the halo.

#### 6.1.1 Monte Carlo simulations

We need to evaluate the yield of different particles per WIMP annihilation. The hadronization and/or decay of the annihilation products are simulated with PYTHIA [10] 6.426. The simulations are done for a set of 18 WIMP masses,  $m_{\chi} = 10, 25, 50, 80.3, 91.2, 100, 150, 176, 200, 250, 350, 500, 750, 1000, 1500, 2000, 3000 and 5000 GeV. We tabulate the yields and then interpolate these tables in DarkSUSY.$ 

The simulations are here simpler than those for annihilation in the Sun/Earth since we don't have a surrounding medium that can stop the annihilation products. We here simulate for 8 'fundamental' annihilation channels  $c\bar{c}$ ,  $b\bar{b}$ ,  $t\bar{t}$ ,  $\tau^+\tau^-$ ,  $W^+W^-$ ,  $Z^0Z^0$ , gg and  $\mu^+\mu^-$ . Compared to the simulations in the Earth and the Sun, we now let pions and kaons decay and we also let antineutrons decay to antiprotons. For each mass we simulate  $2.5 \times 10^6$  annihilations and tabulate the yield of antiprotons, positrons, gamma rays (not the gamma lines), muon neutrinos and neutrino-to-muon conversion rates and the neutrino-induced muon yield, where in the last two cases the neutrino-nucleon interactions has been simulated with PYTHIA as outlined in section 24.1.1

With these simulations, we can calculate the yield for any of these particles for a given particle physics model. In **src**, we only include the channels that are actually simulated. In the different particle physics modules in **src\_model**, the summation over all possible final states and possibly more complex channels, like scalars decaying to other particles is done.

Note that simulations are typically done without specifying a particular polarization state of the final state particles. This is however not entirely correct as the possible polarization states will depend on the particle physics model we have.

Even if the simulations are not performed in this more general way yet, we have set up the structure here to eventually provide the yields in this form. In particular the most general routine to calculate the yields from any of the simulated channels is **dsanyield\_sim\_ls**. Apart from the mass, energy and yield type, this routine also takes are arguments the PDG codes of the final state particles and the polarization state of the final state. In particular, you are required to provide the

quantum numbers

- j = the quantum number for the total angular momentum
- P = the parity of the final state
- l = the quantum number for the orbital angular momentum in the final state
- s = the quantum number for the total spin in the final state

We want to move in a direction where this routine is the one that should be used. While getting there, we also provide a simpler routine which just gives the polarization state in terms of helcity and polarization, **dsanyield\_sim**. This routine takes the PDG code of the final state and the polarization as

- Left-handed or right-handed polarization for fermions.
- Longitudinal or transverse polarization for vector bosons.

This routine for simplicity assumes that the final state particles have the same polarization state.

### aux: General routines

### 7.1 General routines

In **aux/**, we collect routines that are of general interest to many other routines in DarkSUSY. E.g., we have routines to find elements in arrays (used for interpolation), Bessel functions, error functions, spline routines, etc.

### aux\_xdiag: Diagonalization routines

### 8.1 XDIAG

This folder contains a set of routines to perform numerical diagonalization of complex matrices. Even though analytical routines can be used for up to  $5 \times 5$  matrics, they tend to be numerically unstable for matrices occuring in some particle physics models (for example supersymmetry). Hence, we use numerical routines to perform the diagonalization instead.

### cr\_aux: Cosmic rays – general

### 9.1 Cosmic Rays – auxiliary routines

This folder contains auxiliary functions needed by the cosmic ray routines. In particular, it contains different versions of simple integration routines and a set of routines to handle the correct setting and interpreting of labels for the tabulation of confinement times and Green's functions needed for the calculation of cosmic ray fluxes.

### cr axi:

# Cosmic rays – diffusion routines for axisymmetric distributions

#### 10.1 Cosmic ray propagation in axially symmetric halos

There is a clean asymmetry between particles and antiparticles in the standard cosmic ray picture: The bulk of cosmic rays – protons, nuclei and electrons – are mainly "primary" species, i.e. particles accelerated in astrophysical sources and then copiously injected in the interstellar medium; "secondary" components, including antimatter, are instead produced in the interaction of primaries with the interstellar medium during the propagation. It follows that there is a pronounced deficit of antimatter compared to matter in the locally measured cosmic ray flux (about 1 antiproton in  $10^4$  protons). When considering instead a source term due to DM annihilations or decays, a significant particle-antiparticle asymmetry is in general not expected, and antiprotons, positrons and antideuterons turn out to be competitive indirect DM probes.

Charged particles propagate diffusively through the regular and turbulent components of Galactic magnetic fields. This makes it more involved for local measurements to track spectral and morphological imprints of DM sources than, e.g., for the gamma-ray and neutrino channels (though searches for spectral features in CR positron fluxes still lead to very competitive limits [?]). In fact the transport of cosmic rays in the Galaxy is still a debated subject: Most often one refers to the quasi-linear theory picture (with magnetic inhomogeneities as a perturbation compared to regular field lines) in which propagation can be described in terms of a (set of) equation(s) linear in the density of a given species, containing terms describing diffusion in real space, diffusion in momentum space (the so-called reacceleration), convective effects due to Galactic winds, energy or fragmentation losses and primary and secondary sources (see, e.g., Ref. [?] for a review). Dedicated codes have been developed to solve numerically this transport equation, including GALPROP [?], DRAGON [?] and PICARD [?].

Here we follow instead a semi-analytical approach, analogous to that developed for the USINE code [?]. In particular, we model the propagation of antiprotons and antideuterons by considering the steady state equation [19]

$$\frac{\partial N}{\partial t} = 0 = \nabla \cdot (D \nabla N) - \nabla \cdot (\vec{u} N) - \frac{N}{\tau_N} + Q.$$
(10.1)

We solve it for situations where i) the diffusion coefficient D can have an arbitrary dependence on the particle rigidity but can at most take two different values in the Galactic disc and in the diffusive halo, ii) the convective velocity  $\vec{u}$  has a given fixed modulus and is oriented outwards and perpendicular to the disc, *iii*) the loss term due to inelastic collisions has an interaction time  $\tau_N$  which is energy dependent but spatially constant in the disc and going to infinity in the halo (corresponding to a constant target gas density in the disc and no gas in the halo), *iv*) the DM source Q is spatially axisymmetric and has a generic energy dendence. Under these approximations and assuming, as is usually done, that the propagation volume is a cylinder centred at the disc and that particles can freely escape at the boundaries of the diffusion region, Eq. (10.1) can be solved analytically by expanding N in a Fourier-Bessel series; the computation of the flux involves, at each energy, a sum over the series of zeros of a Bessel function of first kind and order zero, and a volume integral of the spatially dependent part in the axisymmetric source term Q (basically the DM density  $\rho_{\chi}$  for decaying DM and its square for pair annihilating DM) times a weight function depending on the given zero in the series (see [19] for further details).

Since the path lengths for antiprotons and antideuterons are rather large, of the order of a few kpc, taking average values for parameters in the transport equation rather than the more realistic modelling that can be implemented in full numerical solutions, has no large impact in case of extended and rather smooth sources such as for DM. Eq. (10.1) neglects reacceleration effects, which may in general be relevant at low energies (rigidities below a few GV); however even this does not have a large impact in case of the species at hand, see, e.g., [?] for a comparison of results with numerical and semi-analytical solutions for cosmic-ray antiprotons. The power of our semianalytic approach is that one can store values of the solution of the transport equation obtained by assuming a given mono-energetic source – provided by the functions dspbtdaxi and dsdbtdaxi for, respectively, antiprotons and antideuterons – and then apply these as weights to any particle source term  $\mathcal{S}_n(E_f)$  as introduced above. For antiprotons and antideuterons, this latter step is done in the functions that compute the local galactic differential fluxes from DM annihilation and decay, dspbdphidtaxi and dsdbdphidtaxi, respectively. Note that while the outputs of dspbtdaxi and dsdbtdaxi are labelled "confinement time" in the code, since they do have a dimension of time and scale the dependence between source and flux, one cannot trade them for what is usually intended as confinement time for standard cosmic ray components, given that the morphology of the DM source is totally different from supernova remnant distributions usually implemented for describing ordinary primary components.

The structure we implemented gives a particularly clear advantage when the code is used to scan over many particle physics DM models, but only over a limited number of propagation parameters and DM density profiles. For such an application, it is useful to tabulate the 'confinement times' (returned by dspbtdaxi and dsdbtdaxi) over a predefined range of energy; this is done in the functions dspbtdaxitab and dsdbtdaxitab when calling the flux routines with an appropriate option (and only in case the DM halo profile currently active is within the halo profile database). Such tabulations can be saved and re-loaded for later use; here the proper table association is ensured by a propagation parameter label setting system in analogy to the one implemented for the halo profile database. Computing such a table on the first call is rather CPU consuming, especially for DM profiles that are singular towards the Galactic center, so in case only a few flux computations are needed it may be better to switch off the tabulation option; this is true also in case the flux is needed at a small number of energy values, since the latest 100 (non-equivalent) calls to dspbtdaxi and dsdbtdaxi are stored in memory (with the corresponding propagation parameters and halo model correctly associated).

The case for positrons is treated analogously, except that energy losses and spatial diffusion are the most important effects for propagating cosmic ray leptons. The transport equation we solve semi-analytically therefore has the form [18]

$$\frac{\partial N}{\partial t} = 0 = \nabla \cdot (D \,\nabla N) + \frac{\partial}{\partial p} \left(\frac{dp}{dt}N\right) + Q, \qquad (10.2)$$

where the functional form of D and Q can be chosen as for antiprotons and antideuterons, and the energy loss rate dp/dt can have a generic momentum dependence but needs again to be spatially constant. Assuming the same topology for the propagation volume and free escape conditions at the vertical boundaries (to compute the local positron flux the radial boundary turns out to be irrelevant), the solution of the propagation equation is given in terms of a Green's function in energy (the function dsepgreenaxi in the code) to be convoluted over the source energy spectrum at emission for a given particle DM candidate. This last step is performed by the function dsepdndpaxi returning the local positron number density, while dsepdphidpaxi converts it to a flux and is the function which should be called from the main file. The method to implement this solution is a slight generalization of the one described [18] and generalizes the one introduced in [?] for a spherically symmetric configuration to an axisymmetric system.

The computation of the Green's function involves a volume integral over the spatially dependent part of the DM source function Q (again basically the DM density  $\rho_{\chi}$  for decaying DM and its square for pair annihilating DM) and the implementation via the so-called method of image charges (again a sum over a series) of the free escape boundary condition. It can again be CPU expensive for singular halo profiles, but its tabulation is always needed since the Green function appears in a convolution. The main limiting factor with respect to full numerical solutions is that one is forced to assume an (spatially) average value for dp/dt. However this has not a severe impact on our results for the local DM-induced positron flux since, especially for energies above 10 GeV, the bulk of the DM contribution to the local flux stems from a rather close-by emission volume; one thus just has to make sure to normalize dp/dt to the mean value for *local* energy losses, as opposed to the mean value in the Galaxy, which are mainly due to the synchrotron and inverse Compton processes.

While the transport equations (10.1) and (10.2) are essentially the same as considered in previous releases of the code, their implementation in the present release is completely new and appears to be numerically more stable. In particular cases with very singular DM profiles still give numerically accurate results and converge faster (in case of antiprotons and antideuterons implementing a procedure which applies to point sources). Note however that the case of very singular DM profiles is also the one in which our models or *any* propagation model is subject to a significant uncertainty related to the underlying physics, since propagation in the Galactic center region is difficult to model and probably rather different from what can be tested in the local neighbourhood by measurements of primary and secondary cosmic rays. Finally, the implementation starting from DarkSUSY 6 is more flexible regarding parameter choices, such as for the rigidity scaling of the diffusion coefficient and energy scaling of energy losses, in a framework which is now fully consistent for antiprotons, antideuterons and positrons.

### cr\_gamma: Cosmic rays – Gamma fluxes

#### 11.1 Gamma rays from the halo – theory

Among the yields of decay or pair annihilations of halo dark matter particles, the role played by gamma-rays could be a major one. Unlike the cases involving charged particles, for gamma-rays it is straightforward to relate the distribution of sources and the expected flux at the earth. Most flux estimated can be obtained just by summing over the contributions along lines of sight (or better, geodesics): gamma-rays have a low enough cross section on gas and dust and therefore the Galaxy is essentially transparent to them (except perhaps in the innermost part, very close to the region where a massive black hole is inferred); absorption by starlight and infrared background becomes effecient only for very far away sources (redshift larger than about 1).

It follows that in case the gamma-ray signal is detectable, this might be the only chance for mapping the fine structure of a dark halo, with a much better resolution for inhomogeneities (clumps) with respect to what is achievable through dynamical measurements or lensing effects. This is especially true for annihilating dark matter. Turning the latter argument around, if the fine structure of the Galactic halo is clumpy, or if a large density enhancement is present towards the Galactic center, as seen in N-body simulations of dark matter halos, this dark matter detection technique may be more promising than indicated by the estimates in which smooth non-singular halo scenarios are considered (recall that the fluxes per unit volume are proportional to the square of the dark matter density locally in space).

Several targets have been discussed as sources of gamma-rays from the annihilation of dark matter particles. An obvious source is the dark halo of our own galaxy [28, 29, 30, 31] and in particular the Galactic center, as the dark matter density profile is expected, in most models, to be picked towards it, possibly with huge enhancements close to te central black hole. The Galactic center is an ideal target for both ground- and space-based gamma-ray telescopes. As satellite experiments have much wider field of views and will provide a full sky coverage, they can in principle test the hypothesis of gamma-rays emitted in clumps of dark matter which may be present in the halo [32, 33, 34, 16, 35, 36]. Another possibility which has been considered is the case of gamma-ray fluxes from external nearby galaxies [37]. Furthermore, it has been proposed to search for an extragalactic flux originated by all cosmological annihilations of dark matter particles [38, 39, 40]. DarkSUSY is suitable to compute the gamma-ray flux from all these (and possibly other) sources.

### 11.2 Continuous gamma yields and line signals

The bulk of the gamma-ray yield from DM annihilation or decay arises from quark jets that when they hadronize/decay give rise to neutral pions which decay to gamma rays. At loop level, it could also be possible to produce monochromatic gamma rays, which could be a smoking gun for dark matter searches. The advantage with the gamma-ray lines is the distinctive spectral signature, which has no plausible astrophysical counterpart.

Compared to the monochromatic flux, the gamma-ray flux produced in  $\pi^0$  decays is much larger but has less distinctive features. The photon spectrum in the process of a pion decaying into  $2\gamma$ is, independent of the pion energy, peaked at half of the  $\pi^0$  mass, about 70 MeV, and symmetric with respect to this peak if plotted in logarithmic variables. Of course, this is true both for pions produced by DM and, e.g., for those generated by cosmic ray protons interacting with the interstellar medium.

When considered together with to the cosmic-ray induced Galactic gamma-ray background, the DM induced signal looks like a component analogous to the secondary flux due to nucleon nucleon interactions: it is drowned into the Bremsstrahlung component at low energy, while it may be the dominant contribution at energies above 1 GeV or so. In fact, if the exotic component is indeed significant an option to disentangle it would be to search for a break in the energy spectrum at about the WIMP mass, where the line feature might be present as well: while the maximal energy for a photon emitted in WIMP pair annihilations is equal to the WIMP mass, the component from cosmic ray protons extends to much higher energies, essentially with the same spectral index as for the proton spectrum (the role played by the third main background component, inverse Compton emission, has still to settled at the time being and may worsen the problem of discrimination against background).

Besides this (weak) spectral feature, another way to disentangle the dark matter signal may be to exploit a directional signature: data with a wide angular coverage should be analyzed to search for a gamma-ray flux component following the shape and density profile of the dark halo, including eventual contributions from clumps.

#### 11.3 Fluxes

Given a density distribution of DM particles, we can define a source function that tells us how many particles that are produced per volume element per time unit (and per energy interval for differential yields) from either annihilation or decay. The DM signal ultimately only depends on the local injection rate of some stable (cosmic ray) particle f, per volume and energy,

$$\frac{d\mathcal{Q}(E_f, \mathbf{x})}{dE_f} = \sum_n \rho_{\chi}^n(\mathbf{x}) \left\langle \mathcal{S}_n(E_f) \right\rangle \,. \tag{11.1}$$

Here,  $\rho_{\chi}$  is the DM density (of the respective component, in case of multi-component DM) and the ensemble average  $\langle ... \rangle$  is taken over the DM velocities; in principle, it depends on the spatial location **x**, but in many applications of interest this can be neglected.

The particle source terms  $S_n(E_f)$  encode the full information about the DM particle physics model. For a typical WIMP DM candidate, e.g., only the annihilation part (n = 2) contributes,

$$\mathcal{S}_2(E_f) = \frac{1}{N_\chi m_\chi^2} \sum_i \sigma_i v \frac{dN_i}{dE_f}, \qquad (11.2)$$

where  $\sigma_i$  is the annihilation cross section of two DM particles into final state *i* and  $dN_i/dE_f$  is the resulting number of stable particles of type *f* per such annihilation and unit energy.  $N_{\chi}$  is a symmetry factor that depends on the nature of DM; if DM is (not) its own antiparticle we have  $N_{\chi} = 2$  ( $N_{\chi} = 4$ ). The right-hand side of the above expression must be further integrated over f(v), the velocity distribution of the *relative* velocity of the two dark matter particles, but in practice it is typically sufficient to evaluate it for v = 0.

For decaying DM, on the other hand, we have

$$S_1(E_f) = \frac{1}{m_\chi} \sum_i \Gamma_i \frac{dN_i}{dE_f}, \qquad (11.3)$$

where  $\Gamma_i$  denotes the partial decay widths. Let us stress, however, that Eq. (11.1) is much more general in that it encapsulates also DM that is *both* annihilating and decaying, multi-component SM, as well as DM models with an internal  $Z^n$  symmetry [41, 42, 43, 44].

For a telescope pointing in the direction  $\psi$ , the expected DM-induced differential flux in gamma rays or neutrinos – i.e. the expected number of particles per unit area, time and energy – from a sky-region  $\Delta \psi$  is thus given by a line-of-sight integral

$$\frac{d\Phi}{dE} = \frac{1}{4\pi} \int_{\Delta\psi} d\Omega \int_{\text{l.o.s.}} d\ell \, \frac{dQ}{dE} \,. \tag{11.4}$$

For an effectively point-like source at distance d, this line-of-sight integral simplifies to

$$\int_{\Delta\psi} d\Omega \int_{\text{l.o.s.}} d\ell \, \frac{d\mathcal{Q}}{dE} \longrightarrow \frac{1}{d^2} \int dV \frac{d\mathcal{Q}}{dE} \,, \tag{11.5}$$

where the integral is over the extention of the source (much smaller than d).

For decaying DM, the above line-of-sight integral always factorizes into the particle source term  $S_1$  given in Eq. (11.3) and a term that only depends on the DM distribution,

$$\frac{d\Phi^{\rm dec}}{dEd\Omega} = \frac{1}{4\pi} J^{\rm dec} S_1 , \qquad J^{\rm dec} \equiv \int_{\rm l.o.s.} d\ell \,\rho \tag{11.6}$$

For annihilating DM, the corresponding factorization strictly speaking *only* holds *if* the annihilation rate is independent of velocity:

$$\frac{d\Phi^{\rm ann}}{dEd\Omega} = \frac{1}{4\pi} J^{\rm ann} S_2 \,, \qquad J^{\rm ann} \equiv \int_{\rm l.o.s.} d\ell \,\rho^2 \,. \tag{11.7}$$

While notable exceptions exist (in particular for resonances [45], *p*-wave annihilation [?] and Sommerfeld-enhanced annihilation [?]), this is a commonly encountered situation and hence of general interest.

#### 11.4 Gamma rays from the halo – routines

DarkSUSY provides the functions dscrgaflux\_dec and dscrgaflux\_v0ann that take  $J^{dec}$  (or  $J^{ann}$ ) as input and return the fluxes given in Eqs. (11.7) and (11.6). Here, the subscript \_v0ann refers to the fact that, for the purpose of those routines,  $S_2$  is evaluated in the limit of vanishing relative velocity of the annihilating DM pair. While this is the only situation of practical interest in many DM models, future DarkSUSY versions will offer support for a full velocity dependence of this quantity. In analogy with dscrsource\_line, the core library furthermore provides routines dscrgaflux\_line\_dec and dscrgaflux\_line\_v0ann (and correspondingly for neutrinos) that return strength, width and location of *monochromatic* ('line') signals. The latter is a convenient change with respect to previous versions of the code, as one cannot generally know how many lines there are (this depends on the particle physics module).

If a halo label is available, one can also more conveniently call dsgafluxsph instead. This function directly returns the gamma-ray flux, for that halo, automatically calculating the required line-of-sight integrals and adding decaying and annihilating DM components depending on which particle model is initialized.

### cr\_nu: Cosmic rays – Neutrino fluxes

### 12.1 Neutrino fluxes from the halo – theory

Usually, the flux of neutrinos from annihilation of WIMPs in the Milky Way halo is too small to be detectable, but for some clumpy or cuspy models, it might be detectable. The calculation of the neutrino-flux follows closely the calculation of the continuous gamma ray flux, with the main addition that neutrino interactions close to the detector are also included. Hence, both the neutrino flux and the neutrino-induced muon flux can be obtained. The neutrino to muon conversion rate in the Earth can also be obtained.
# cr\_ps: Cosmic rays – point sources

## 13.1 Cosmic Ray propagation for point sources

In this directory, we collect the relevant cosmic ray routines for DM point sources like ordinary DM clumps or mini-spikes around intermediate-mass black holes.

# dd: Direct detection

### 14.1 Direct detection – theory

Specific choices of nuclear structure functions can be selected by calling dsddset('sf', label), where the character variable label indicates the set of structure functions. For the default option ('best'), e.g., the code automatically picks the best currently available structure function (depending on the nucleus). This mean, in order Fourier-Bessel, Sum-of-Gaussians, Fermi, Lewin-Smith. The function returning the value of  $\tilde{\sigma}_{\chi T}$  is to be provided by an interface function dsddsigma(v,Er,A,Z,sigij, ierr) residing in the particle physics module, where on input v=v,  $Er=E_R$ , A=A, Z=Z and on output the 27×27 array sigij contains the (partial) equivalent cross sections  $\tilde{\sigma}_{ij}$  in cm<sup>2</sup> and the integer ierr contains a possible error code. The order of the entries in sigij corresponds to that of the independent nonrelativistic operators  $\mathcal{O}_i$ ; for the first 11 entries, we use the same operators and convention as in Ref. [46], while for the last 16 entries we add the additional operators discussed in Ref. [47]. In particular, sigii(1,1) is the usual spin-independent cross section and sigii(4,4) is the usual spin-dependent cross section. In addition, the direct detection module in DarkSUSY provides utility functions that can be used in the computation of the cross section. For example, the subroutine dsddgg2sigma(v, er,A,Z,gg,sigij,ierr) computes the (partial) equivalent scattering cross sections  $\tilde{\sigma}_{ij}$  for nucleus (A, Z) at relative velocity v and recoil energy  $E_R$  starting from values of the  $G_i^N$  constants in gg, with nuclear structure functions set by the previous call to dsddset. The actual nuclear recoil event rate as given in

$$\frac{dR}{dE_R} = \sum_T c_T \frac{\rho_\chi^0}{m_T m_\chi} \int_{v > v_{\min}} \frac{d\sigma_{\chi T}}{dE_R} \frac{f(\mathbf{v}, t)}{v} d^3 v \,, \tag{14.1}$$

finally, is computed by the function dsdddrde. The latter two functions are independent of the specific particle physics implementation and hence are contained in the core library. In the above expression, the sum runs over nuclear species in the detector,  $c_T$  being the detector mass fraction in nuclear species *i*.  $m_T$  is the nuclear (target) mass, and  $\mu_{\chi T} = m_{\chi} m_T / (m_{\chi} + m_T)$  is the reduced DM–nucleus mass. Furthermore,  $\rho_{\chi}^0$  is the local DM density, **v** the DM velocity relative to the detector,  $v = |\mathbf{v}|$ , and  $f(\mathbf{v}, t)$  is the (3D) DM velocity distribution. In order to impart a recoil energy  $E_R$  to the nucleus, the DM particle needs a minimal speed of  $v_{\min} = \sqrt{M_T E_R / 2\mu_{\chi T}^2}$ .

# dmd\_astro: Astrophysical source functions

In the folder src/dmd\_astro we provide a set of wrapper routines that all call the halo driver function dsdmsdriver in a specific way. The rest of the code communicates with dsdmsdriver, and hence the halo model(s), only via these functions and never by calling dsdmsdriver directly. The routines in src/dmd\_astro therefore provide examples of functions that *cannot be replaced* individually in a consistent way, but only as a whole set (along with dsdmsdriver, in case the user wants to change the structure of the driver itself).

# dmd\_mod: Dark matter distributions

### 16.1 Dark matter distributions – theory

All the dark matter detection rates depend in one way or another on the properties of the Milky Way dark matter halo. We will here outline the halo model that by default is included with DarkSUSY.

Observationally, the distribution of DM on scales relevant for DM searches is only poorly constrained. The situation is somewhat improved when instead referring to the results of large *N*-body simulations of gravitational clustering, which consistently find that DM halos on average are well described by Einasto [8] or Navarro-Frenk-White profiles [9], with the halo mass being essentially the only free parameter (after taking into account that the halo concentration strongly correlates with the halo mass [48]). On the other hand, there is a considerable halo-to-halo scatter associated to these findings, so that it remains challenging to make concrete predictions for individual objects – in particular if they are located in cosmologically somewhat 'special' environments like in the case of the Milky Way and its embedding in the Local Group. Even worse, baryonic physics can have a large impact on the DM profiles, especially on their inner parts most relevant for indirect detection, and even though hydrodynamic simulations taking into account such effect have made tremendous progress in recent years [49, 50, 51, 52], there is still a significant uncertainty related to the modelling of the underlying processes. In light of this situation, there is a considerable degree of freedom concerning halo models and the DM density profiles, and a code computing observables related to DM should be able to fully explore this freedom.

#### 16.1.1 Rescaling of the WIMP density

While it is natural to assume that the DM particles described by a given particle module implemented in DarkSUSY make up most of the DM in our galaxy, they may also just constitute a sub-dominant part of a multi-component realization of DM. What is more, there might be both a thermal contribution to the cosmological DM abundance – as computed by the relic density routines in DarkSUSY– and a non-thermal contribution, e.g. via out-of-equilibrium production or via the decay of heavier particles. In this context, it is important to remember that DM detection rates only depend on the *local* DM density of that particular DM candidate. For the case of direct detection, e.g., the rate scales linearly with the local density at earth, while for indirect detection it scales linearly or quadratically with the local DM density at the point of of decay or or annihilation, respectively.

In previous versions of the code, the ratio of thermal relic abundance returned by dsrdomega and observed cosmological DM abundance was used to internally rescale the results from rate calculations. For the reasons given above, this is not fully satisfactory and in any case obscures the origin of this rescaling. Starting from DarkSUSY 6, this is therefore no longer the case. All rate routines now assume that the local DM density equals the local density in the particular DM candidate realized in the particle module – unless they explicitly take the local DM density as an input parameter (which in that case refers to the local density in the particles described by the respective DarkSUSY module). An example for such an exception are the neutrino telescope routines, because the combined effect of DM capture and annihilation makes the dependence on the local DM density more complicated. In all other cases, e.g. if DM rate routines just take a halo label as input, the user has to make sure to rescale the rates, as described above, *by hand* to reflect possible sub-dominant DM populations.

### 16.2 Implementation in DarkSUSY

The implementation of dark matter halo models and related quantities in the library ds\_core follows a new and highly flexible scheme, compared to earlier versions of the code, avoiding pre-defined hardcoded functions. For convenience a few pre-defined options are provided, however these can be either complemented by other profiles eventually needed, or the entire sample configuration can be simply replaced linking to a user-defined setup - in both cases without editing routines provided in this release of the code. A further improvement compared to previous versions of DarkSUSY is that different dark matter density profiles, possibly referring to different dark matter detection targets, can be defined at the same time: E.g., one can easily switch back and forth from a computation of the local positron flux induced by dark matter annihilations or decays in the Milky Way halo to the computation of the gamma-ray flux from an external halo or a Milky Way satellite within the same particle physics scenario. Finally the present implementation simplifies the task of keeping track of consistent definitions for related quantities, such as, e.g., a proper connection between the dark matter profile and the source function for a given dark matter yield (see Chapter 6), or calling within an axisymmetric coordinate system a spherically symmetric function (and preventing the opposite).

At the basis of the implementation, there is the subroutine dsdmsdriver routine which acts as an interface to quantities related to the dark matter density profiles. This routine must contain a complete set of instructions on how to retrieve the different observables: E.g. it checks the scaling of the various DM source functions (Chapter 6 with the DM density  $\rho_{\chi}$  – namely  $\rho$  for decaying dark matter and  $\rho^2$  for dark matter pair annihilations (in case the effect of substructures is neglected) – and passes this information to the routines for propagations of charged cosmic rays in the Galaxy (Chapter 10), assuming that such source function is axisymmetric; line of sight integration routines (needed, e.g. for the computation of gamma-ray fluxes, see Chapter 11) call this same subroutine, but may assume instead that the corresponding source function is spherically symmetric. The routine dsdmsdriver must contain the specification on whether the dark matter density profile is spherically symmetric or axisymmetric, and in the first case provide a consistent numerical output to both calls, in the latter return an error to the second call (since a spherically symmetric profile was expected). It may also be useful to use the dsdmsdriver routine for initialization calls, for instance to set parameters for a given parametric density profile, and test calls, for instance to print which dark matter profile is currently active within a set of available profiles. The input/output structure of the routine is rather general, with the first entry being however fixed to an integer flag specifying the action of the routine; currently available values and relative action are implicitly defined (through integer variables such as 'idmddensph' referring to the spherical dark matter density profile) in an include file and are global parameter. Such set of definitions should not be changed, but can be enlarged in case further profile-related quantities would be needed.

While a specific dsdmsdriver routine should match the user needs in the problem at hand, the present release provides a sample version, illustrating the flexibility of the setup. In particular the version included in the library ds\_core assumes that the dark matter density profile is spherically

symmetric and does not include dark matter substructure; it allows to choose as dark matter profile one among three parametric profiles, namely the Einasto [8], the Navarro-Frenk-White [9] and the Burkert [?] profiles, or a profile interpolated from a table of values of the dark matter density at a given radius. Besides providing parameters as needed in case of parametric profiles, to complete the initialization of a profile one should also specify: *i*) an inner truncation radius, namely some  $r_{ic}$  fixing  $\rho(r) = \rho(r_{ic})$  for any  $r < r_{ic}$  (the choice has an impact on predictions for dark matter rates only for very singular dark matter profiles; choosing a value which is not too small allows for a faster numerical convergence of some rate computations); *ii*) an outer truncation radius, namely some  $r_{oc}$  beyond which the profile is assumed to be zero; *iii*) the distance from the observer of the center of the profile, corresponding to the Sun galactocentric distance only in case the profile refers to the Milky Way; *iv*) whether it is a profile that refers to the Milky Way and hence for which rates that are Milky Way specific, such as the local contribution to antimatter fluxes, can be computed; *v*) whether it is a profile to be saved in a halo profile database for later use.

Regarding this last point, the code implements a procedure of associating the set of entries fully specifying a halo profile (namely the choice of the parametric profile, the corresponding parameters and and the entries i)-v) above) to a given *input label*, and this can be reloaded at any time when needed; in particular all indirect detection flux routines have the label among their input parameters, so that in case of several dark matter detection targets or several profiles for the same targets it becomes unambiguous which profile is being considered. On the other hand it may be the case that the user needs to loop over many different profiles without keeping track of all of them for later reuse (e.g., in a scan over parameter space in the estimate of line of sight integrals towards a dwarf satellite); in such case the profile can be defined as "temporary", with only the latest set temporary profile available at any given time. For halos that are stored in the halo profile database, one can save and/or read from disc tabulated quantities, such as, e.g., the Green function needed for the computation of the local positron flux, for temporary profiles tables can (or in some cases need) to be computed running the code but are overwritten any time the temporary profile is changed. While in the previous releases of DarkSUSY, halo parameters were typically set via common blocks to be included in the main file, the default dsdmsdriver.f implements a procedure in which, when initializing a given halo profile, parameters are given as an input in association with a corresponding parameter label, and profile settings are specified as character strings appearing within the profile label.

Along with the default dsdmsdriver.f routine, which is unfortunately rather involved since it allows for several different options, in the present release we provide example main files which illustrate a few of the possible user needs. Those are described in the 'quick start' part of this manual, see Section 2.2.1, and cover examples of how to use pre-defined halo profiles, read in tabulated profiles as well as how to create completely new ones.

#### **16.2.1** Dark matter distributions – routines

The main routines in this directory are

**dsdmsdriver** Main driver function for setting the halo profile and retrieving observables. All calls to the halo models go through this routine.

**dsdmssethm** Wrapper routine for dsdmsdriver to set a halo model.

**dsdmsselecthm** Wrapper routine for dsdmsdriver to select an already defined halo model.

# dmd\_vel: Dark matter velocity distributions

### 17.1 Dark matter phase-space distributions – theory

The DM velocity and density profiles cannot be chosen independently, in principle, but have to satisfy consistency relations. For a spherically symmetric and isotropic system, e.g., the two profiles are related by the Eddington equation [?, ?]. A fully self-consistent implementation of phase-space distributions will be available with a later DarkSUSY version, at which point the directory dmd\_vel will become obsolete.

Until then, the user can freely choose a DM velocity distribution among those provided in  $src/dmd_vel$  – but should keep in mind this consistency requirement when comparing direct detection rates (which require the local DM velocity profile) to, e.g., the gamma-ray flux from the galactic center (which requires choosing a density profile). Concretely, it is the function dshmuDF that returns the 3D distribution function  $f(\mathbf{v})$  needed by the direct detection routines. It allows to switch between various pre-implemented functional forms, including tabulated velocity profiles, but can of course also be replaced by an arbitrary function supplied by the user (c.f. Section 3.6).

One of these options is the often adopted truncated gaussian, which in the detector frame moving at speed  $v_O$  relative to the galactic halo reads

$$f(v) = \frac{1}{\mathcal{N}_{\text{cut}}} \frac{v^2}{uv_O\sigma} \left\{ \exp\left[-\frac{(u-v_O)^2}{2\sigma^2}\right] - \exp\left[-\frac{\min(u+v_O, v_{\text{cut}})^2}{2\sigma^2}\right] \right\}$$
(17.1)

for  $v_{\rm esc} < v < \sqrt{v_{\rm esc}^2 + (v_O + v_{\rm cut})^2}$  and zero otherwise, with  $u = \sqrt{v^2 + v_{\rm esc}^2}$  and

$$\mathcal{N}_{\rm cut} = \frac{v_{\rm cut}}{\sigma} \exp\left(-\frac{v_{\rm cut}^2}{2\sigma^2}\right) - \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{v_{\rm cut}}{\sqrt{2}\sigma}\right).$$
(17.2)

As default, we have taken the halo line-of-sight (one-dimensional) velocity dispersion  $\sigma = 120 \text{ km/s}$ ,<sup>\*</sup> the galactic escape speed  $v_{\text{cut}} = 600 \text{ km/s}$ , the relative Earth-halo speed  $v_O = 264 \text{ km/s}$  (a yearly average) and the Earth escape speed  $v_{\text{esc}} = 11.9 \text{ km/s}$ . These parameters can be changed by the user.

<sup>\*</sup>Other authors write  $\exp(-3v^2/2\overline{v}^2)$ , in which case  $\overline{v} = \sqrt{3}\sigma$ .

# ini: Initialization routines

## 18.1 Initialisation routines

This directory contains general initialisation routines that need to be called to prepare calculations with DarkSUSY, but which are independent of the particle physics. Most importantly, it contains the subroutine dsinit, which should be called at the beginning of any program using DarkSUSY. In particular, this routine calls various more specific initialisation routines that are relevant only for specific applications (such as the relic density calculation) and therefore reside in the respective directory in src/. It also calls dsinit\_module, which initializes the specific particle module that the user has chosen to link to when compiling the main program. Another subroutine called by dsinit is ini/dsreadnuclides. It reads in names and properties of nuclei and stores them in common blocks, to which both direct detection and solar/earth capture routines need access.

Lastly, the directory /ini also contains two files dsdirname.c.in and dsvername.c.in. These are needed to provide a system-wide reference to the installation directory and version of DarkSUSY, respectively. These refer to include files that are set at configure time, dsdir.h and dsver.h. The user should never have to modify these manually as they are determined at the configure stage.

# kd: Kinetic decoupling

## 19.1 Kinetic decoupling and microhalos (kd) – theory

Even after *chemical decoupling*, which sets the DM relic density (see Section 20.1.1), DM frequently scatters with the very abundant standard model particles and thereby stays in local thermal equilibrium with the heat bath until the temperature has dropped by another factor of between 10 and a few 1000; after *kinetic decoupling*, even these scattering events cease and DM no longer interacts with standard model particles. Inhomogeneities in the DM density can only develop after this has happenened, and the DM particles have sufficiently cooled down so that free streaming becomes negligible. The scale of kinetic decoupling can therefore directly be translated into a cutoff in the power spectrum of (dark) matter fluctuations and thus the size of the smallest (at least when not taking into account primordial black holes) gravitationally bound objects in the universe.

#### 19.1.1 Kinetic decoupling

The process of kinetic decoupling is governed by the full Boltzmann equation for the WIMP distribution function  $f(\mathbf{x}, \mathbf{p})$ , which in a flat Friedmann-Robertson Walker spacetime reads

$$E(\partial_t - H \mathbf{p} \cdot \nabla_{\mathbf{p}}) f = C[f], \qquad (19.1)$$

where C[f] is known as the collision term. The Boltzmann equation quoted in Eq. (20.1) for the description of the (chemical) DM freeze-out process is actually just the first moment of this expression, i.e. obtained by integrating it over  $\int d^3p$  (after dividing it by E). As was realized int [53, 15], kinetic decoupling can be described to a very high precision by considering, instead, the second moment of Eq. (19.1). For this purpose, one introduces the WIMP "temperature"  $T_{\chi}$ ,

$$\int \frac{d^3 p}{(2\pi)^3} \mathbf{p}^2 f(\mathbf{p}) \equiv 3 \, m_\chi T_\chi n_\chi \,, \tag{19.2}$$

as a parameter that characterizes the deviation from thermal equilibrium (for which  $T_{\chi} = T$  holds). After kinetic decoupling, the DM 'temperature' will simply decrease due to the expansion of the universe and, for non-relativistic DM, scale like  $T_{\chi} \propto a^{-2}$ . It is thus natural to define the moment of decoupling as the transition between these two regimes [15]. Allowing for the scattering partners to have a different temperature  $(T_{\tilde{\chi}})$  than the photons (T), this implies

$$T_{\chi}(T) = \begin{cases} T_{\tilde{\gamma}}(T) & \text{for } T \gtrsim T_{\text{kd}} \\ T_{\tilde{\gamma}}(T_{\text{kd}}) \left(\frac{a(T_{\text{kd}})}{a(T)}\right)^2 & \text{for } T \lesssim T_{\text{kd}} \end{cases}$$
(19.3)

For practical purposes, one may now further introduce

$$x \equiv m_{\chi}/T, \qquad (19.4)$$

$$y \equiv m_{\chi} T_{\chi} s^{-2/3} .$$
 (19.5)

Multiplying Eq.(19.1) by  $\mathbf{p}^2/E$ , integrating it over  $\mathbf{p}$  and keeping only the leading order terms<sup>\*</sup> in  $\mathbf{p}^2/m_{\chi}^2$  then results in [15, 55]

$$\frac{d\log y}{d\log x} = \left(1 - \frac{1}{3}\frac{d\log g_{*S}}{d\log x}\right)\frac{\gamma(T_{\tilde{\gamma}})}{H(T)}\left(\frac{y_{\rm eq}}{y} - 1\right).$$
(19.6)

Here,  $g_{*S}$  is the number of effective entropy degrees of freedom,  $y_{eq}$  is the value of y in thermal equilibrium and  $\gamma$  denotes the momentum transfer rate,

$$\gamma(T_{\tilde{\gamma}}) = \frac{1}{48\pi^3 g_{\chi} T_{\tilde{\gamma}} m_{\chi}^3} \sum_{i} \int d\omega \, k^4 \left( 1 \mp g_i^{\pm} \right) g_i^{\pm}(\omega) \, |\mathcal{M}|_{t=0}^2 \sup_{\substack{s=m_{\chi}^2 + 2m_{\chi}\omega + m_{\tilde{\gamma}}^2}},\tag{19.7}$$

where the sum runs over all DM scattering partners (counting, e.g., electrons and positrons separately),  $k \equiv |\mathbf{k}|$  is their momentum and  $\omega$  their energy. The  $g_i$  denote the SM distribution functions, which are assumed to be thermal (note, however, that no assumption has been made about the form of the WIMP distribution function f). The scattering amplitude squared in this expression,  $|\mathcal{M}|^2$ , is understood to be *summed* over all internal (spin or color) degrees of freedom, including initial ones. If it is not Taylor expandable around t = 0, one has to make the replacement [56, 57]

$$\left|\mathcal{M}\right|_{\substack{t=0\\s=m_{\chi}^{2}+2m_{\chi}\omega+m_{\tilde{\gamma}}^{2}}}^{2} \longrightarrow \left\langle\left|\mathcal{M}\right|^{2}\right\rangle_{t} \equiv \frac{1}{8k^{4}} \int_{-4k_{CM}^{2}}^{0} dt(-t) \left|\mathcal{M}\right|^{2}$$
(19.8)

in the above expression. We may easily check that the asymptotic behaviour for  $T_{\chi}$  described by Eq. (19.6) is consistent with the expectation outlined above: At large T, we have  $\gamma \gg H$ , thus enforcing  $T_{\chi} = T$ ; when T becomes small, the WIMPs completely decouple from the thermal bath and y stays constant, i.e.  $T_{\chi} \propto s^{2/3} \propto a^{-2}$ . A practical way to determine the *kinetic decoupling* temperature  $T_{\rm kd}$  as defined in Eq. (19.3) is by solving the above differential equation until y stays constant (indicated by the ' $x \to \infty$ '):

$$x_{\rm kd} = \frac{m_{\chi}}{T_{\rm kd}} \equiv g_{\rm eff}(T_{\rm kd}) |y|_{x \to \infty} .$$
(19.9)

#### 19.1.2 The smallest protohalos

Before kinetic decoupling, WIMPs are tightly coupled to the heat bath, so any small-scale perturbations in the DM fluid would immediately be washed out. For temperatures  $T \leq T_{\rm kd}$ , however, this is no longer the case and perturbations in the DM density start to devolop under the influence of gravity; however, the remaining viscous coupling between the two fluids and, subsequently, the free-streaming of the WIMPs generate an exponential cutoff in the power spectrum [58], with a characteristic comoving damping scale  $k_{\rm fs}$ . The WIMP mass contained in a sphere of the corresponding size is thus given by [15]

$$M_{\rm fs} \approx \frac{4\pi}{3} \rho_{\chi} \left(\frac{\pi}{k_{\rm fs}}\right)^3 = 4.0 \times 10^{-6} \left(\frac{1 + \ln\left(g_{\rm eff}^{1/4} T_{\rm kd}/30 \,\,{\rm MeV}\right)/18.6}{(m_{\chi}/100 \,\,{\rm GeV})^{1/2} \,g_{\rm eff}^{1/4} \left(T_{\rm kd}/30 \,\,{\rm MeV}\right)^{1/2}}\right)^3 M_{\odot} \,.$$
(19.10)

<sup>\*</sup> For very early decoupling, next-order correction terms may become relevant [54].

Acoustic oscillations also have to be taken into account as a damping mechanism and lead to a similar exponential cutoff in the power spectrum [59, 60]. In this case, the characteristic damping mass is given by the total amount of DM inside the horizon at the time of kinetic decoupling:

$$M_{\rm ao} \approx \frac{4\pi}{3} \left. \frac{\rho_{\chi}}{H^3} \right|_{T=T_{\rm kd}} = 3.4 \times 10^{-6} \left( \frac{T_{\rm kd} g_{\rm eff}^{1/4}}{50 \,{\rm MeV}} \right)^{-3} M_{\odot} \,.$$
(19.11)

Note that  $T_{\rm kd}$  in the above expressions only holds when using the definition given in Eq. (19.3); for an alternative definition, the expected magnitude of the cutoff mass has to be correspondingly re-scaled.

In general, the actual cutoff in the power spectrum is given by  $M_{\rm cut} = \max [M_{\rm fs}, M_{\rm ao}]$ ; which of the two physically independent damping mechanisms is more efficient depends on the particle nature of the WIMP. The expected mass for the smallest gravitationally bound objects in the universe is then also simply given by  $M_{\rm cut}$ . Numerically, the formation of protohalos with masses down to the cutoff scale has been confirmed and their evolution could be followed until a redshift of  $z \sim 26$  [61]; the further survival probabilities of these objects, however, as well as the resulting consequences for the indirect detection of DM, are subject to a presently still ongoing discussion.

### **19.2** Kinetic decoupling – routines

Before using any of the routines provided in src/kd for the first time, one has to call **dskdset** in order to make some necessary initializations; in particular, a call to this routine ensures that the relevant tables for the relativistic degrees of freedom in the early universe are correctly read in. Typically, the subroutines of greatest interest will be **dskdtkd** and **dskdmcut**, and there should be no need to call any of the other routines directly.

**dskdtkd** numerically solves the Boltzmann equation (19.6) and determines  $T_{\rm kd}$  as given in Eq. (19.9). Here, special care is taken to accurately handle potential resonances in the scattering amplitude; to this end, **dskdboltz\_init** identifies the location of all relevant resonances and passes this information to **dskdgammarate** where the integral of Eq. (19.7) is performed.

Finally, **dskdmcut** returns the mass cutoff in the power spectrum, with an input parameter determining whether it is  $M_{\rm fs}$  or  $M_{\rm ao}$ ; the default call results in  $M_{\rm cut} = \max[M_{\rm fs}, M_{\rm ao}]$ , i.e. the mass of the smallest protohalos.

There are three interface functions that a particle physics module must provide for the inetic decoupling routines in src/ to work: dskdm2 returns the full scattering matrix element squared, evaluated at t = 0 or averaged over t, while dskdm2simp returns only the leading contribution for small  $\omega$  (expressed as a simple power-law in  $\omega$ , in which case there exists an analytic solution for  $T_{\rm kd}$  [53]). Lastly, the particle physics module must provide a routine dskdparticles which, in analogy to the routine dsrdparticles for the case of chemical decoupling discussed above, sets the location of resonances in  $|\mathcal{M}|^2$ .

# rd: Relic density

## 20.1 Relic density – theoretical background

#### 20.1.1 The Boltzmann equation and thermal averaging

Griest and Seckel [62] have worked out the Boltzmann equation when coannihilations are included. We start by reviewing their expressions and then continue by rewriting them into a more convenient form that resembles the familiar case without coannihilations. This allows us to use similar expressions for calculating thermal averages and solving the Boltzmann equation whether coannihilations are included or not. The implementation in DarkSUSY is based upon the work done in [13].

#### 20.1.2 Review of the Boltzmann equation with coannihilations

Consider annihilation of N DM particles  $\chi_i$  (i = 1, ..., N) with masses  $m_i$  and internal degrees of freedom (statistical weights)  $g_i$ . Also assume that  $m_1 \leq m_2 \leq \cdots \leq m_{N-1} \leq m_N$  and that *R*-parity is conserved. Note that for the mass of the lightest stable of the particles we will use the notation  $m_{\chi}$  and  $m_1$  interchangeably.

The evolution of the number density  $n_i$  of particle *i* is

$$\frac{dn_i}{dt} = -3Hn_i - \sum_{j=1}^N \langle \sigma_{ij} v_{ij} \rangle \left( n_i n_j - n_i^{\text{eq}} n_j^{\text{eq}} \right) 
- \sum_{j \neq i} \left[ \langle \sigma'_{Xij} v_{ij} \rangle \left( n_i n_X - n_i^{\text{eq}} n_X^{\text{eq}} \right) - \langle \sigma'_{Xji} v_{ij} \rangle \left( n_j n_X - n_j^{\text{eq}} n_X^{\text{eq}} \right) \right] 
- \sum_{j \neq i} \left[ \Gamma_{ij} \left( n_i - n_i^{\text{eq}} \right) - \Gamma_{ji} \left( n_j - n_j^{\text{eq}} \right) \right].$$
(20.1)

The first term on the right-hand side is the dilution due to the expansion of the Universe. H is the Hubble parameter. The second term describes  $\chi_i \chi_j$  annihilations, whose total annihilation cross section is

$$\sigma_{ij} = \sum_{X} \sigma(\chi_i \chi_j \to X).$$
(20.2)

The third term describes  $\chi_i \rightarrow \chi_j$  conversions by scattering off the cosmic thermal background,

$$\sigma'_{Xij} = \sum_{Y} \sigma(\chi_i X \to \chi_j Y)$$
(20.3)

being the inclusive scattering cross section. The last term accounts for  $\chi_i$  decays, with inclusive decay rates

$$\Gamma_{ij} = \sum_{X} \Gamma(\chi_i \to \chi_j X).$$
(20.4)

In the previous expressions, X and Y are (sets of) standard model particles involved in the interactions,  $v_{ij}$  is the 'relative velocity' defined by

$$v_{ij} = \frac{\sqrt{(p_i \cdot p_j)^2 - m_i^2 m_j^2}}{E_i E_j}$$
(20.5)

with  $p_i$  and  $E_i$  being the four-momentum and energy of particle *i*, and finally  $n_i^{\text{eq}}$  is the equilibrium number density of particle  $\chi_i$ ,

$$n_i^{\rm eq} = \frac{g_i}{(2\pi)^3} \int d^3 \mathbf{p}_i f_i$$
 (20.6)

where  $\mathbf{p}_i$  is the three-momentum of particle *i*, and  $f_i$  is its equilibrium distribution function. In the Maxwell-Boltzmann approximation it is given by

$$f_i = e^{-E_i/T}.$$
 (20.7)

The thermal average  $\langle \sigma_{ij} v_{ij} \rangle$  is defined with equilibrium distributions and is given by

$$\langle \sigma_{ij} v_{ij} \rangle = \frac{\int d^3 \mathbf{p}_i d^3 \mathbf{p}_j f_i f_j \sigma_{ij} v_{ij}}{\int d^3 \mathbf{p}_i d^3 \mathbf{p}_j f_i f_j}$$
(20.8)

Normally, the decay rate of particles  $\chi_i$  other than the lightest which is stable is much faster than the age of the universe. Since we have assumed *R*-parity conservation, all of these particles decay into the lightest one. So its final abundance is simply described by the sum

$$n = \sum_{i=1}^{N} n_i.$$
 (20.9)

For n we get the following evolution equation

$$\frac{dn}{dt} = -3Hn - \sum_{i,j=1}^{N} \langle \sigma_{ij} v_{ij} \rangle \left( n_i n_j - n_i^{\rm eq} n_j^{\rm eq} \right)$$
(20.10)

where the terms on the second and third lines in Eq. (20.1) cancel in the sum.

The scattering rate of DM particles off particles in the thermal background is much faster than their annihilation rate, because the scattering cross sections  $\sigma'_{Xij}$  are of the same order of magnitude as the annihilation cross sections  $\sigma_{ij}$  but the background particle density  $n_X$  is much larger than each of the DM particle densities  $n_i$  when the former are relativistic and the latter are non-relativistic, and so suppressed by a Boltzmann factor. In this case, the  $\chi_i$  distributions remain in thermal equilibrium, and in particular their ratios are equal to the equilibrium values,

$$\frac{n_i}{n} \simeq \frac{n_i^{\rm eq}}{n^{\rm eq}}.\tag{20.11}$$

We then get

$$\frac{dn}{dt} = -3Hn - \langle \sigma_{\rm eff} v \rangle \left( n^2 - n_{\rm eq}^2 \right) \tag{20.12}$$

where

$$\langle \sigma_{\text{eff}} v \rangle = \sum_{ij} \langle \sigma_{ij} v_{ij} \rangle \frac{n_i^{\text{eq}}}{n^{\text{eq}}} \frac{n_j^{\text{eq}}}{n^{\text{eq}}}.$$
(20.13)

#### 20.1.3 Thermal averaging

So far the reviewing. Now let's continue by reformulating the thermal averages into more convenient expressions.

We rewrite Eq. (20.13) as

$$\langle \sigma_{\text{eff}} v \rangle = \frac{\sum_{ij} \langle \sigma_{ij} v_{ij} \rangle n_i^{\text{eq}} n_j^{\text{eq}}}{n_{\text{eq}}^2} = \frac{A}{n_{\text{eq}}^2} \,. \tag{20.14}$$

For the denominator we obtain, using Boltzmann statistics for  $f_i$ ,

$$n^{\rm eq} = \sum_{i} n_i^{\rm eq} = \sum_{i} \frac{g_i}{(2\pi)^3} \int d^3 p_i e^{-E_i/T} = \frac{T}{2\pi^2} \sum_{i} g_i m_i^2 K_2\left(\frac{m_i}{T}\right)$$
(20.15)

where  $K_2$  is the modified Bessel function of the second kind of order 2.

The numerator is the total annihilation rate per unit volume at temperature T,

$$A = \sum_{ij} \langle \sigma_{ij} v_{ij} \rangle n_i^{\text{eq}} n_j^{\text{eq}} = \sum_{ij} \frac{g_i g_j}{(2\pi)^6} \int d^3 \mathbf{p}_i d^3 \mathbf{p}_j f_i f_j \sigma_{ij} v_{ij}$$
(20.16)

It is convenient to cast it in a covariant form,

$$A = \sum_{ij} \int W_{ij} \frac{g_i f_i d^3 \mathbf{p}_i}{(2\pi)^3 2E_i} \frac{g_j f_j d^3 \mathbf{p}_j}{(2\pi)^3 2E_j}.$$
 (20.17)

 $W_{ij}$  is the (unpolarized) annihilation rate per unit volume corresponding to the covariant normalization of 2E colliding particles per unit volume.  $W_{ij}$  is a dimensionless Lorentz invariant, related to the (unpolarized) cross section through<sup>\*</sup>

$$W_{ij} = 4p_{ij}\sqrt{s\sigma_{ij}} = 4\sigma_{ij}\sqrt{(p_i \cdot p_j)^2 - m_i^2 m_j^2} = 4E_i E_j \sigma_{ij} v_{ij}.$$
 (20.18)

Here

$$p_{ij} = \frac{\left[s - (m_i + m_j)^2\right]^{1/2} \left[s - (m_i - m_j)^2\right]^{1/2}}{2\sqrt{s}}$$
(20.19)

is the momentum of particle  $\chi_i$  (or  $\chi_j$ ) in the center-of-mass frame of the pair  $\chi_i \chi_j$ .

Averaging over initial and summing over final internal states, the contribution to  $W_{ij}$  of a general *n*-body final state is

$$W_{ij}^{n-\text{body}} = \frac{1}{g_i g_j S_f} \sum_{\text{internal d.o.f.}} \int |\mathcal{M}|^2 (2\pi)^4 \delta^4 (p_i + p_j - \sum_f p_f) \prod_f \frac{d^3 \mathbf{p}_f}{(2\pi)^3 2E_f},$$
(20.20)

where  $S_f$  is a symmetry factor accounting for identical final state particles (if there are K sets of  $N_k$  identical particles, k = 1, ..., K, then  $S_f = \prod_{k=1}^K N_k$ !). In particular, the contribution of a two-body final state can be written as

$$W_{ij\to kl}^{2-\text{body}} = \frac{p_{kl}}{16\pi^2 g_i g_j S_{kl} \sqrt{s}} \sum_{\text{internal d.o.f.}} \int |\mathcal{M}(ij \to kl)|^2 \, d\Omega, \tag{20.21}$$

where  $p_{kl}$  is the final center-of-mass momentum,  $S_{kl}$  is a symmetry factor equal to 2 for identical final particles and to 1 otherwise, and the integration is over the outgoing directions of one of the final particles. As usual, an average over initial internal degrees of freedom is performed.

<sup>\*</sup>The quantity  $w_{ij}$  in Ref. [63] is  $W_{ij}/4$ .

#### CHAPTER 20. RD: RELIC DENSITY

We now reduce the integral in the covariant expression for A, Eq. (20.17), from 6 dimensions to 1. Using Boltzmann statistics for  $f_i$  (a good approximation for  $T \leq m$ )

$$A = \sum_{ij} \int g_i g_j W_{ij} e^{-E_i/T} e^{-E_j/T} \frac{d^3 \mathbf{p}_i}{(2\pi)^3 2E_i} \frac{d^3 \mathbf{p}_j}{(2\pi)^3 2E_j},$$
(20.22)

where  $\mathbf{p}_i$  and  $\mathbf{p}_j$  are the three-momenta and  $E_i$  and  $E_j$  are the energies of the colliding particles. Following the procedure in Ref. [12] we then rewrite the momentum volume element as

$$d^{3}\mathbf{p}_{i}d^{3}\mathbf{p}_{j} = 4\pi |\mathbf{p}_{i}|E_{i}dE_{i} 4\pi |\mathbf{p}_{j}|E_{j}dE_{j} \frac{1}{2}d\cos\theta \qquad (20.23)$$

where  $\theta$  is the angle between  $\mathbf{p}_i$  and  $\mathbf{p}_j$ . Then we change integration variables from  $E_i$ ,  $E_j$ ,  $\theta$  to  $E_+$ ,  $E_-$  and s, given by

$$\begin{cases} E_{+} = E_{i} + E_{j} \\ E_{-} = E_{i} - E_{j} \\ s = m_{i}^{2} + m_{j}^{2} + 2E_{i}E_{j} - 2|\mathbf{p}_{i}||\mathbf{p}_{j}|\cos\theta, \end{cases}$$
(20.24)

whence the volume element becomes

$$\frac{d^3 \mathbf{p}_i}{(2\pi)^3 2E_i} \frac{d^3 \mathbf{p}_j}{(2\pi)^3 2E_j} = \frac{1}{(2\pi)^4} \frac{dE_+ dE_- ds}{8},\tag{20.25}$$

and the integration region  $\{E_i \ge m_i, E_j \ge m_j, |\cos \theta| \le 1\}$  transforms into

$$s \ge (m_i + m_j)^2,$$
 (20.26)

$$E_+ \ge \sqrt{s},\tag{20.27}$$

$$\left| E_{-} - E_{+} \frac{m_{j}^{2} - m_{i}^{2}}{s} \right| \le 2p_{ij} \sqrt{\frac{E_{+}^{2} - s}{s}}.$$
(20.28)

Notice now that the product of the equilibrium distribution functions depends only on  $E_+$  and not  $E_-$  due to the Maxwell-Boltzmann approximation, and that the invariant rate  $W_{ij}$  depends only on s due to the neglect of final state statistical factors. Hence we can immediately integrate over  $E_-$ ,

$$\int dE_{-} = 4p_{ij}\sqrt{\frac{E_{+}^2 - s}{s}}.$$
(20.29)

The volume element is now

$$\frac{d^3 \mathbf{p}_i}{(2\pi)^3 2E_i} \frac{d^3 \mathbf{p}_j}{(2\pi)^3 2E_j} = \frac{1}{(2\pi)^4} \frac{p_{ij}}{2} \sqrt{\frac{E_+^2 - s}{s}} dE_+ ds$$
(20.30)

We now perform the  $E_+$  integration. We obtain

$$A = \frac{T}{32\pi^4} \sum_{ij} \int_{(m_i + m_j)^2}^{\infty} ds g_i g_j p_{ij} W_{ij} K_1\left(\frac{\sqrt{s}}{T}\right)$$
(20.31)

where  $K_1$  is the modified Bessel function of the second kind of order 1.

We can take the sum inside the integral and define an effective annihilation rate  $W_{\text{eff}}$  through

$$\sum_{ij} g_i g_j p_{ij} W_{ij} = g_1^2 p_{\text{eff}} W_{\text{eff}}$$
(20.32)

with

$$p_{\text{eff}} = p_{11} = \frac{1}{2}\sqrt{s - 4m_1^2}.$$
(20.33)

In other words

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W_{ij} = \sum_{ij} \sqrt{\frac{[s - (m_i - m_j)^2][s - (m_i + m_j)^2]}{s(s - 4m_1^2)}} \frac{g_i g_j}{g_1^2} W_{ij}.$$
 (20.34)

Because  $W_{ij}(s) = 0$  for  $s \leq (m_i + m_j)^2$ , the radicand is never negative.

In terms of cross sections, this is equivalent to the definition

$$\sigma_{\rm eff} = \sum_{ij} \frac{p_{ij}^2}{p_{11}^2} \frac{g_i g_j}{g_1^2} \sigma_{ij}.$$
 (20.35)

Eq. (20.31) then reads

$$A = \frac{g_1^2 T}{32\pi^4} \int_{4m_1^2}^{\infty} ds p_{\text{eff}} W_{\text{eff}} K_1\left(\frac{\sqrt{s}}{T}\right)$$
(20.36)

This can be written in a form more suitable for numerical integration by using  $p_{\text{eff}}$  instead of s as integration variable. From Eq. (20.33), we have  $ds = 8p_{\text{eff}}dp_{\text{eff}}$ , and

$$A = \frac{g_1^2 T}{4\pi^4} \int_0^\infty dp_{\text{eff}} p_{\text{eff}}^2 W_{\text{eff}} K_1\left(\frac{\sqrt{s}}{T}\right)$$
(20.37)

with

$$s = 4p_{\rm eff}^2 + 4m_1^2 \tag{20.38}$$

So we have succeeded in rewriting A as a 1-dimensional integral.

From Eqs. (20.37) and (20.15), the thermal average of the effective cross section results

$$\langle \sigma_{\rm eff} v \rangle = \frac{\int_0^\infty dp_{\rm eff} p_{\rm eff}^2 W_{\rm eff} K_1\left(\frac{\sqrt{s}}{T}\right)}{m_1^4 T \left[\sum_i \frac{g_i}{g_1} \frac{m_i^2}{m_1^2} K_2\left(\frac{m_i}{T}\right)\right]^2}.$$
(20.39)

This expression is very similar to the case without coannihilations, the differences being the denominator and the replacement of the annihilation rate with the effective annihilation rate. In the absence of coannihilations, this expression correctly reduces to the formula in Gondolo and Gelmini [12].

The definition of an effective annihilation rate independent of temperature is a remarkable calculational advantage. As in the case without coannihilations, the effective annihilation rate can in fact be tabulated in advance, before taking the thermal average and solving the Boltzmann equation.

In the effective annihilation rate, coannihilations appear as thresholds at  $\sqrt{s}$  equal to the sum of the masses of the coannihilating particles. We show an example for neutralino DM (in the mssm module) in Fig. 20.1 where it is clearly seen that the coannihilation thresholds appear in the effective invariant rate just as final state thresholds do. For the same example, Fig. 20.2 shows the differential annihilation rate per unit volume  $dA/dp_{\text{eff}}$ , the integrand in Eq. (20.37), as a function of  $p_{\text{eff}}$ . We have chosen a temperature  $T = m_{\chi}/20$ , a typical freeze-out temperature. The Boltzmann suppression contained in the exponential decay of  $K_1$  at high  $p_{\text{eff}}$  is clearly visible. At higher temperatures the peak shifts to the right and at lower temperatures to the left. For the particular model shown in Figs. 20.1–20.2, the relic density results  $\Omega_{\chi}h^2 = 0.030$  when coannihilations are included and  $\Omega_{\chi}h^2 = 0.18$  when they are not. Coannihilations have lowered  $\Omega_{\chi}h^2$  by a factor of 6.

### 20.1.4 Reformulation of the Boltzmann equation

We now follow Gondolo and Gelmini [12] to put Eq. (20.12) in a more convenient form by considering the ratio of the number density to the entropy density,

$$Y = \frac{n}{s}.$$
 (20.40)



Figure 20.1: The effective invariant annihilation rate  $W_{\text{eff}}$  as a function of  $p_{\text{eff}}$  for an example mssm model. The final state threshold for annihilation into  $W^+W^-$  and the coannihilation thresholds, as given by Eq. (20.34), are indicated. The  $\chi_2^0\chi_2^0$  coannihilation threshold is too small to be seen.



Figure 20.2: Total differential annihilation rate per unit volume  $dA/dp_{\text{eff}}$  for the same model as in Fig. 20.1, evaluated at a temperature  $T = m_{\chi}/20$ , typical of freeze-out. Notice the Boltzmann suppression at high  $p_{\text{eff}}$ .

Consider

$$\frac{dY}{dt} = \frac{d}{dt} \left(\frac{n}{s}\right) = \frac{\dot{n}}{s} - \frac{n}{s^2} \dot{s}$$
(20.41)

where dot means time derivative. In absence of entropy production,  $S = R^3 s$  is constant (R is the scale factor). Differentiating with respect to time we see that

$$\dot{s} = -3\frac{\dot{R}}{R}s = -3Hs \tag{20.42}$$

which yields

$$\dot{Y} = \frac{\dot{n}}{s} + 3H\frac{n}{s}.\tag{20.43}$$

Hence we can rewrite Eq. (20.12) as

$$\dot{Y} = -s \langle \sigma_{\rm eff} v \rangle \left( Y^2 - Y_{\rm eq}^2 \right).$$
(20.44)

The right-hand side depends only on temperature, and it is therefore convenient to use temperature T instead of time t as independent variable. Defining  $x = m_1/T$  we have

$$\frac{dY}{dx} = -\frac{m_1}{x^2} \frac{1}{3H} \frac{ds}{dT} \langle \sigma_{\text{eff}} v \rangle \left( Y^2 - Y_{\text{eq}}^2 \right).$$
(20.45)

where we have used

$$\frac{1}{\dot{T}} = \frac{1}{\dot{s}}\frac{ds}{dT} = -\frac{1}{3Hs}\frac{ds}{dT}$$
(20.46)

which follows from Eq. (20.42). With the Friedmann equation in a radiation dominated universe

$$H^2 = \frac{8\pi G\rho}{3},$$
 (20.47)

where G is the gravitational constant, and the usual parameterization of the energy and entropy densities in terms of the effective degrees of freedom  $g_{\text{eff}}$  and  $h_{\text{eff}}$ ,

$$\rho = g_{\text{eff}}(T) \frac{\pi^2}{30} T^4, \quad s = h_{\text{eff}}(T) \frac{2\pi^2}{45} T^3,$$
(20.48)

we can cast Eq. (20.45) into the form [12]

$$\frac{dY}{dx} = -\sqrt{\frac{\pi}{45G}} \frac{g_*^{1/2} m_1}{x^2} \langle \sigma_{\rm eff} v \rangle \left( Y^2 - Y_{\rm eq}^2 \right)$$
(20.49)

where  $Y_{eq}$  can be written as

$$Y_{\rm eq} = \frac{n_{\rm eq}}{s} = \frac{45x^2}{4\pi^4 h_{\rm eff}(T)} \sum_i g_i \left(\frac{m_i}{m_1}\right)^2 K_2\left(x\frac{m_i}{m_1}\right), \qquad (20.50)$$

using Eqs. (20.15), (20.40) and (20.48). The parameter  $g_*^{1/2}$  is defined as

$$g_*^{1/2} = \frac{h_{\text{eff}}}{\sqrt{g_{\text{eff}}}} \left( 1 + \frac{T}{3h_{\text{eff}}} \frac{dh_{\text{eff}}}{dT} \right)$$
(20.51)

For  $g_{\rm eff}$ ,  $h_{\rm eff}$  and  $g_*^{1/2}$ , the user can choose different implementations by a call to the routine dsrdset (see that routine for further details). The default for those effective relativistic degrees of freedom is to use the results from Drees et al [64].

To obtain the relic density we integrate Eq. (20.54) from x = 0 to  $x_0 = m_{\chi}/T_0$  where  $T_0$  is the photon temperature of the Universe today. The relic density today in units of the critical density is then given by

$$\Omega_{\chi} = \rho_{\chi}^0 / \rho_{\rm crit} = m_{\chi} s_0 Y_0 / \rho_{\rm crit} \tag{20.52}$$

where  $\rho_{\text{crit}} = 3H^2/8\pi G$  is the critical density,  $s_0$  is the entropy density today and  $Y_0$  is the result of the integration of Eq. (20.54). With a background radiation temperature of  $T_0 = 2.726$  K we finally obtain

$$\Omega_{\chi}h^2 = 2.755 \times 10^8 \frac{m_{\chi}}{\text{GeV}} Y_0.$$
(20.53)

## 20.2 Relic density – numerical integration of the density equation

Let us write the evolution equation for the density,

$$\frac{dY}{dx} = -\sqrt{\frac{\pi}{45G}} \frac{g_*^{1/2} m_1}{x^2} \langle \sigma_{\rm eff} v \rangle \left( Y^2 - Y_{\rm eq}^2 \right)$$
(20.54)

as

$$\frac{dY}{dx} = \lambda(Y^2 - q^2), \qquad (20.55)$$

where  $\lambda$  contains the annihilation rate and q represents the thermal-equilibrium density.

This equation is stiff and an explicit method, like Euler or Runge-Kutta, fails to converge. To obtain a numerical solution, we use an adaptive implicit trapezoidal method which we explain in the following. Basically we discretize the equation first with a trapezoidal then with an Euler method, and adapt the step size according to the difference in the updated function values.

For simplicity we denote the right hand wide of eq. (20.55) as f(x). We further write  $f_i = f(x_i)$ and similarly for the other functions  $\lambda(x)$  and q(x). Given  $Y_i = Y(x_i)$  we find  $Y_{i+1} = Y(x_{i+1})$  with  $x_{i+1} = x_i + h$  as follows.

First we discretize the evolution equation as

$$Y_{i+1} - Y_i = h \frac{f_i + f_{i+1}}{2}.$$
(20.56)

We insert

$$f_i = \lambda_i \left( Y_i^2 - q_i^2 \right), \tag{20.57}$$

$$f_{i+1} = \lambda_{i+1} \left( Y_{i+1}^2 - q_{i+1}^2 \right), \qquad (20.58)$$

and solve the resulting quadratic equation for  $Y_{i+1}$  to obtain

$$Y_{i+1} = \frac{c}{1 + \sqrt{1 + uc}},\tag{20.59}$$

where

$$c = 2Y_i + u \left[ (q_{i+1}^2 + \rho q_i^2) - \rho Y_i^2 \right], \qquad (20.60)$$

$$u = h\lambda_{i+1}, \tag{20.61}$$

$$\rho = \lambda_i / \lambda_{i+1}. \tag{20.62}$$

In the expression for c we have explicitly indicated the order of evaluation which we found avoids round-off errors. If in eq. (20.59) 1 + uc is negative, we simply reduce the step h to h/2 and try again. Secondly we discretize the evolution equation as

$$Y_{i+1} - Y_i = h f_{i+1}. (20.63)$$

We insert the expression for  $f_{i+1}$  and solve the quadratic equation for  $Y_{i+1}$  to obtain

$$Y'_{i+1} = \frac{1}{2} \frac{c'}{1 + \sqrt{1 + uc'}},\tag{20.64}$$

where

$$c' = 4\left(Y_i + uq_{i+1}^2\right). \tag{20.65}$$

Again if in eq. (20.64) 1 + uc' < 0, we reduce the step h to h/2 and try again.

We then adapt the step size according to the relative difference of  $Y_{i+1}$  and  $Y'_{i+1}$ ,

$$d = \left| \frac{Y_{i+1} - Y'_{i+1}}{Y_{i+1}} \right|.$$
(20.66)

If the difference is larger than a prefixed  $\epsilon$ , set at 0.01, we reduce the step size h to  $hs/\sqrt{\epsilon}$  but never to less than h/10. s is a safety factor set to 0.9. If  $d < \epsilon$ , we increase the step size by a factor  $s/\sqrt{\epsilon}$  but never by more than a factor of 5. We do not allow the step size to become smaller than  $h_{\min} = 10^{-9}$ . Error code 5 is reported if this happens. Error code 4 occurs when  $x_{i+1}$  is numerically equal to  $x_i$  because of round-off. Error code 6 occurs when the number of steps exceeds a maximum of 100000. Finally the initial step size is taken to be 0.01.

### 20.3 Relic density – routines

In src/rd, the general relic density routines are found. These routines can be used for any dark matter candidate and e.g. the interface to neutralino dark matter is in  $src_models/mssm/rd$ .

The main routine for relic density calculations is dsrdomega. It calculates and returns the relic density as well as the approximate temperature of freeze-out. It taks a few arguments: i) one option argument that typically (depending on the particle physics module) determines how co-annihilations are included, ii) one fast argument that determines how careful the Boltzmann solver should be in calculating the relic density. See the header of dsrdomega for more details.

#### 20.3.1 Global parameters

All internal settings of the relic density routines are set in common blocks in dsrdcom.h. The most important parameters that can be changed by the user are

<u>.</u>	
Purpose:	Provide a set of parameters, with which the internal behaviour of the relic
	density routines can be changed.
Parameters	
tharsi i	Size of the coannihilation, resonance and threshold arrays (default=50). Increase this size if you have more than 50 coannihilating particles, more than 50 resonances or more than 50 thresholds.
rdluerr i	Logical unit number where error messages are printed.
rdtag c*12	Idtag that is printed in case of errors.
cosmin r8	
waccd r8	
dpminr r8	
dpthr r8	
wdiffr r8	

Important parameters in dsrdcom.h

wdifft r8 ...
hstep r8 ...
rdt\_max r8 Maximum number of seconds to spend on relic density calculation. If the time
limit is exceeded, dsrdomega returns with an error flag and the result 0. The
time is the total CPU time (i.e. summed up over all cores/threads) and the
limit is approximate as it is only checked before a new point is added to the
W<sub>eff</sub> tabulation.

### 20.3.2 Brief description of the internal routines

Below, the remaining routines related to the relic density calculation are briefly mentioned. For more details, we refer to the routines themselves.

Routine	Purpose		
dsrdaddpt	To add one point in the $W_{\text{eff}}$ - $p_{\text{eff}}$ table.		
dsrdcom	To initialize parameters in the common blocks in dsrdcom.h. If you want to change these parameters yourself, include dsrdcom.h in your code and change the parameters you want.		
dsrddof	Returns the degrees of freedom as a function of the temperature in the early Universe.		
dsrddpmin	To return the allowed minimal distance in $p_{\text{eff}}$ between two points in the $W_{\text{eff}}$ - $p_{\text{eff}}$ plane. The returned value depends on if there is a resonance present or not at the given $p_{\text{eff}}$ .		
dsrdens	Routine to solve the Boltzmann equation and return the relic density. Called by dsrdomega.		
dsrdeqn	To solve the relic density equation by means of an implicit trapezoidal method with adaptive stepsize and termination.		
dsrdfunc	To return the invariant annihilation rate times the thermal distribution.		
dsrdfuncs	To provide dsrdfunc in a form suitable for numerical integration.		
dsrdlny	To return $\ln W_{\text{eff}}$ for a given $p_{\text{eff}}$ .		
dsrdnormlz	To return a unit vector in a given direction.		
dsrdqad	To calculate the relic density with a quick-and-dirty method. It uses the approximative expressions in Kolb & Turner with the cross section expaned in $v$ .		
dsrdgrkck	To numerically integrate a function with a Runge-Kutta method		
dsrdrhs	To calculate terms on the right-hand side in the Boltzmann equation.		
dsrdset	To set the control parameters for the relic density calculation. Currently, only the choice of effective degrees of freedom is implemented through dsrdset; the other parameters are passed as arguments to dsrdomega.		
dsrdspline	To set up the table $W_{\text{eff}}$ - $p_{\text{eff}}$ for spline interpolation.		
dsrdstart	To sort and store information about coannihilations, resonances and thresholds in common blocks.		
dsrdtab	To set up the table $W_{\text{eff}}$ - $p_{\text{eff}}$ .		
dsrdthav	To calculate the thermally averaged annihilation cross section at a given temperature.		
dsrdthclose			
dsrdthlim	To determine the end-points for the thermal average integration.		
dsrdthtest	To check if a given entry in the $W_{\text{eff}}$ - $p_{\text{eff}}$ table is at a threshold.		
dsrdwfunc	To write out dsrdfunc for a given $x = m_{\chi}/T$ .		

dsrdwintp	To return the invariant rate $W_{\text{eff}}$ for any given $p_{\text{eff}}$ by performing a spline
	interpolation in the $W_{\text{eff}}$ - $p_{\text{eff}}$ table.
dsrdwintpch	To check the spline interpolation in the $W_{\text{eff}}$ - $p_{\text{eff}}$ table and compare with a linear interpolation.
dsrdwintrp	To write out a table of the invariant rate $W_{\text{eff}}$ and some internal integration variables and expressions.
dsrdwres	To write out the table $W_{\text{eff}}$ - $p_{\text{eff}}$ .

# se\_aux: Auxiliary routines for WIMP annihilation in the Sun/Earth

## 21.1 Sun and Earth models – auxiliary routines

This directory contains parameterizations and semi-analytical calculations of auxiliary fluxes, like Earth atmospheric fluxes. They are not up to date with the latest calculations and are provided as is for convenience.

# se\_mod: Sun and Earth models

(JE: I am responsible for updating this part)

## 22.1 Sun and Earth models – theory

We need to have accurate models for the Sun and the Earth, both in terms of mass distributions, but also in terms chemical compositions etc. We here include a set of Sun and Earth models.

The models here are used for the capture rate calculation for capture of WIMPs in the Sun and the Earth, but they are also used for external packages like **WimpSim** which simulates neutrino interactions and oscillations on the way out of the Sun/Earth. Those simulation results are included as tables in se\_yield.

### 22.2 Sun and Earth models – routines

The main routines that could be of interest to most users are

# se\_nu: Capture and annihilation in the Sun/Earth

### 23.1 Neutrinos from the Sun and Earth – theory

. Note. This chapter is not up to date with the latest additions to the code, but provided as is as a reference right now.

There are several indirect methods for detection of WIMPs. One of the most promising [65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 17] is to make use of the fact that scattering of halo neutralinos by the Sun and the planets, in particular the Earth, during the several billion years that the Solar system has existed, will have trapped these neutralinos within these astrophysical bodies. Being trapped within the Solar or terrestrial material, they will sink towards the center, where a considerable enrichment and corresponding increase of annihilation rate will occur.

Searches for neutralino annihilation into neutrinos will be subject to extensive experimental investigations in view of the new neutrino telescopes (AMANDA, IceCube, Baikal, NESTOR, ANTARES) planned or under construction [80]. A high-energy neutrino signal in the direction of the centre of the Sun or Earth is an excellent experimental signature which may stand up against the background of neutrinos generated by cosmic-ray interactions in the Earth's atmosphere.

There are several different approximations one could do, or processes to include when calculating the capture rates in the Earth/Sun and many of these are coded into DarkSUSY. The default in DarkSUSY is always to use the best calculations available, but more approximate (older) routines are also available, as well as more speculative signals, like the Damour-Krauss signal (not included by default). If you want to use something else than the defaults, or want to call more internal rotuines (more internal than dsntrates or dsntdiffrates), you should read the following sections carefully.

#### 23.1.1 Neutrino yield from annihilations

The differential neutrino flux from neutralino annihilation is

$$\frac{dN_{\nu}}{dE_{\nu}} = \frac{\Gamma_A}{4\pi D^2} \sum_f B_{\chi}^f \frac{dN_{\nu}^f}{dE_{\nu}}$$
(23.1)

where  $\Gamma_A$  is the annihilation rate, D is the distance of the detector from the source (the central region of the Earth or the Sun), f is the neutralino pair annihilation final states, and  $B_{\chi}^{f}$  are the branching ratios into the final state f.  $dN_{\nu}^{f}/dE_{\nu}$  are the energy distributions of neutrinos generated

by the final state f and are obtained from the PYTHIA simulations described in section ??. (JE: Update reference and description of WimpSim simulations.)

In comparison with calculations using the results of [81] (e.g. [82, 75, 83, 78, 77, 84, 85]), this Monte Carlo treatment of the neutrino propagation through the Sun does not need the simplifying assumptions previously made, namely neutral currents are no more assumed to be much weaker than charged currents and energy loss is no more considered continuous.

The neutrino-induced muon flux may be detected in a neutrino telescope by measuring the muons that come from the direction of the centre of the Sun or Earth. For a shallow detector, this usually has to be done in the case of the Sun by looking (as always the case for the Earth) at upward-going muons, since there is a huge background of downward-going muons created by cosmic-ray interactions in the atmosphere. There is always in addition a more isotropic background coming from muon neutrinos created on the other side of the Earth in such cosmic-ray events (and also from cosmic-ray interactions in the outer regions of the Sun). The flux of muons at the detector is given by

$$\frac{dN_{\mu}}{dE_{\mu}} = N_A \int_{E_{\mu}^{\text{th}}}^{\infty} dE_{\nu} \int_0^{\infty} d\lambda \int_{E_{\mu}}^{E_{\nu}} dE'_{\mu} P(E_{\mu}, E'_{\mu}; \lambda) \frac{d\sigma_{\nu}(E_{\nu}, E'_{\mu})}{dE'_{\mu}} \frac{dN_{\nu}}{dE_{\nu}}, \qquad (23.2)$$

where  $\lambda$  is the muon range in the medium (ice or water for the large detectors in the ocean or at the South Pole, or rock which surrounds the smaller underground detectors),  $d\sigma_{\nu}(E_{\nu}, E'_{\mu})/dE'_{\mu}$  is the weak interaction cross section for production of a muon of energy  $E'_{\mu}$  from a parent neutrino of energy  $E_{\nu}$ , and  $P(E_{\mu}, E'_{\mu}; \lambda)$  is the probability for a muon of initial energy  $E'_{\mu}$  to have a final energy  $E_{\mu}$  after passing a path–length  $\lambda$  inside the detector medium.  $E^{\text{th}}_{\mu}$  is the detector threshold energy, which for "small" neutrino telescopes like Baksan, MACRO and Super-Kamiokande is around 1 GeV. Large area neutrino telescopes in the ocean or in Antarctic ice typically have thresholds of the order of tens of GeV, which makes them sensitive mainly to heavy neutralinos (above 100 GeV) [17].

The integrand in Eq. (23.2) is weighted towards high neutrino energies, both because the cross section  $\sigma_{\nu}$  rises approximately linearly with energy and because the average muon energy, and therefore the range  $\lambda$ , also grow approximately linearly with  $E_{\nu}$ . Therefore, final states which give a hard neutrino spectrum (such as heavy quarks,  $\tau$  leptons and W or Z bosons) are usually more important than the soft spectrum arising from light quarks and gluons.

#### 23.1.2 Evolution of the number density in the Earth/Sun

Neutralinos are steadily being trapped in the Sun or Earth by scattering, whereas annihilations take them away. Let N(t) be the total number of neutralinos trapped, at time t, in the core of, for example, the Earth. The annihilation rate of neutralino pairs can be written as

$$\Gamma_a(t) = \frac{1}{2} C_a N^2(t) .$$
(23.3)

The evolution of N(t) is the result of the competition between capture and annihilation:

$$\frac{dN}{dt} = C_c(t) - C_a N^2 \tag{23.4}$$

The constant  $C_c$  is the capture rate, and  $C_a$  entering equations (23.3) and (23.4) is linked to the annihilation cross-section  $\sigma_a$ , and to some effective volumes  $V_j$ , j = 1, 2, taking into account the quasi-thermal distribution of neutralinos in the Earth core:

$$C_a = \langle \sigma_a \, v \rangle \, \frac{V_2}{V_1^2} \,, \tag{23.5}$$

$$V_j \simeq 2.3 \times 10^{25} \left(\frac{j \, m_X}{10 \, \text{GeV}}\right)^{-3/2} \, \text{cm}^3 \,.$$
 (23.6)

This has the solution for the annihilation rate implemented in DarkSUSY

$$\Gamma_A = \frac{C_c}{2} \tanh^2\left(\frac{t}{\tau}\right),\tag{23.7}$$

where the equilibration time scale  $\tau = 1/\sqrt{C_c C_a}$ . In most cases for the Sun, and in the cases of observable fluxes for the Earth,  $\tau$  is much smaller than a few billion years, and therefore equilibrium is often a good approximation ( $\dot{N}(t) = 0$ ). This means that it is the capture rate which is the important quantity that determines the neutrino flux. However, in the program we keep the exact formula (23.7), with some modifications discussed in Sec. ??).

#### 23.1.3 Approximate capture rate expressions

The capture rate induced by scalar (spin-independent) interactions between the neutralinos and the nuclei in the interior of the Earth or Sun is the most difficult one to compute, since it depends sensitively on the Higgs mass, form factors, and other poorly known quantities. However, this spin-independent capture rate calculation is the same as for direct detection treated in Chapter 14. Therefore, there is a strong correlation between the neutrino flux expected from the Earth (which is mainly composed of spin-less nuclei) and the signal predicted in direct detection experiments [17, 86]. It seems that even the large (kilometer-scale) neutrino telescopes planned, when searching for neutralino annihilation in the Earth, will not be competitive with the next generation of direct detection experiments when it comes to detecting neutralino dark matter. However, the situation concerning the Sun is more favourable. Due to the low counting rates for the spin-dependent interactions in terrestrial detectors, high-energy neutrinos from the Sun constitute a competitive and complementary neutralino dark matter search. Of course, even if a neutralino is found through direct detection, it will be extremely important to confirm its identity and investigate its properties through indirect detection. In particular, the mass can be determined with reasonable accuracy by looking at the angular distribution of the detected muons [87, 88].

For the Sun, dominated by hydrogen, the axial (spin-dependent) cross section is important and relatively easy to compute. A reasonably good approximation is given by [89]

$$\frac{C_{\odot}^{\rm sd}}{(1.3 \cdot 10^{23} \,\mathrm{s}^{-1}) \,(270 \,\mathrm{km}\,\mathrm{s}^{-1}/\bar{v})} = \left(\frac{\rho_{\chi}}{0.3 \,\,\mathrm{GeV}\,\mathrm{cm}^{-3}}\right) \left(\frac{100 \,\mathrm{GeV}}{m_{\chi}}\right) \left(\frac{\sigma_{p\chi}^{\rm sd}}{10^{-40} \,\,\mathrm{cm}^{2}}\right)$$
(23.8)

where  $\sigma_{p\chi}^{\rm sd}$  is the cross section for neutralino-proton elastic scattering via the axial-vector interaction,  $\bar{v}$  is the dark-matter velocity dispersion, and  $\rho_{\chi}$  is the local dark matter mass. The capture rate in the Earth is dominated by scalar interactions, where there may be kinematic and other enhancements, in particular if the mass of the neutralino almost matches one of the heavy elements in the Earth. For this case, a more detailed analysis is called for, which is available in [90] with convenient approximations in [89]. In fact, also for the Sun the spin-independent contribution can be important, in particular iron may contribute non-negligibly. For the Sun, the approximation in [89] is also available,

$$\frac{C_{\odot}^{\rm si}}{(4.8 \cdot 10^{22} \,\mathrm{s}^{-1}) \,(270 \,\mathrm{km}\,\mathrm{s}^{-1}/\bar{v})} = \left(\frac{\rho_{\chi}}{0.3 \,\mathrm{GeV}\,\mathrm{cm}^{-3}}\right) \left(\frac{100 \,\mathrm{GeV}}{m_{\chi}}\right) \times \sum_{A} \left(\frac{\sigma_{A}^{\rm si}}{10^{-40} \,\mathrm{cm}^{2}}\right) F_{A}(m_{\chi}) f_{A} \phi_{A} S\left(m_{\chi}/m_{A}\right)/m_{A},$$
(23.9)

where  $f_A$  is the mass fraction of element A and  $\phi_A$  is the typical gravitational potential (relative to the surface) for that element. I.e. an element that is concentrated in the core will have a higher  $\phi_A$ 

		Average parameters	
Element	Mass number $(A)$	$f_i$	$\phi_i$
Hydrogen, H	1	0.670	3.15
Helium-4, <sup>4</sup> He	4	0.311	3.40
Carbon, C	12	0.00237	2.85
Nitrogen, N	14	0.00188	3.83
Oxygen, O	16	0.00878	3.25
Neon, Ne	20	0.00193	3.22
Magnesium, Mg	24	0.000733	3.22
Silicon, Si	28	0.000798	3.22
Sulphur, S	32	0.000550	3.22
Iron, Fe	56	0.00142	3.22

Table 23.1: The composition of the Sun with average parameters to be used in the approximative relations given in [89]. These values are updated with the solar model of [91] and differs slightly from the values used in [89].

	Mass	Mass	fraction	Average p	arameters
Element	number $(A)$	Core	Mantle	$f_i$	$\phi_i$
Oxygen, O	16	0.0	0.440	0.298	1.20
Silicon, Si	28	0.06	0.210	0.162	1.24
Magnesium, Mg	24	0.0	0.228	0.154	1.20
Iron, Fe	56	0.855	0.0626	0.319	1.546
Calcium, Ca	40	0.0	0.0253	0.0171	1.20
Phosphor, P	30	0.002	0.00009	0.00071	1.56
Sodium, Na	23	0.0	0.0027	0.00183	1.20
Sulphur, S	32	0.019	0.00025	0.0063	1.59
Nickel, Ni	59	0.052	0.00196	0.0181	1.57
Aluminum, Al	27	0.0	0.0235	0.0159	1.20
Chromium, Cr	52	0.009	0.0026	0.0047	1.44

Table 23.2: The composition of the Earth's core and mantle. The core mass fractions are from [92][Table 4] and the mantle mass fractions are from [92][Table 2]. The average mass fractions and potentials in the last two columns are weighted averages assuming a core mass of  $1.93 \cdot 10^{24}$  kg and a mantle mass of  $4.04 \cdot 10^{24}$  kg with average potentials (relative to the surface) of 1.6 in the core and 1.2 in the mantle [90].

than an element at the surface. A is the atomic number of the element and  $M_A$  is its mass. The factor S is a kinematical suppression factor [89, 74]. In the next subsection we will go through the compositions of the Earth/Sun that we use.

The approximate capture rate expressions above are coded into the routines dsntcapsun and dsntcapearth. More accurate expressions will follow in the coming subsections.

### 23.1.4 Earth and Sun composition

When the capture rates are calculated, we need to know the composition and density of the Earth/Sun as a function of depth.

In [89] they used average mass fractions and potentials for the location of the various elements in the Sun. We have updated these to the BP2000 [91] values instead, as given in Table 23.1

For the Earth, we have also implemented more accurate density profiles and more up-to date chemical distributions within the Earth. We use the estimates for the Earth composition given in [92][Table 2 for the mantle and Table 4 for the core]. In Table 23.2 we list these values together with



Figure 23.1: In a) the density profile and in b) the escape velocity in the Earth is shown.

the average parameters  $f_i$  and  $\phi_i$  that should be used in the expressions for the approximate capture rates in the previous section. Note that using these average parameters instead of integrating over the full radius is equivalent to putting all the elements of the give type at the gravitational potential  $\phi_i$ .

We also need the density profile of the Earth, and for this we use the values in [93]. Using this density profile, we can calculate the gravitational potential,  $\phi(r)$  inside the Earth and from this one the escape velocity v inside the Earth,

$$v = 11.2 \sqrt{\frac{\phi(r)}{\phi(R_{\oplus})}} \text{ km/s.}$$
(23.10)

In Fig. 23.1 we show the density profile and escape velocity inside the Earth.

#### 23.1.5 More accurate capture rate expressions

Another complicating factor when calculating the capture rates is the integration over the velocity distribution. In [90], parts of the integrations are performed analytically for a Gaussian veolocity distributions. These expressions are also coded in DarkSUSY for the Earth and give a more accurate calculation of the capture rate in the Earth than the approximations given above. The routine dsntcapearth2 performs these calculations for the Earth.

### 23.1.6 Accurate capture rates in the Earth for general velocity distributions

If one wants even more accurate and general expressions for the capture rates in the Sun/Earth, we have also implemented the full expressions in [90], but without assuming that the velocity distribution is a Gaussian (or Maxwell-Boltzmann). These routines are now the default in DarkSUSY.

We will here outline how these expressions look like for the Earth and how they can be used both for a Maxwell-Boltzmann distribution and for a general velocity distribution. The expressions will of course look analogously for the Sun. We start with the general case and study the special case of a Maxwell-Boltzmann distribution in the next section.

We will divide the Earth into shells and calculate the capture from element i in each shell individually. At the end we will integrate over all the shells and sum over all the elements in the Earth. The capture rate from element i per unit shell volume is given by [90][Eq. (2.8)]

$$\frac{dC_i}{dV} = \int_0^{u_{max}} du \frac{f(u)}{u} w \Omega_{v,i}^-(w)$$
(23.11)

where f(u) is the velocity distribution (normalized such that  $\int_0^\infty f(u) = n_{\chi}$  where  $n_{\chi}$  is the number density of WIMPs. The expression  $\Omega_{v,i}^-(w)$  is related to the probability that we scatter to orbits below the escape velocity. w is the velocity at the given shell and it is related to the velocity at infinity u and the escape velocity v by

$$w = \sqrt{u^2 + v^2}.$$
 (23.12)

The upper limit of integration is a priori set to  $u_{max} = \infty$ , but we will see below that due to kinematical reasons we can set it to a lower value (Eq. (23.17) below). If we allow for a form factor suppression of the form [90][Eq. (A3)]

$$|F(q^2)|^2 = \exp\left(-\frac{\Delta E}{E_0}\right) \tag{23.13}$$

with [90][Eq. (A4)]

$$E_0 = \frac{3\hbar^2}{2m_\chi R^2}$$
(23.14)

we can evaluate  $w\Omega_{v,i}^{-}(w)$  and arrive at the expression [90][Eq. (A6)]

$$w\Omega_{v,i}^{-}(w) = \sigma_{i}n_{i}\frac{\mu_{+}^{2}}{\mu}2E_{0}\left[e^{-\frac{m_{\chi}u^{2}}{2E_{0}}} - e^{-\frac{\mu}{\mu_{+}^{2}}m_{\chi}\frac{u^{2}+v^{2}}{2E_{0}}}\right]\Theta\left(\frac{\mu}{\mu_{+}^{2}} - \frac{u^{2}}{u^{2}+v^{2}}\right)$$
(23.15)

where we have introduced

$$\mu = \frac{m_{\chi}}{m_i} \quad ; \quad \mu_{\pm} = \frac{\mu \pm 1}{2} \tag{23.16}$$

with  $m_i$  the mass of element *i*. The Heaviside step function  $\Theta$  plays the role of only including WIMPs that can scatter to a velocity lower then the escape velocity *v*. To simplify our calculations we can drop this step function in Eq. (23.15) and instead set the upper limit of integration in Eq. (23.11) to

$$u_{max} = \sqrt{\frac{\mu}{\mu_-^2}} v \tag{23.17}$$

We also need the scattering cross section on element i, which can be written as [89][Eq. (9-25)]

$$\sigma_i = \sigma_p A_i^2 \frac{(m_\chi m_i)^2}{(m_\chi + m_i)^2} \frac{(m_\chi + m_p)^2}{(m_\chi m_p)^2}$$
(23.18)

where  $A_i$  is the atomic number of the element,  $m_p$  is the proton mass and  $\sigma_p$  is the scattering cross section on protons.

We now have what we need to calculate the capture rate. In Eq. (23.11) we integrate over the velocity for our chosen velocity distribution. We then integrate this equation over the radius of the Earth and sum over all the different elements in the Earth,

$$C = \int_0^{R_{\oplus}} dr \sum_i \frac{dC_i}{dV} 4\pi r^2$$
(23.19)

Note that we have not assumed anything special about our velocity distribution, it doesn't even have to be isotropic since the distribution of elements evenly in the shells will make an anisotropic distribution on average to behave as an isotropic one.

The routines that calculate the capture rates with these general (and accurate) expressions are dsntcapearthnum and dsntcapsunnum. As these calculations are somewhat time-consuming, we have also added a possibility to tabulate the result and interpolate in these tables. To use (or create, if the table files are missing) instead call dsntcapearthtab and dsntcapsuntab. These last two routines are the default in DarkSUSY. The velocity distribution used is determined by a switch when the halo model is set (i.e. when dshmset is called).

# 23.1.7 Accurate capture rates for the Earth for a Maxwell-Boltzmann velocity distribution

We will here give some more information on how the approximations introduced in the beginning of this chapter are derived from the general expressions in the preceeding section.

If the velocity distribution is of Maxwell-Boltzmann type we can greatly simplify our expressions above as we can perform the integration over velocity analytically. The integration over radius can also be further simplified by using the average mass fractions  $f_i$  and potentials  $\phi_i$  in Tables 23.1– 23.2.

If the velocity distribution in the halo is Maxwell-Boltzmann, it looks like

$$f_h(u)du = n_\chi \frac{4}{\sqrt{\pi}} \left(\frac{3}{2}\right)^{\frac{3}{2}} \frac{u^2}{\bar{v}^3} e^{-\frac{3}{2}\frac{u^2}{\bar{v}^2}} du$$
(23.20)

where  $\bar{v}$  is the three-dimensional velocity dispersion and  $n_{\chi}$  is the number density of WIMPs in the halo. However, the solar system moves through the halo with a velocity  $v_*$  and the distribution on observer with this velocity through the halo will see is

$$f_*(u) = f_h(u)e^{-\frac{3}{2}\frac{v_*^2}{\bar{v}^2}}\frac{\sinh\left(\frac{3uv_*}{\bar{v}^2}\right)}{\frac{3uv_*}{\bar{v}^2}} = n_\chi \sqrt{\frac{3}{2\pi}}\frac{u}{\bar{v}v_*} \left[e^{-\frac{3}{2}\frac{(u-v_*)^2}{\bar{v}^2}} - e^{-\frac{3}{2}\frac{(u+v_*)^2}{\bar{v}^2}}\right]$$
(23.21)

Now one would naively believe that this is not the distribution that an observer at the Earth will see. First of all, the Earth is moving with respect to the Sun and secondly, the WIMPs have gained speed by the gravitational attraction of the Sun when they reach the Earth. Both of these arguments are true and the distribution of WIMPs in the halo will not look like Eq. (23.21) to an observer on the Earth. However, Gould [94] showed that WIMPs from the halo can diffuse into the solar system due to gravitational interactions with the planets and this distribution of WIMPs will roughly look like as if the Earth was in free space moving through the halo with the velocity of the solar system, i.e. Eq. (23.21). We will later scrutinize this statement, as it turns out that it does not quite hold, but as a first guess it is a reasonable approximation. For the Sun, though, the velocity distribution give above is the correct one for a Maxwell-Boltzmann distribution.

With the distribution Eq. (23.21) we can analytically perform the integration over velocity in Eq. (23.11). After some algebra we arrive at [90][Eq. (A10)]

$$\frac{dC_i}{dV} = \left(\frac{8}{3\pi}\right)^{\frac{1}{2}} \frac{\sigma_i n_i n_\chi \bar{v}}{2b\eta} \\
\left[\frac{e^{-a\hat{\eta}^2}}{\sqrt{1+a}} \left[2\widetilde{\mathrm{erf}}(\hat{\eta}) - \widetilde{\mathrm{erf}}(\hat{A}_+) + \widetilde{\mathrm{erf}}(\hat{A}_-)\right] \\
-\frac{e^{-b\tilde{\eta}^2}}{\sqrt{1+b}} e^{-(a-b)A^2} \left[2\widetilde{\mathrm{erf}}(\tilde{\eta}) - \widetilde{\mathrm{erf}}(\check{A}_+) + \widetilde{\mathrm{erf}}(\check{A}_-)\right]\right]$$
(23.22)

where erf is the modified error function,

$$\widetilde{\operatorname{erf}}(x) = \frac{\sqrt{\pi}}{2} \operatorname{erf}(x) \qquad ; \qquad \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-y^2} dy. \tag{23.23}$$

Following Gould [90], we have in Eq. (23.22) introduced the following shorthand notation:

$$\eta = \sqrt{\frac{3}{2}} \frac{v_{\pm}^{2}}{v^{2}} ; \quad a = \frac{m_{c}hi\bar{v}^{2}}{3E_{0}} ; \quad b = \frac{\mu}{\mu_{+}^{2}}a \hat{\eta} = \frac{\eta}{\sqrt{1+a}} ; \quad \check{\eta} = \frac{\eta}{\sqrt{1+b}} A^{2} = \frac{3}{2} \frac{v^{2}}{\bar{v}^{2}} \frac{\mu}{\mu_{-}^{2}} ; \quad \hat{A} = A\sqrt{1+a} ; \quad \check{A} = A\sqrt{1+b} \hat{A}_{\pm} = \hat{A} \pm \hat{\eta} ; \quad \check{A}_{\pm} = \check{A} \pm \check{\eta}$$

$$(23.24)$$

If we wish, we can now integrate Eq. (23.22) over radius just like in the previous section, but we can without loosing too much accuracy, replace this integration with a sum over the elements in the Earth with their respective typical location. I.e. we can write

$$C = \sum_{i} \frac{dC_i}{dV} \frac{1}{n_i} \frac{f_i M_{\oplus}}{m_i}$$
(23.25)

where we instead of the number density  $n_i$  use the total number of atoms of the given type  $f_i M_{\oplus}/m_i$ . Note that for each element in the sum we should evaluate this expression at the given typical gravitational potential  $\phi_i$  of the element, i.e. with the escape velocity given by Eq. (23.10). The mass fractions  $f_i$  and typical potentials  $\phi_i$  are listed in Table 23.2 (and analogously in Table 23.1 for the Sun). This approximation introduces an error of no more than about 1–2% for a Maxwell-Boltzmann distribution<sup>\*</sup>

The capture rate evaluated with the expressions shown here are encoded into the routine dsnt-capearth2. Note that we have not coded the corresponding approximate expressions for the Sun. Instead, as given in the preceeding section, we now have more accurate expressions for both the Sun and the Earth.

#### 23.1.8 Effects of WIMP diffusion in the solar system

As the Earth has a rather low escape velocity, the Earth will only be able to capture WIMPs that have a rather low velocity with respect to the Earth. However, WIMPs from the halo have gained speed in the gravitational potential from the Sun and will essentially be impossible to capture by the Earth. Hence, the Earth will only capture WIMPs that have diffused around in the solar system (by gravitational interactions with the other planets). Gould showed [94] that effectively this diffusion will lead to the same phase space distribution at the Earth as if the Earth was in free space (i.e. neglecting the solar potential). However, numerical simulations of asteroids showed that they are thrown into the Sun due to perturbations of the orbits by other planets, see e.g. [95]. These analyses led to worried that maybe the population of WIMPs diffusing around in the solar system is not as big as thought [96]. In [97], Lundberg and Edsjö investigated this issue with detailed numerical simulations of WIMP orbits in the solar system, showing that the annihilation rate in the Earth is typically reduced by up to two orders of magnitude. In DarkSUSY, we include these results for the neutrino rates from the Earth by using the velocity distribution at the Earth (as obtained in [97]). This velocity distribution is then used as input for our numerical capture rate routines instead of the usual approximation of using the halo velocity distribution directly. Using these new velocity distributions for the Earth is the default in DarkSUSY.

<sup>\*</sup>Note that it is not advisable to use this approximation for general velocity distributions. If one e.g. has a lower limit on possible velocities,  $u_{min}$ , for heavy WIMPs capture will then only be possible very close to the central core. Replacing the actual distribution of potentials  $\phi(r)$  with the typical value  $\phi_i$  may then introduce larger errors. We will encounter these kind of distributions shortly.

### 23.2 Neutrinos from Sun and Earth – routines

This folder contains routines to calculate the neutrino-induced muon flux from the Earth and the Sun in various models.

There are several different methods of calculation available (determined by secalcmet in dssecom.h). Method 1 uses the approximate formulae for the capture rates in the Earth/Sun from the Jungman, Kamionkowski and Griest review [89]. Method 2, uses the same expression for the Sun, but the full expression from Gould [90] for capture in the Earth. Method 4 uses a full numerical integration over the velocity distribution (instead of assuming that it is Gaussian) and method 5, finally, also performs a full numerical integration over the momentum transfer in the form factors (instead of assuming exponential form factors). The default is to use method 5, but instead of using the numerical routines directly, tables are used. One can always force using the numerical routines instead of the tables if one so wishes. There are several options regarding how many elements to incluce in the Sun summation over elements ('lo', 'med' and 'hi'). The easiest way to select method is by calling dssenu\_set. For the Earth, method 5 is not expected to make a large difference due to the smaller momentum transfers in the Earth and method 5 reverts back to method 4 for the Earth.

A call to dssenu\_set('default') is made in dsinit, but can be changed by the user by calling dssenu\_set after dsinit.

To calculate the neutrino-induced muon flux from the Earth, you call **dssenu** rates.

# se\_yield: Yields from annihilation in the Sun/Earth

#### (JE: I am responsible for updating this part)

### 24.1 Muon yields from annihilation in the Earth/Sun – theory

We need to take into account all processes that yield muon neutrinos from annihilation in the Earth/Sun. To do this, we use a Monte Carlo, WimpSim [98], to simulate annihilations in the center of the Sun/Earth, neutrino oscillations and neutrino interactions on the way out of the Sun/Earth and to the detector.

#### 24.1.1 Monte Carlo simulations with WimpSim

We need to evaluate the yield of different particles per neutralino annihilation. The hadronization and/or decay of the annihilation products are simulated with PYTHIA [10] 6.426 and we here describe how the simulations are done. For annihilation in the Sun/Earth the simulations are done for a set of 22 neutralino masses,  $m_{\chi} = 3$ , 6, 10, 25, 50, 80.3, 91.2, 100, 150, 176, 200, 250, 350, 500, 750, 1000, 1500, 2000, 3000, 5000, 7500 and 10000 GeV. We tabulate the yields and then interpolate these tables in DarkSUSY.

We are mainly interested in the flux of high energy muon neutrinos and neutrino-induced muons at a neutrino telescope. We simulate 13 'fundamental' annihilation channels, for each mass (where kinematically allowed) above. In Table 24.1 we list the 'fundamental' channels for which simulations are run and the full set of more complex channels. Pions and kaons get stopped before they decay and are thus made stable in the PYTHIA simulations so that they don't produce any neutrinos. For annihilation channels containing Higgs bosons, we can calculate the yield from these fundamental channels by letting the Higgs bosons decaying in flight (see below). We also take into account the energy losses of *B*-mesons in the Sun and the Earth by following the approximate treatment of [81] but with updated *B*-meson interaction cross sections as given in [99]. We also take neutrinointeractions on the way out of the Sun into account by considering the charged-current interaction as a neutrino-loss and the neutral current interactions are simulated with nusigma [100]. The neutrino-nucleon charged current interactions close to the detector are also simulated with nusigma and finally the multiple Coulomb scattering of the muon on its way to the detector is calculated using distributions from [101]. We have used the CTEQ6 structure functions in these simulations. We also take into account neutrino oscillations with a full three-neutrino Monte Carlo. All of these processes are put into the simulation package WimpSim [98] that can be downloaded separately. Results from simulation runs with this package are included with DarkSUSY. For more details on these simulations, see [102].

71

For each annihilation channel and mass we simulate  $10^7$  annihilations and tabulate the final results as a neutrino-yield, neutrino-to-lepton conversion rate, a muon yield and hadronic shower yields differential in energy and angle from the center of the Sun/Earth. We also tabulate the integrated yield above a given threshold and below an opening angle  $\theta$ . We assumed throughout that the surrounding medium is water with a density of  $1.0 \text{ g/cm}^3$ . Hence, the neutrino-to-muon conversion rates have to be multiplied by the density of the medium. In the muon fluxes, the density cancels out (to within a few percent). All results are summarized as yield tables that can be loaded and interpolated in with DarkSUSYThis is done with the function dswayieldf. There are three kinds of yields (two-dimensional in opening angle  $\theta$  and energy), kind=1 gives integrated vields, kind=2 gives differential vields and kind=3 gives vields integrated in angle, but differential in  $\theta$ . For each kind, there are 26 different types of yield available according to Table 24.4. As a default, only type 3-4, 9-10 and 13-14 are included in the DarkSUSY download, as these are the most commonly used types. If you need any other types, download the auxiliary data files from http://www.darksusy.org, unpack them in share/DarkSUSY in the DarkSUSY root directory, and then do a usual configure and make to install them. Also note that the kind=3 yields are not tabulated directly, but are instead calculated and tabulated when the simulation tables are read in during DarkSUSY initialization.

With these simulations, we can calculate the yield for any of these particles for a given MSSM model. For the Higgs bosons, which decay in flight, an integration over the angle of the decay products with respect to the direction of the Higgs boson is performed. Given the branching ratios for different annihilation channels it is then straightforward to compute the yield above any given energy threshold and within any angular region around the Sun or the center of the Earth. The routine dswayieldone calculates the yield for one channel, i.e. even these complex channels containing Higgs bosons, whereas the main routine dswayield calculates the total yield for a given model. Note that the WIMP annihilation yield routines do not know about SUSY at all, so before they are called, a routine dswasetup is called to set up the annihilation branching ratios for the WIMP and decay channels for the Higgs bosons. In Tables 24.2 and 24.3, the Higgs decay width channels are given. If these routines are used with other particle physics models, replace dswasetup with a routine appropriate for your particle physics model and then call dswayield as usual.

Annihilation channel		Internal channel	Internal channel array index
ch	Particles	chi	chii
1	$S_{1}^{0}S_{1}^{0}$	-	-
2	$S_{1}^{0}S_{2}^{0}$	-	-
3	$S_{2}^{0}S_{2}^{0}$	-	-
4	$S_{3}^{0}S_{3}^{0}$	-	-
5	$S_{1}^{0}S_{3}^{0}$	-	-
6	$S_{2}^{0}S_{3}^{0}$	-	-
7	$S^-S^+$	-	-
8	$Z^{0}S_{1}^{0}$	-	-
9	$Z^{0}S_{2}^{0}$	-	-
10	$Z^{0}S_{3}^{0}$	-	-
11	$W^{-}S^{+}/W^{+}W^{-}$	-	-
12	$Z^0Z^0$	9	9
13	$W^+W^-$	8	8
14	$ u_e \bar{ u}_e$	12	11
15	$e^+e^-$	-	-
16	$ u_{\mu} ar{ u}_{\mu}$	13	12
17	$\mu^+\mu^-$	10	-
18	$ u_{ au} \overline{ u}_{ au}$	14	13
19	$\tau^+\tau^-$	11	10
20	$u \bar{u}$	2	2
21	$d\overline{d}$	1	1
22	$c\bar{c}$	4	4
23	$s\bar{s}$	3	3
24	$t\bar{t}$	6	6
25	$bar{b}$	5	5
26	gg	7	7
27	qqg	-	-
28	$\gamma\gamma$	-	-
29	$Z^0\gamma$	-	-

Table 24.1: The annihilation channels ch used in dswayieldone. Also shown are the internal channel numbers chi used for the fundamental channels used in the simulations (used by routine dswayieldf). To save some additional space with the data files in memory, there are also array index channel numbers chi that are only used internally to access the right elements of the yield arrays. S denotes scalars (Higgs bosons).

Decay width channel	
dch	Particles
1-29	Same as the annihilation channels in Table 24.1.
30	Sfermions
31	Neutralinos
32	Charginos

Table 24.2: The neutral scalar (Higgs) decay width channels used. In DarkSUSY these are stored in the array hdwidth(i,j) where i is the decay channel index above and j is the Higgs number (1–3 for  $H_1^0$ ,  $H_2^0$  and  $H_3^0$  respectively). For the wa routines, these decay branching ratios (partial width divided by total width) are stored in dswas0br(i,j).
Decay width channel	
dch	Particles
1	ud
2	$u\bar{s}$
3	$uar{b}$
4	$c ar{d}$
5	$c\bar{s}$
6	$car{b}$
7	$t \bar{d}$
8	$t\bar{s}$
9	$tar{b}$
10	$\nu_e e^+$
11	$ u_{\mu}\mu^+$
12	$ u_{ au}  au^+$
13	$W^{+}S_{1}^{0}$
14	$W^{+}S_{2}^{0}$
15	$W^{+}S_{3}^{0}$
20	Sfermions
21	Neutralinos and charginos

Table 24.3: The (positively) charged scalar (Higgs) decay width channels used. In DarkSUSY these are stored in the array hdwidth(i,4) where i is the decay channel index above. For the wa routines, these decay branching ratios (partial width divided by total width) are stored in dswascbr(i).

Yield type		
type	Yield	Unit
1	$\nu_e$	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
2	$\bar{ u}_e$	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
3	$ u_{\mu}$	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
4	$ar{ u}_{\mu}$	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
5	$ u_{ au}$	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
6	$ar{ u}_{ au}$	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
7	$e^-$ at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3} \text{ annihilation}^{-1}$
8	$e^+$ at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3} \text{ annihilation}^{-1}$
9	$\mu^-$ at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
10	$\mu^+$ at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
11	$\tau^-$ at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
12	$\tau^+$ at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3} \text{ annihilation}^{-1}$
13	$\mu^-$ at an imaginary plane at detector	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
14	$\mu^+$ at an imaginary plane at detector	$10^{-30} \text{ m}^{-2} \text{ annihilation}^{-1}$
15	hadronic shower from $\nu_e$ CC int. at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3} \text{ annihilation}^{-1}$
16	hadronic shower from $\bar{\nu}_e$ CC int. at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3} \text{ annihilation}^{-1}$
17	hadronic shower from $\nu_{\mu}$ CC int. at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
18	hadronic shower from $\bar{\nu}_{\mu}$ CC int. at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
19	hadronic shower from $\nu_{\tau}$ CC int. at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
20	hadronic shower from $\bar{\nu}_{\tau}$ CC int. at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
21	hadronic shower from $\nu_e$ NC int. at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
22	hadronic shower from $\bar{\nu}_e$ NC int. at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
23	hadronic shower from $\nu_{\mu}$ NC int. at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3} \text{ annihilation}^{-1}$
24	hadronic shower from $\bar{\nu}_{\mu}$ NC int. at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3}$ annihilation <sup>-1</sup>
25	hadronic shower from $\nu_{\tau}$ NC int. at neutrino-nucleon vertex	$10^{-30}$ m <sup>-3</sup> annihilation <sup>-1</sup>
26	hadronic shower from $\bar{\nu}_{\tau}$ NC int. at neutrino-nucleon vertex	$10^{-30} \text{ m}^{-3} \text{ annihilation}^{-1}$

Table 24.4: The yield types available from the wa routines. All of these yields are at the detector (currently IceCube). Note that the units are for integrated yields (kind=1), for differential yields (kind=2), the units should be multiplied by  $\text{GeV}^{-1}$  degree<sup>-1</sup>. CC int. = charged current interactions. NC int. = neutral current interactions.

# si: Self-interactions

### 25.1 Dark matter self-interactions (si) – theory

The CDM paradigm, resting on cold and collisionless dark matter, describes cosmological structure formation with remarkable accuracy at scales larger than about one Mpc. At smaller cosmological scales, on the other hand, this paradigm is less well tested, and currently still allows for DM self-interactions as strong as the interaction between nucleons – and thus much stronger than current limits on DM interacting with SM particles. The possibility that DM could be relatively strongly interacting with itself [103] and thereby leave imprints on cosmological observables related to structure formation has therefore seen significant interest in recent years, both from an astrophysical and a model-building perspective. Self-interacting DM (SIDM) could even address [104, 105, 106, 107, 108, 109, 110, 111] the most pressing potential small-scale problems of ACDM cosmology [112], referred to as 'core-vs.-cusp" [113, 114], 'too-big-to-fail' [115, 116], 'diversity' [117, 118] and 'missing satellites' problems [119, 120] (the latter problem is addressed for SIDM with late kinetic decoupling, which is a natural combination in many models [55]). Those specific discrepancies with the ACDM paradigm may of course well turn out to be either due to poorly modelled baryonic effects or observational uncertainties – but these examples serve as evidence that SIDM can leave observable imprints even in the absence of interactions with the SM. An unambiguous detection of such signatures, which are not expected in for example traditional WIMP models of DM, would significantly narrow down the range of possible particle explanations for the nature of DM.

The traditional reference quantity for the impact of DM self-interactions on the halo structure is the momentum transfer cross section,

$$\sigma_T \equiv \int d\Omega \left(1 - \cos\theta\right) \frac{d\sigma}{d\Omega} \,, \tag{25.1}$$

where  $\sigma$  is the standard cross section for DM-DM scattering. This has the advantage of regulating large forward-scattering amplitudes, which should not affect the DM distribution (the same goes for backward scattering which, however, is treated symmetrically in this prescription. See Ref. [121, 122] for a more detailed discussion). The size of this quantity that is of cosmological relevance is very roughly given by

$$\sigma_T/m_\chi \sim 1 \,\mathrm{cm}^2/\mathrm{g}\,. \tag{25.2}$$

Cross sections in this ballpark, in other words, may leave observable imprints and possibly address the various ACD; small-scale structure problems mentioned above, while much smaller cross sections have no impact on the structure of DM halos. Much larger values of  $\sigma_T/m_{\chi}$  are ruled out, on the other hand, in particular from observations of galaxy clusters. For a detailed review that summarizes both various constraints on DM self-interactions and models that have been discussed in the literature, we refer to Ref. [123].

The details of the DM self-interactions depend on the underlying particle theory. A simple contact interaction term in the Lagrangian, for example, would lead to a constant (velocity-independent)  $\sigma_T$ . Another well-motivated option is that the interaction is mediated by a massive particle with mass  $m_{\rm med}$ , which leads to a Yukawa potential between the two DM particles:

$$V(r) = \pm \frac{\alpha_{\chi}}{r} e^{-m_{\rm med}r} \,. \tag{25.3}$$

Here,  $\alpha_{\chi} = g_{\chi}^2/(4\pi)$  describes the coupling strength between mediator and DM particles, and the different signs refer to attractive (-) and repulsive (+) potentials. Scalar mediators only generate attractive potentials, while for vector mediators this depends on the particles involved in the scattering: for (e.g. Dirac) DM scattering with anti-DM, the force is attractive, otherwise it is repulsive. An interesting phenomenological aspect for the scattering of nonrelativistic particles in such a potential is the resulting strong velocity-dependence of  $\sigma_T$ ; this allows to achieve large scattering cross sections for the relatively small velocities of ~30 km/s typically encountered in dwarf galaxies (where one observes potential discrepancies with the  $\Lambda$ CDM expectations) while evading the strong bounds at cluster scales for velocities of ~1000 km/s. The transfer cross section resulting from scattering in a Yukawa potential has been extensively studied, and depends on the scattering regime ( $v_{\rm rel}$  is the relative velocity between the DM particles):

• In the **Born regime** ( $\alpha_{\chi} m_{\chi} \lesssim m_{\text{med}}$ ),  $\sigma_T$  can be calculated perturbatively, leading (for both attractive and repulsive potentials) to [124]

$$\sigma_T^{\text{Born}} = \frac{8\pi \alpha_\chi^2}{m_\chi^2 v_{\text{rel}}^4} \left( \ln \left[ 1 + \frac{m_\chi^2 v_{\text{rel}}^2}{m_{\text{med}}^2} \right] - \frac{m_\chi^2 v_{\text{rel}}^2}{m_{\text{med}}^2 + m_\chi^2 v_{\text{rel}}^2} \right) \,. \tag{25.4}$$

• In the classical regime  $(m_{\chi} v_{\rm rel} \gtrsim m_{\rm med})$ , a large numbers of partial waves contributes such that non-perturbative effects are important. In principle, this can be computed by numerically solving the Schrödinger equation for each case and then summing all contributions [121]. A computationally much more efficient method is to use parameterizations [125] of numerical results from the Plasma literature that have been obtained for screened Coulomb scattering [126, 127]. For an attractive potential, these are given by

$$\sigma_T^- = \frac{\pi}{m_{\rm med}^2} \times \begin{cases} 2\beta^2 \ln[1+\beta^{-2}] & \text{for } \beta \lesssim 10^{-2} \\ \frac{7\beta^{1.8}+1960(\beta/10)^{10.3}}{1+1.4\beta+0.006\beta^4+160(\beta/10)^{10}} & \text{for } 10^{-2} \lesssim \beta \lesssim 10^2 \\ 0.81 \left(1+\ln\beta-(2\ln\beta)^{-1}\right)^2 & \text{for } \beta \gtrsim 10^2 \end{cases}$$
(25.5)

where  $\beta \equiv 2\alpha_{\chi} m_{\rm med} / (m_{\chi} v_{\rm rel}^2)$ , while for a repulsive potential we have

$$\sigma_T^+ = \frac{\pi}{m_{\rm med}^2} \times \begin{cases} 2\beta^2 \ln[1+\beta^{-2}] & \text{for } \beta \lesssim 10^{-2} \\ \frac{8\beta^{1.8}}{1+5\beta^{0.9}+0.85\beta^{1.6}} & \text{for } 10^{-2} \lesssim \beta \lesssim 10^4 \\ (\ln 2\beta - \ln \ln 2\beta)^2 & \text{for } \beta \gtrsim 10^4 \end{cases}$$
(25.6)

• Finally, there is the **resonant regime** (for  $m_{\chi}v_{\rm rel} \leq m_{\rm med}$  or  $\alpha_{\chi}m_{\chi} \gtrsim m_{\rm med}$ ). Full numerical solutions can be obtained by solving the Schrödinger equation explicitly in this regime. These are well described by the following analytic expressions that result from approximating the Yukawa potential with a Hulthén potential [121]

$$\sigma_T^{\text{Hulthn}} = \frac{16\pi}{m_\chi v_{\text{rel}}^2} \sin^2 \left( \text{Arg} \left[ \frac{i\Gamma[i\Theta v_{\text{rel}}]}{\Gamma[\lambda_+]\,\Gamma[\lambda_-]} \right] \right) \,, \tag{25.7}$$

where  $\Theta \equiv \frac{m_{\chi}}{\sqrt{2\zeta(3)}m_{\text{med}}}$  and  $\lambda_{\pm} \equiv 1 + i\Theta v_{\text{rel}}/2 \pm \sqrt{\alpha_{\chi}\Theta - \Theta^2 v_{\text{rel}}^2/4}$  for an attractive potential and  $\lambda_{\pm} \equiv 1 + i\Theta v_{\text{rel}}/2 \pm i\sqrt{\alpha_{\chi}\Theta + \Theta^2 v_{\text{rel}}^2/4}$  for a repulsive potential.

#### 25.2 Self-interactions – routines

In DarkSUSY,  $\sigma_T(v_{\rm rel})/m_{\chi}$  is provided by an interface function dssigtm returned by the particle module which, from the perspective of the core library, can take any functional form. The function dssigtmav then computes the velocity average  $\langle \sigma_T \rangle/m_{\chi}$ , assuming a Maxwellian velocity distribution of the DM particles; this gives a better estimate of the effect of DM self-interactions than just evaluating dssigtm directly for a typical halo velocity. The core library furthermore provides several auxiliary routines, to be used by any particle module, for the commonly encountered specific transfer cross sections in the presence of a Yukawa potential as discussed above. In particular, dssisigmatborn returns the expression given in Eq. (25.4), dssisigmatclassical those given in Eqs. (25.5, 25.6), and dssisigmatres those given in Eq. (25.7). For the latter, we adopt the complex gamma function as implemented in the CERN library (based on Ref. [128]), and use analytic expansions where a naive usage of these routines is problematic due to limited numerical precision (which is relevant for a significant fraction of the physically interesting parameter range).

# Part III

# Particle physics modules in src\_models

# Basic principles and common routines

The general concept of a particle physics module, and how it communicates with the core library via interface functions, was already introduced in Chapter 3 – see in particular Section 3.2, and Table 3.3 for an automatically updated list of all interface functions that the core library is aware of. Note that there is no *principal* restriction on which interface functions a particle module must provide: the main program will determine at compilation time whether it needs a functionality of the core library that requires certain interface functions to exist.

Every particle physics module must provide an independent representation of the full particle content of the respective particle theory. How this is done is fully flexible, and completely up to the module. In practice, however, there are common frameworks, like the Standard Model, that appear repeatedly. For convenience, we therefore store auxiliary routines and setups that *may* be used by more than just one particle module in src\_models/common. Common blocks and header files included by more than one particle module are found in src\_model/include. In order to keep up the modularity of the code, routines in src\_models/common thus *only* include files in src\_model/include.

### 26.1 common/aux: auxiliary routines

Here we collect various routines that belong to particle physics, and hence do not reside in ds\_core, but are not only useful for one specific particle module. Currently, the most important are

- A set of functions to handle *unique model IDs* for each particle model, which is essential in order not to repeat identical calculations when changing the model and thus to optimize numerical performance. A new such model ID should be assigned with dsnewidnumber whenever a new particle model is initialized (for the modules provided with the DarkSUSY release, this is automatically done in dsmodelsetup). Any routine in src\_models can test with dsidnumberset whether this ID number is indeed set, and retrieve its value with dsidnumber (which then can be compared to the locally stored value of the ID number that was valid at the time when the respective routine was called the previous time).
- The function dsanthreshold returns the correction factor to a 2-body rate close to a kinematical threshold, resulting from the fact that one or both of the particles may be slightly off-shell. This implements the simplified treatment presented in Ref. [27], c.f. their Fig. 5, and hence assumes that the decay products of the virtual particle are (effectively) massless.

#### 26.2 common/sm: standard model

The routines collected in this folder provide a convenient shortcut to include the most basic properties of the standard model in a BSM particle module. Currently, the most important functionalities collected here are given by

- An initialization routine dsinit\_sm, which can be called directly from the corresponding dsinit\_module. After a call to this routine, the functions dsmass(kPDG) and dswidth(kPDG) correctly return masses and widths of the standard model particles kPDG being the PDG code [101] as stored centrally in src\_models/include/dssmparam.h.
- Running quark masses are provided up to the 4-loop level, with SM contributions only, both for pole and  $\overline{MS}$  masses [129].
- The SM contributions to the strong coupling constant are also provided up to the 4-loop level.
- Another convenience function is dsgf2s2thw, which calculates  $\sin^2(\theta_W)$  at the  $m_Z$  scale: in praxis, its most important application is to ensure that a particle module can adopt a consistent relation between  $m_Z$  and  $m_W$ , which for example can be crucial in order to keep large numerical cancellations under control.

Let us stress that standard model physics in DarkSUSY is by far not restricted to the routines collected in common/sm. Much is presently still contained in the mssm module. It will be (further) disentangled from the SUSY-specific parts as new modules are added to the code that require this functionality.

# Chapter 27 The empty model

The empty module cannot be used to perform any real calculations. Its purpose is for debugging and testing only, with the main advantage being that it contains (empty versions of) all interface functions that the core library is aware of (for a full list, see Table 3.3). In some cases, this makes it a good starting point to create new particle physics modules.

# Generic decaying dark matter

This module provides a simple phenomenological template for a generic decaying dark matter model. It is fully specified by the DM mass  $m_{\chi}$  and the total decay rate  $\Gamma$ , along with a list of decay channels. The latter are specified by their branching ratios and the PDG codes of the final state standard model particles. Non-standard model final states can also be included, and are treated as invisible decays.

The phenomenology of this module is restricted to indirect DM searches, with full access to the corresponding routines in ds\_core. DM scattering with nuclei is assumed to be negligible, as well thermal production of DM in the early universe.

# Generic WIMPs

Similar to the case of generic\_decayingDM, the module generic\_wimp provides a simple phenomenological template for a large class of DM candidates. Rather than being based on an actual particle physics model, it mostly serves to provide an illustration of how the functionalities of DarkSUSY can be used in phenomenological studies of âĂŸvanillaâĂŹ WIMP DM, when only providing the absolute minimum of input parameters.

A generic WIMP model in DarkSUSY is set up by a call to dsgivemodel\_generic\_wimp, and hence fully defined by the input parameters of that routine: the mass  $m_{\chi}$  of the DM particle and a flag stating whether the DM particle is its own anti-particle or not; a constant annihilation rate  $\sigma v$  in the CMS frame, along with the dominant annihilation channel into SM particles (as usual stated in terms of PDG codes); finally, the spin-independent scattering cross section <sub>SI</sub> of DM with nucleons.

This simple setup allows to use essentially the full functionality of the core library, from relic density calculation to direct detection rates and cosmic ray propagation and indirect detection signals in general.

# The Minimal Supersymmetric Standard Model

The implementation of the minimal supersymmetric standard model (MSSM) in the module mssm mainly follows that of DarkSUSY versions 4 [1] (see also Section 30.8 in this manual). In particular, the conventions for the superpotential and soft supersymmetry-breaking potential are the same as implemented in [21], and thus similar to [130, 131]. The full set of input parameters to be provided at the weak scale thus consists of the pseudoscalar mass  $(m_A)$ , the ratio of Higgs vacuum expectation values  $(\tan \beta)$ , the Higgsino  $(\mu)$  and gaugino  $(M_1, M_2, M_3)$  mass parameters, trilinear couplings  $(A_{Eaa}, A_{Uaa}, A_{Daa}, \text{ with } a = 1, 2, 3)$  as well as soft sfermion masses  $(M_{Qaa}^2, M_{Laa}^2, M_{Uaa}^2, M_{Daa}^2, M_{Daa}^2)$  $M_{Eaa}^2$ , with a = 1, 2, 3.\* Internally, those values are stored in mssm common blocks. The user may either provide them directly or by setting up pre-defined phenomenological MSSM models with a reduced number of parameters through a call to a routine like dsgive model or dsgive model25 (followed by a call to dsmodelsetup). The former sets up the simplest of those models, defined by the input parameters  $\mu$ ,  $M_2$ ,  $m_A$ , tan  $\beta$ , a common scalar mass  $m_0$ , and trilinear parameters  $A_t$  and  $A_b$ ;  $M_1$  and  $M_3$  are then calculated by assuming the GUT condition, and the remaining MSSM parameters are given by  $\mathbf{M}_Q = \mathbf{M}_U = \mathbf{M}_D = \mathbf{M}_E = \mathbf{M}_L = m_0 \mathbf{1}, \ \mathbf{A}_U = \text{diag}(0, 0, A_t),$  $\mathbf{A}_D = \text{diag}(0, 0, A_b), \mathbf{A}_E = \mathbf{0}$ . Similarly, dsgive model25 sets up a pMSSM model with 25 free parameters (see the header of that file for details). Alternatively, all those values can be set by reading in an SLHA file, or providing GUT scale parameters in the case of cMSSM models (via an interface to the ISASUGRA code, as included in ISAJET [132, 133]).

Compared to previous versions of the code, DarkSUSY 6 has a new interface to read and write SUSY Les Houches Accord (SLHA) files [134, 135].

All particle and sparticle masses are stored in a common block array mass(). For neutralino masses, we include the leading loop corrections [136, 137, 138] but neglect the relatively small corrections for charginos [136] (in both cases, masses cannot be negative in our convention).<sup>†</sup> Likewise, all mixing matrices and decay widths are available as common block arrays. The latter are currently only computed for the Higgs particles (via an interface to the FeynHiggs [139, 140, 141, 142, 143] package), while the other sparticles have fictitious widths of 0.5% of the sparticle mass (for the sole purpose of regularizing annihilation amplitudes close to poles). Again, the conventions for masses and mixings follow exactly those of Ref. [1], to which we refer for further details.

<sup>\*</sup> Note that currently only diagonal matrices are allowed. While not being the most general ansatz possible, this implies the absence of flavour changing neutral currents at tree-level in all sectors of the model.

 $<sup>^{\</sup>dagger}$  Unless, of course, those values are provided by an SLHA file. This comment also applies to the following simplifications concerning both sparticle masses and widths.

# 30.1 mssm/ac: Accelerator constraints

#### **30.1.1** Accelerator bounds

DarkSUSY contains a set of routines for a rough check whether a given model is excluded by accelerator constraints. These routines are called **dsacbnd[number]**. The policy is that when we update DarkSUSY with new accelerator constraints, we keep the old routine, and add a new routine with the last number incremented by one. Which routine that is called is determined by calling **dsacset** with a tag determining which routine to call. To check the accelerator constraints, then call **dsacbnd** which calls the right routine for you. Upon return, **dsacbnd** returns an exclusion flag, warning. If zero, the model is OK, if non-zero, the model is likely excluded. The cause for the exclusion is coded in the bits of excl according to table 30.1.

We stress that these bounds are in most cases only *approximate* limits: DarkSUSY generally focusses on theoretical predictions for such observables, given a DM model realization, rather than on the implementation of experimental likelihoods and the possibility to derive statistically well-defined limits from those. For the latter, we instead refer to packages like DarkBit [144] (or ColliderBit [145] for accelerator-based constraints)

	excl		
Bit set	Octal value	Decimal value	Reason for exclusion
0	1	1	Chargino mass
1	2	2	Gluino mass
2	4	4	Squark mass
3	10	8	Slepton mass
4	20	16	Invisible $Z$ width
5	40	32	Higgs mass in excluded region
6	100	64	Neutralino mass
7	200	128	$b  ightarrow s \gamma$
8	400	256	$\rho$ parameter
9	1000	512	$(g-2)_{\mu}$
10	2000	1024	$B_s \rightarrow \mu^+ \mu^-$
11	4000	2048	squark-gluino
12	10000	4096	Higgs mass does not fit observed Higgs

Table 30.1: The bits of excl are set to indicate by which process this particular model is excluded. Check if a bit is set with btest(excl,bit).

Compared to previous DarkSUSY versions, we use in particular updated limits from HiggsBounds [146] on the mass of the MSSM Higgs bosons, as well as approximate bounds on squark and gluino masses from LHC 8 TeV data [147]. For  $b \to s\gamma$ , we keep our genuine routines in mssm/ac\_bsg for this rare decay (see Ref. [1] for a more detailed description) but now use as a default the result from Superlso [148]; we compare this to the current limit of  $\mathcal{B}(B \to X_s\gamma) = (3.27\pm0.14)\times10^{-4}$  as adopted in FlavBit [149], based on data from BarBar and Belle [150, 151, 152]. Superlso also computes the rate for the rare leptonic decay  $B_s^0 \to \mu^+\mu^-$ , which we compare to the LHCb measurement of  $\mathcal{B}(B_s^0 \to \mu^+\mu^-) = (3.0 \pm 0.6^{+0.3}_{-0.2}) \times 10^{-9}$  [153]. Finally,  $a_{\mu} \equiv (g-2)_{\mu}/2$  is calculated by dsgm2muon, based on [154]; in dsacbnd, this is compared to the observed valued of  $a_{\mu, obs} = (11659208.9 \pm 6.3) \times 10^{-10}$  [155] after subtracting the SM expectation as specified in PrecisionBit [156].

# 30.2 mssm/an: (Co-)annihilation cross sections

#### **30.2.1** Annihilation cross sections – theory

For the relic density calculations, we need all possible (co)annihilation cross sections between neutralinos, charginos and sfermions.

#### 30.2.1.1 Annihilation cross sections

We have calculated all two-body final state cross sections at tree level involving neutralinos, charginos, sneutrinos, sleptons and squarks in the initial state. A complete list is given below.

Since we have so many different diagrams contributing, we have to use some method where the diagrams can be calculated efficiently. To achive this, we calculate the diagrams with general expressions for vertices, masses etc so that they can be reused for other processes. How we do this in practice differs a bit between different sets of annihilation diagrams.

For neutralino-neutralino, neutralino-chargino and chargino-chargino annihilation, we classify the diagrams according to their topology (s-, t- or u-channel) and to the spin of the particles involved. We then compute the helicity amplitudes for each type of diagram analytically with REDUCE [157] using general expressions for the vertex couplings.

The strength of the helicity amplitude method is that the analytical calculation of a given type of diagram has to be performed only once and the sum of the contributing diagrams for each set of initial and final states can be done numerically afterwards.

For the diagrams involving sfermions, FORM is used to analytically calculate the amplitudes. This output is then converted into Fortran with a PERL script, **form2f** [158].

#### 30.2.1.2 Coannihilation diagrams

All Feynman diagrams for which we calculate the annihilation cross section are listed in the coming sections. s(x), t(x) and u(x) denote a tree-level Feynman diagram in which particle x is exchanged in the s-, t- and u-channel respectively.

The convention used in this list of included coannihilation diagrams is that if a sfermion is denoted  $\tilde{f}$ , then it's antiparticle is denoted  $\tilde{f}^*$ .

#### 30.2.1.3 Neutralino and chargino annihilation

Indices i, j, k run from 1 to 4, and indices c, d, e from 1 to 2.  $u, \tilde{u}, d, \tilde{d}, \nu, \tilde{\nu}, \ell, \tilde{\ell}, \tilde{f}$  and  $\tilde{f}$  are generic notations for up-type quarks, up-type squarks, down-type quarks, down-type squarks, neutrinos, sneutrinos, leptons, sleptons, fermions and sfermions. A sum of diagrams over (s)fermion generation indices and over the neutralino and chargino indices k and e is understood (no sum over indices i, j, c, d).

#### Neutralino-neutralino annihilation

Initial state	Final state	Feynman diagrams
	$H_1H_1, H_1H_2, H_2H_2, H_3H_3$	$t(\chi_k^0), u(\chi_k^0), s(H_{1,2})$
	$H_1H_3, H_2H_3$	$t(\chi_k^0), u(\chi_k^0), s(H_3), s(Z^0)$
	$H^-H^+$	$t(\chi_e^+), u(\chi_e^+), s(H_{1,2}), s(Z^0)$
	$Z^0 H_1,  Z^0 H_2$	$t(\chi_k^0), u(\chi_k^0), s(H_3), s(Z^0)$
$\chi_i^0 \chi_j^0$	$Z^{0}H_{3}$	$t(\chi_k^0), u(\chi_k^0), s(H_{1,2})$
0	$W^-H^+, W^+H^-$	$t(\chi_e^+), u(\chi_e^+), s(H_{1,2,3})$
	$Z^{0}Z^{0}$	$t(\chi_k^0),  u(\chi_k^0),  s(H_{1,2})$
	$W^-W^+$	$t(\chi_e^+), u(\chi_e^+), s(H_{1,2}), s(Z^0)$
_	$far{f}$	$t(\tilde{f}_{L,R}), u(\tilde{f}_{L,R}), s(H_{1,2,3}), s(Z^0)$

CHAPTER 30. THE MINIMAL SUPERSYMMETRIC STANDARD MODEL

Neutralino-chargino annihilation

Initial state	Final state	Feynman diagrams
	$H^+H_1, H^+H_2$	$t(\chi_k^0),  u(\chi_e^+),  s(H^+),  s(W^+)$
	$H^+H_3$	$t(\chi_k^0),  u(\chi_e^+),  s(W^+)$
	$W^+H_1, W^+H_2$	$t(\chi_k^0),  u(\chi_e^+),  s(H^+),  s(W^+)$
	$W^+H_3$	$t(\chi_k^0),  u(\chi_e^+),  s(H^+)$
$\chi_c^+ \chi_i^0$	$H^+Z^0$	$t(\chi_k^0),  u(\chi_e^+),  s(H^+)$
	$\gamma H^+$	$t(\chi_c^+), s(H^+)$
	$W^+Z^0$	$t(\chi_k^0),  u(\chi_e^+),  s(W^+)$
	$\gamma W^+$	$t(\chi_c^+),  s(W^+)$
	$u ar{d}$	$t(\tilde{d}_{L,R}), u(\tilde{u}_{L,R}), s(H^+), s(W^+)$
	$ u \overline{\ell}$	$t(\tilde{\ell}_{L,R}), u(\tilde{\nu}_L), s(H^+), s(W^+)$

Chargino-chargino annihilation

Initial state	Final state	Feynman diagrams
	$H_1H_1, H_1H_2, H_2H_2, H_3H_3$	$t(\chi_e^+), u(\chi_e^+), s(H_{1,2})$
	$H_1H_3, H_2H_3$	$t(\chi_e^+), u(\chi_e^+), s(H_3), s(Z^0)$
	$H^+H^-$	$t(\chi_k^0),  s(H_{1,2}),  s(Z^0, \gamma)$
	$Z^0H_1,  Z^0H_2$	$t(\chi_e^+),  u(\chi_e^+),  s(H_3),  s(Z^0)$
	$Z^0H_3$	$t(\chi_e^+),  u(\chi_e^+),  s(H_{1,2})$
	$H^+W^-, W^+H^-$	$t(\chi_k^0),  s(H_{1,2,3})$
$\chi_c^+ \chi_d^-$	$Z^{0}Z^{0}$	$t(\chi_e^+),  u(\chi_e^+),  s(H_{1,2})$
	$W^+W^-$	$t(\chi_k^0),  s(H_{1,2}),  s(Z^0, \gamma)$
	$\gamma\gamma$ (only for $c = d$ )	$t(\chi_c^+),  u(\chi_c^+)$
	$Z^0\gamma$	$t(\chi_d^+),  u(\chi_c^+)$
	$uar{u}$	$t(\tilde{d}_{L,R}), s(H_{1,2,3}), s(Z^0, \gamma)$
	$ u \overline{ u}$	$t(\tilde{\ell}_{L,R}),  s(Z^0)$
	$ar{d}d$	$t(\tilde{u}_{L,R}), s(H_{1,2,3}), s(Z^0, \gamma)$
	$\bar{\ell}\ell$	$t(\tilde{\nu}_L), s(H_{1,2,3}), s(Z^0, \gamma)$
	$H^+H^+$	$t(\chi_{k}^{0}), u(\chi_{k}^{0})$
$\chi_c^+ \chi_d^+$	$H^+W^+$	$t(\chi_k^0),  u(\chi_k^0)$
- 30	$W^+W^+$	$t(\chi_k^0),  u(\chi_k^0)$

#### 30.2.1.4 Squark-squark annihilation

Note: The tables below are not entirely up to date, more processes are included than shown in the tables.

We will here denote squarks as  $\tilde{q}_a^i$  and  $\tilde{q}_b^j$  where *i* and *j* are the family indices and *a* and *b* are the mass eigenstate indices (running from 1 to 2). *k* and *l* will also be used as family indices for

processes including more squarks. Colour indices are suppressed.  $\tilde{u}^i$  is used as a generic notation for any up-type squark where *i* denotes the family index. Down-type squarks are denoted analogously.

Note that we will not (except in rare occations) show processes for  $\tilde{\nu}$  and  $\tilde{\ell}$  separately since they can easily be obtained from the squark processes by replacing  $\tilde{u}$  with  $\tilde{\nu}$  and  $\tilde{d}$  with  $\tilde{\ell}$  (and noting that we only have one mass eigenstate for the  $\tilde{\nu}$ . Also note that the  $\tilde{\nu} - \tilde{\ell}$ -sector is assumed not to be flavour-changing.

Initial state	Final state	Diagrams	Note
$\tilde{d}^i_a \tilde{d}^{i*}_b$	$\gamma\gamma, Z\gamma$	$t(\tilde{d}_{1,2}^i),  u(\tilde{d}_{1,2}^i),  p$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	ZZ	$t(\tilde{d}_{1,2}^i),  u(\tilde{d}_{1,2}^i),  p,  s(H_1, H_2)$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$W^-W^+$	$p, s(H_1, H_2, Z, \gamma), t(\tilde{u}_{1,2}^k)$	k = 1, 2, 3
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$ZH_2, ZH_1$	$t(\tilde{d}_{1,2}^i),  u(\tilde{d}_{1,2}^i),  s(Z,H_3)$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$ZH_3$	$t(\tilde{d}_{1,2}^i),  u(\tilde{d}_{1,2}^i),  s(H_1, H_2)$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$\gamma H_2,  \gamma H_1,  \gamma H_3$	$t(\tilde{d}^{i}_{1,2}),  u(\tilde{d}^{i}_{1,2})$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$H_2H_2, H_1H_1, H_1H_2$	$t(\tilde{d}_{1,2}^i), u(\tilde{d}_{1,2}^i), p, s(H_1, H_2)$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$H_2H_3, H_1H_3$	$s(Z, H_3), t(\tilde{d}_{1,2}^i), u(\tilde{d}_{1,2}^i)$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$H_3H_3$	$s(H_1, H_2), p, t(\tilde{d}_{1,2}^i), u(\tilde{d}_{1,2}^i)$	
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$W^-H^+$	$s(H_1, H_2, H_3), t(\tilde{u}_{1,2}^k)$	k = 1, 2, 3
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$H^-H^+$	$s(H_1, H_2, Z, \gamma), p, t(\tilde{u}_{1,2}^k)$	k = 1, 2, 3
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$f\bar{f} \ (f \neq d^i)$	$s(H_{1}^{\star}, H_{2}^{\star}, H_{3}^{\star}, Z, \gamma^{\star}, g^{\ddagger}), t(\chi_{c}^{+})^{\dagger}$	†) $f = u^k \ (k = 1, 2, 3), \star$
			Not for $f = \nu$ , ‡) Only for
			squarks/quarks
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$d^i ar d^i$	$s(H_1, H_2, H_3, Z, \gamma, g^{\dagger}), t(\tilde{\chi}_k^0, \tilde{g}^{\dagger})$	†) Only for squarks
$ ilde{d}^i_a  ilde{d}^{i*}_b$	Zg	$t( ilde{d}^{i}_{1,2}), u( ilde{d}^{i}_{1,2}), p$	Only for squarks
$ ilde{d}^i_a  ilde{d}^{i*}_b$	gg	$t( ilde{d}^i_{1,2}), u( ilde{d}^i_{1,2}), s(g), p$	Only for squarks
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$g\gamma$	$t( ilde{d}^{i}_{1,2}), u( ilde{d}^{i}_{1,2}), p$	Only for squarks
$ ilde{d}^i_a  ilde{d}^{i*}_b$	$gH_1, gH_2, gH_3$	$t( ilde{d}^{i}_{1,2}), u( ilde{d}^{i}_{1,2})$	Only for squarks

#### $\tilde{d}^i_a \tilde{d}^{i*}_b$ annihilation

# $\tilde{d}_a^i \tilde{d}_b^{j*}$ annihilation $(i \neq j)$

Initial state	Final state	Diagrams	Note
$ ilde{d}^i_a  ilde{d}^{j*}_b$	$W^+W^-$	$t(\tilde{u}_{1,2}^k)^\dagger$	Not included at present
$ ilde{d}^i_a  ilde{d}^{ ilde{j}*}_b$	$W^+H^-$	$t( ilde{u}_{1,2}^{k'})^{\dagger}$	Not included at present
$ ilde{d}^i_a  ilde{d}^{j*}_b$	$H^+H^-$	$t(\tilde{u}_{1,2}^{k'})^{\dagger}$	Not included at present
$ ilde{d}^i_a  ilde{d}^{*j}_b$	$d^i \bar{d}^j$	$t( ilde{\chi}_k^0, ilde{g}^\dagger)$	†) Only for squarks
$ ilde{d}^i_a  ilde{d}^{*j}_b$	$u^k \bar{u}^l$	$t(\tilde{\chi}_c^+)$	Only $k = i, l = j$ at present

### $\tilde{d}^i_a \tilde{d}^i_b$ annihilation

Initial state	Final state	Diagrams	Note
$ ilde{d}^i_a  ilde{d}^i_b$	$d^i d^i$	$t(\tilde{\chi}^0_k, \tilde{g}^\dagger),  u(\tilde{\chi}^0_k, \tilde{g}^\dagger)$	†) Only for squarks

### $\tilde{d}_a^i \tilde{d}_b^j$ annihilation $(i \neq j)$

Initial state	Final state	Diagrams	Note
$ ilde{d}^i_a  ilde{d}^j_b$	$d^i d^j$	$t(\tilde{\chi}_k^0, \tilde{g}^\dagger)$	†) Only for squarks

Initial state	Final state	Diagrams	Note
$\tilde{u}_a^i \tilde{u}_b^{i*}$	$\gamma\gamma^{\dagger},Z\gamma^{\dagger}$	$t(\tilde{u}_{1,2}^i),  u(\tilde{u}_{1,2}^i),  p$	†) Not for $\tilde{\nu}$
$\tilde{u}^i_a \tilde{u}^{i*}_b$	ZZ	$t(\tilde{u}_{1,2}^i),  u(\tilde{u}_{1,2}^i),  p,  s(H_1, H_2)$	
$\tilde{u}^i_a \tilde{u}^{i*}_b$	$W^-W^+$	$p,  s(H_1, H_2, Z, \gamma^{\dagger}),  u( ilde{d}^k_{1,2})$	$k = 1, 2, 3, \dagger$ Not for $\tilde{\nu}$
$\tilde{u}^i_a \tilde{u}^{i*}_b$	$ZH_2, ZH_1$	$t( ilde{u}^i_{1,2}), u( ilde{u}^i_{1,2}), s(Z,H_3^\dagger)$	†) Not for $\tilde{\nu}$
$\tilde{u}_a^i \tilde{u}_b^{i*}$	$ZH_3$	$t(\tilde{u}_{1,2}^i)^\dagger,  u(\tilde{u}_{1,2}^i)^\dagger,  s(H_1, H_2)$	†) Not for $\tilde{\nu}$
$\tilde{u}^i_a \tilde{u}^{i*}_b$	$\gamma H_2^\dagger,\gamma H_1^\dagger,\gamma H_3^\dagger$	$t( ilde{u}^i_{1,2}), u( ilde{u}^i_{1,2})$	†) Not for $\tilde{\nu}$
$\tilde{u}_a^i \tilde{u}_b^{i*}$	$H_2H_2, H_1H_1, H_1H_2$	$t(\tilde{u}_{1,2}^i), u(\tilde{u}_{1,2}^i), p, s(H_1, H_2)$	
$\tilde{u}_a^i \tilde{u}_b^{i*}$	$H_2H_3, H_1H_3$	$s(Z, H_3^{\dagger}), t(\tilde{u}_{1,2}^i)^{\dagger}, u(\tilde{u}_{1,2}^i)^{\dagger}$	†) Not for $\tilde{\nu}$
$\tilde{u}^i_a \tilde{u}^{i*}_b$	$H_3H_3$	$s(H_1, H_2),  p,  t(\tilde{u}_{1,2}^i)^\dagger,  u(\tilde{u}_{1,2}^i)^\dagger$	†) Not for $\tilde{\nu}$
$\tilde{u}_a^i \tilde{u}_b^{i*}$	$W^-H^+$	$s(H_1, H_2, H_3^{\dagger}),  u(\tilde{d}_{1,2}^k)$	$k = 1, 2, 3, \dagger$ Not for $\tilde{\nu}$
$\tilde{u}_a^i \tilde{u}_b^{i*}$	$H^+H^-$	$s(H_1, H_2, Z, \gamma^{\dagger}),  p,  t(\tilde{d}_{1,2}^k)$	$k = 1, 2, 3, \dagger$ Not for $\tilde{\nu}$
$\tilde{u}^i_a \tilde{u}^{i*}_b$	$f\bar{f} \ (f \neq u^i)$	$s(H_1^{\times}, H_2^{\times}, H_3^{\dagger \times}, Z, \gamma^{\dagger \times}, g^{\ddagger}), t(\chi_c^+)^{\star}$	†) Not for $\tilde{\nu}, \star$ ) If $f = d^k$
			$(k = 1, 2, 3), \ddagger)$ Only for
			squarks/quarks, $\times$ ) Not for
ãi ãi*	aiāi	$a(H^{\times} H^{\times} H^{\times} T^{\times} 2 \alpha^{\times} a^{\ddagger}) t(\tilde{x}^{0} \tilde{a}^{\ddagger})$	$\nu$ ) Not for $\mu$ (t) Only for
$u_a u_b$	u u	$S(\Pi_1, \Pi_2, \Pi_3, Z, \gamma, g^*), \iota(\chi_k, g^*)$	squarks
$\tilde{u}_{a}^{i}\tilde{u}_{b}^{i*}$	Za	$t(\tilde{u}_{1,2}^{i}), u(\tilde{u}_{1,2}^{i}), p$	Only for squarks
$\tilde{u}_{a}^{i}\tilde{u}_{b}^{i*}$	qq	$t(\tilde{u}_{1,2}^{i}), u(\tilde{u}_{1,2}^{i}), s(q), p$	Only for squarks
$\tilde{u}_{a}^{i}\tilde{u}_{b}^{i*}$	$g\gamma$	$t(\tilde{u}_{1,2}^{i}), u(\tilde{u}_{1,2}^{i}), p$	Only for squarks
$ ilde{u}_{a}^{ar{i}} ilde{u}_{b}^{ar{i}*}$	$gH_1, gH_2, gH_3$	$t(\tilde{u}_{1,2}^{i}), u(\tilde{u}_{1,2}^{i})$	Only for squarks

# $\tilde{u}^i_a \tilde{u}^{i*}_b$ annihilation

# $\tilde{u}_a^i \tilde{u}_b^{j*}$ annihilation $(i \neq j)$

Initial state	Final state	Diagrams	Note
$\tilde{u}^i_a \tilde{u}^{j*}_b$	$W^+W^-$	$t(\tilde{d}_{1,2}^k)^\dagger$	Not included at present, †) Not for $\tilde{\ell}$
$\tilde{u}_a^i \tilde{u}_b^{j*}$	$W^+H^-$	$t( ilde{d}^{k}_{1,2})^{\dagger}$	Not included at present, †) Not for $\tilde{\ell}$
$ ilde{u}^i_a  ilde{u}^{j*}_b$	$H^+H^-$	$t(\tilde{d}_{1,2}^k)^\dagger$	Not included at present, †) Not for $\tilde{\ell}$
$ ilde{u}^i_a  ilde{u}^{j*}_b$	$u^i \bar{u}^j$	$t( ilde{\chi}_k^{0},g^\dagger)$	†) Only for squarks
$\tilde{u}_a^i \tilde{u}_b^{j*}$	$d^k \overline{d^l}$	$t(\tilde{\chi}_c^+)$	Only $k = i, l = j$ at present

# $\tilde{u}^i_a \tilde{u}^i_b$ annihilation

Initial state	Final state	Diagrams	Note
$ ilde{u}^i_a  ilde{u}^i_b$	$u^i u^i$	$t( ilde{\chi}^0_k, ilde{g}^\dagger),u( ilde{\chi}^0_k, ilde{g}^\dagger)$	†) Only for squarks

# $\tilde{u}_a^i \tilde{u}_b^j$ annihilation $(i \neq j)$

Initial state	Final state	Diagrams	Note
$ ilde{u}^i_a  ilde{u}^j_b$	$u^i u^j$	$t( ilde{\chi}_k^0,  ilde{g}^\dagger)$	†) Only for squarks

# $\tilde{u}^i_a \tilde{d}^{i*}_b$ annihilation

Initial state	Final state	Diagrams	Note
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$H^+H_1, H^+H_2$	$t(\tilde{d}_{1,2}^i),  u(\tilde{u}_{1,2}^i),  p,  s(W^+, H^+)$	
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$H^+H_3$	$t(\tilde{d}_{1,2}^{i}),  u(\tilde{u}_{1,2}^{i})^{\dagger},  p,  s(W^{+})$	†) Not for $\tilde{\ell}$
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$\gamma H^+$	$t(\tilde{u}_{1,2}^{i})^{\dagger},  u(\tilde{d}_{1,2}^{i}),  s(H^{+})$	†) Not for $\tilde{\ell}$
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$ZH^+$	$t(\tilde{u}_{1,2}^{i}),  u(\tilde{d}_{1,2}^{i}),  s(H^{+})$	
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$W^+H_1, W^+H_2$	$t(\tilde{d}_{1,2}^i),  u(\tilde{u}_{1,2}^i),  s(W^+, H^+)$	
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$W^+H_3$	$t(\tilde{d}_{1,2}^i),  u(\tilde{u}_{1,2}^i)^\dagger,  s(H^+)$	†) Not for $\tilde{\ell}$
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$W^+\gamma$	$t(\tilde{d}_{1,2}^i),  u(\tilde{u}_{1,2}^i)^\dagger,  s(W^+),  p$	†) Not for $\tilde{\ell}$
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$W^+Z$	$t(\tilde{d}_{1,2}^i),  u(\tilde{u}_{1,2}^i),  s(W^+),  p$	
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$u^k \overline{d}^l$	$s(H^+,W^+)^{\star}, t(\tilde{\chi}^0_m,\tilde{g}^{\dagger})\delta^{ik}\delta^{il}$	†) Not for $\tilde{\ell}, \star$ ) Only $k = l$
~		~.	at present
$ ilde{u}^i_a d^{i*}_b$	$W^+g$	$t(d_{1,2}^i),  u(\tilde{u}_{1,2}^i),  p$	Only for squarks
$ ilde{u}^i_a  ilde{d}^{i*}_b$	$gH^+$	$t( ilde{u}_{1,2}^{i}),  u( ilde{d}_{1,2}^{i})$	Only for squarks

CHAPTER 30. THE MINIMAL SUPERSYMMETRIC STANDARD MODEL

 $\tilde{u}_a^i \tilde{d}_b^{j*}$  annihilation  $(i \neq j)$  For squarks we can have the following processes

Initial state	Final state	Diagrams	Note
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$H^+H_1, H^+H_2$	$t(\tilde{d}_{1,2}^j),  u(\tilde{u}_{1,2}^i),  p,  s(W^+, H^+)$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$H^+H_3$	$t(\tilde{d}_{1,2}^{j}),  u(\tilde{u}_{1,2}^{i}),  p,  s(W^+)$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$H^+\gamma$	$t(\tilde{d}_{1,2}^{j}),  u(\tilde{u}_{1,2}^{i}),  s(H^+)$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$H^+Z$	$t(\tilde{d}_{1,2}^{j}),  u(\tilde{u}_{1,2}^{i}),  s(H^+)$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$W^+H_1, W^+H_2$	$t(\tilde{d}_{1,2}^{j}),  u(\tilde{u}_{1,2}^{i}),  s(W^+, H^+)$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$W^+H_3$	$t(\tilde{d}_{1,2}^{j}),  u(\tilde{u}_{1,2}^{i}),  s(H^+)$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$W^+\gamma$	$t(\tilde{d}_{1,2}^{j}),  u(\tilde{u}_{1,2}^{i}),  s(W^{+}),  p$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$W^+Z$	$t(\tilde{d}_{1,2}^{j}),  u(\tilde{u}_{1,2}^{i}),  s(W^{+}),  p$	Not included at present
$ ilde{u}^i_a  ilde{d}^{j*}_b$	$u^k \bar{d}^l$	$s(H^+,W^+)^{\dagger},t(\tilde{\chi}^0_m,\tilde{g})\delta^{ik}\delta^{jl}$	†) Not included at present

whereas for sneutrinos and sleptons, we can only have the process

Initial state	Final state	Diagrams	Note
$\tilde{ u}^i \tilde{\ell}_b^{j*}$	$ u^i ar{\ell}^j$	$t( ilde{\chi}_k^0)$	

# $\tilde{u}^i_a \tilde{d}^i_b$ annihilation

Initial state	Final state	Diagrams	Note
$ ilde{u}^i_a  ilde{d}^i_b$	$u^k d^l$	$t(\tilde{\chi}_m^0, \tilde{g}^{\dagger}) \delta^{ik} \delta^{il},  u(\tilde{\chi}_c^+)^{\star}$	†) Only for squarks, $\star$ )
			Only $i = k = l$ at present

# $\tilde{u}_a^i \tilde{d}_b^j$ annihilation $(i \neq j)$

Initial state	Final state	Diagrams	Note
$ ilde{u}^i_a  ilde{d}^j_b$	$u^k d^l$	$t(\tilde{\chi}^0_m, \tilde{g}^{\dagger}) \delta^{ik} \delta^{jl},  u(\tilde{\chi}^+_c)^{\times \star}$	†) Only for squarks, $\times$ )
			For $\tilde{\nu}\tilde{\ell}$ only when $i =$
			$l, j = k, \star$ ) Only included
			when $i = l, j = k$ at
			present

#### 30.2.1.5 Squark-neutralino annihilation

Note: The tables below are not entirely up to date, more processes are included than shown in the tables.

We will here denote squarks as  $\tilde{u}_a^i$  and  $\tilde{d}_a^i$  where *i* is the family index and *a* is the mass eigenstate index (running from 1 to 2).

Initial state	Final state	Diagrams	Note
$ ilde{u}^i_a  ilde{\chi}^0_i$	$\gamma u^i$	$s(u^i), t(\tilde{u}^i_{1,2})$	Only for squarks
$ ilde{u}^i_a  ilde{\chi}^0_i$	$Zu^i$	$s(u^{i}), t(\tilde{u}_{1,2}^{i}), u(\tilde{\chi}_{k}^{0})$	
$ ilde{u}^i_a  ilde{\chi}^0_i$	$H_1 u^i, H_2 u^i$	$s(u^{i})^{\dagger}, t(\tilde{u}_{1,2}^{i}), u(\tilde{\chi}_{k}^{0})$	†) Only for squarks
$ ilde{u}^i_a ilde{\chi}^0_j$	$H_3 u^i$	$s(u^{i})^{\dagger}, t(\tilde{u}_{1,2}^{i})^{\dagger}, u(\tilde{\chi}_{k}^{0})$	†) Only for squarks
$ ilde{u}^i_a  ilde{\chi}^{\dot{0}}_j$	$W^+ d^k$	$s(u^{i}), t(\tilde{d}^{k}_{1,2}), u(\tilde{\chi}^{+}_{c})$	$k = 1, 2, 3,  \tilde{u}_a^{i*} \tilde{\chi}_i^0 \rightarrow$
$ ilde{u}^i_a ilde{\chi}^0_j$	$H^+ d^k$	$s(u^i), t(\tilde{d}^k_{1,2}), u(\tilde{\chi}^+_c)$	$ \begin{array}{l} W^- \vec{d}^k \text{ in the code} \\ k = 1, 2, 3,  \tilde{u}^{i*}_a \tilde{\chi}^0_j \rightarrow H^- \vec{d}^k \end{array} $
$\tilde{u}_a^i \tilde{\chi}_j^0$	$gu^i$	$s(u^i),t(\tilde{u}^i_{1,2})$	in the code

#### $\tilde{u}_a^i \tilde{\chi}_j^0$ annihilation

### $\tilde{d}^i_a \tilde{\chi}^0_j$ annihilation

Initial state	Final state	Diagrams	Note
$\tilde{d}^i_a \tilde{\chi}^0_j$	$\gamma d^i$	$s(d^i), t(\tilde{d}^i_{1,2})$	
$ ilde{d}^i_a  ilde{\chi}^{\dot{0}}_j$	$Zd^i$	$s(d^{i}), t(\tilde{d}^{i}_{1,2}), u(\tilde{\chi}^{0}_{k})$	
$ ilde{d}^i_a  ilde{\chi}^{0}_j$	$H_1 d^i, H_2 d^i$	$s(d^{i}), t(\tilde{d}^{i}_{1,2}), u(\tilde{\chi}^{0}_{k})$	
$ ilde{d}^i_a  ilde{\chi}^{0}_j$	$H_3 d^i$	$s(d^{i}), t(\tilde{d}^{i}_{1,2}), u(\tilde{\chi}^{0}_{k})$	
$ ilde{d}^i_a  ilde{\chi}^0_j$	$W^- u^k$	$s(d^{i}), t(\tilde{u}_{1,2}^{k}), u(\tilde{\chi}_{c}^{+})$	k = 1, 2, 3
$ ilde{d}^i_a  ilde{\chi}^{\dot{0}}_j$	$H^- u^k$	$s(d^{i}), t(\tilde{u}_{1,2}^{k}), u(\tilde{\chi}_{c}^{+})$	k=1,2,3
$ ilde{d}^i_a  ilde{\chi}^0_j$	$gd^i$	$s(d^i), t(\tilde{d}^i_{1,2})$	

#### 30.2.1.6 Squark-chargino annihilation

# Note: The tables below are not entirely up to date, more processes are included than shown in the tables.

We will here denote squarks as  $\tilde{q}_a^i$  where *i* is the family index and *a* is the mass eigenstate index (running from 1 to 2).

#### $\tilde{u}^i_a \tilde{\chi}^+_c$ annihilation

Initial state	Final state	Diagrams	Note
$\tilde{u}_a^i \tilde{\chi}_c^+$	$W^+ u^k$	$t(\tilde{d}_{1,2}^l),  u(\tilde{\chi}_c^0)\delta^{ik}$	Only $k = l = i$ at present
$\tilde{u}_a^i \tilde{\chi}_c^+$	$H^+ u^k$	$t(\tilde{d}^l_{1,2}),  u(\tilde{\chi}^0_c)\delta^{ik}$	Only $k = l = i$ at present

 $\tilde{u}_a^{i*}\tilde{\chi}_c^+$  annihilation

Initial state	Final state	Diagrams	Note
$\tilde{u}_a^{i*}\tilde{\chi}_c^+$	$Z \bar{d}^k$	$s(\bar{d}^k), t(\tilde{u}_{1,2}^i), u(\tilde{\chi}_c^+)$	k = 1, 2, 3
$\tilde{u}_a^{i*}\tilde{\chi}_c^+$	$\gamma d^k$	$s(\bar{d}^k), t(\tilde{u}_{1,2}^i)^{\dagger}, u(\tilde{\chi}_c^+)$	$k = 1, 2, 3, \dagger$ ) Only for squarks
$ ilde{u}_a^{i*} ilde{\chi}_c^+$	$H_1 \bar{d}^k,  H_2 \bar{d}^k$	$s(\overline{d^k}), t(\tilde{u}_{1,2}^i), u(\tilde{\chi}_c^+)$	k = 1, 2, 3
$ ilde{u}_a^{i*} ilde{\chi}_c^+$	$H_3 \bar{d}^k$	$s(\bar{d}^k), t(\tilde{u}_{1,2}^i)^{\dagger}, u(\tilde{\chi}_c^+)$	$k = 1, 2, 3, \dagger$ ) Only for squarks
$\tilde{u}_a^{i*}\tilde{\chi}_c^+$	$W^+ \bar{u}^k$	$s(ar{d^l}), u( ilde{\chi}^0_c)\delta^{ik}$	Only $k = l = i$ at present
$\tilde{u}_a^{i*}\tilde{\chi}_c^+$	$H^+ \bar{u}^k$	$s(ar{d^l}), u( ilde{\chi}^0_c)\delta^{ik}$	Only $k = l = i$ at present
$ ilde{u}_a^{i*} ilde{\chi}_c^+$	$g \overline{d}^k$	$s(\bar{d}^k),t(\tilde{u}^i_a)$	k = 1, 2, 3, only for squarks

#### $\tilde{d}_a^i \tilde{\chi}_c^+$ annihilation

Initial state	Final state	Diagrams	Note
$\tilde{d}^i_a \tilde{\chi}^+_c$	$Zu^k$	$s(u^k), t(\tilde{d}^i_{1,2}), u(\tilde{\chi}^+_c)$	Only $k = i$ at present
$ ilde{d}^i_a  ilde{\chi}^+_c$	$\gamma u^k$	$s(u^k)^{\dagger}, t(\tilde{d}^i_{1,2}), u(\tilde{\chi}^+_c)$	Only $k = i$ at present, $\dagger$ ) Only for squarks
$ ilde{d}^i_a  ilde{\chi}^+_c$	$H_1 u^k, H_2 u^k$	$s(u^k)^{\dagger}, t(\tilde{d}^i_{1,2}), u(\tilde{\chi}^+_c)$	Only $k = i$ at present, $\dagger$ ) Only for squarks
$\tilde{d}^i_a \tilde{\chi}^+_c$	$H_3 u^k$	$s(u^k)^{\dagger}, t(\tilde{d}^i_{1,2}), u(\tilde{\chi}^+_c)$	Only $k = i$ at present, $\dagger$ ) Only for squarks
$ ilde{d}^i_a  ilde{\chi}^+_c$	$W^+ d^k$	$s(u^l),  u(\tilde{\chi}^0_c) \delta^{ik}$	Only $k = l = i$ at present
$\tilde{d}^i_a \tilde{\chi}^+_c$	$H^+ d^k$	$s(u^l),  u(\tilde{\chi}^0_c) \delta^{ik}$	Only $k = l = i$ at present
$\tilde{d}^i_a \tilde{\chi}^+_c$	$gu^k$	$s(u^k),  t(\tilde{d}^i_a)$	Only $k = i$ at present, only for squarks

#### $\tilde{d}_a^{i*}\tilde{\chi}_c^+$ annihilation

Initial state	Final state	Diagrams	Note
$\tilde{d}_a^{i*}\tilde{\chi}_c^+$	$W^+ \overline{d}^k$	$t(\tilde{u}_{1,2}^l),  u(\tilde{\chi}_c^0)\delta^{ik}$	Only $k = l = i$ at present
$\tilde{d}_a^{i*}\tilde{\chi}_c^+$	$H^+ \overline{d}^k$	$t(\tilde{u}_{1,2}^{l'}),  u(\tilde{\chi}_c^0)\delta^{ik}$	Only $k = l = i$ at present

#### 30.2.1.7 Degrees of freedom

We have to be careful with the internal degrees of freedom, g, of the particles. We can either treat e.g. a  $\chi_i^+$  and a  $\chi_i^-$  as two separate particles with two degrees of freedom each, or we can treat them as one particle  $\chi_i^{\pm}$  with four degrees of freedom. The latter approach has an advantage that we simplify our expressions for the effective annihilation cross sections when coannihilations are needed. Hence, we use that approach here. For a more detailed discussion about this, see Section 30.12.2.

#### **30.2.2** Annihilation routines - general remarks

The annihilation cross section routines is divided into several parts, mostly for historical reasons. The layout is roughly as follows:

- src/an Here we keep the main routins for both neutralino- neutralino annihilation cross sections and the effective annihilation cross section in the relic density calculations. The steering routines for neutralino and chargino coannihilations are also kept here.
- src/anstu Here keep the t-, u- and s- diagram expressions for fermion-fermion coannihilations (i.e. neutralino and chargino coannihilations).

src/as Here all the coannihilation cross sections including sfermions are kept.

We will here describe the src/an-routines.

#### **30.2.2.1** General routines

The general routine to call for an effective annihilation cross section (to be used for relic density calculations) is **dsanwx**, which returns the invariant annihilation rate (integrated over  $\cos \theta$ ). The actual cross section, differential in  $\cos \theta$  is calculated by **dsandwdcos** which includes all the coannihilations needed. This is set up in **mssm/rd/dsrdparticles** (typically called from **dsrdomega** in the **core** library) which determines which coannihilating particles to include.

For other applications where the annihilation rate is needed, e.g. annihilation in the galactic halo, one can call the specific annihilation rate routine directly. The main one is **dsandwdcosnn** for neutralino-neutralino annihilation. To simplify this task, we supply a routine **dssigmav0** which calls **dsandwdcosnn** for neutralino-neutralino annihilation at zero relative velocity and returns the result, either as the total annihilation cross section, or the cross section for a specific channel. See the header of **dssigmav** for details.

#### 30.2.2.2 Neutralino and chargino (co)annihilation cross sections

The routines **dsandwdcosnn**, **dsandwdcoscn** and **dsandwdcoscc** calculate the annihilation cross sections (returning the invariant annihilation rate) for neutralino-neutralino, neutralino-chargino and chargino-chargino annihilations. Which particles the cross section is calculated for is given by particle indices as defined in **mssm/include/dsmssm.h**.

All the annihilation routines return the invariant rate instead of the cross section. The invariant annihilation rate between particle i and j is defined as

$$W_{ij} = 4p_{ij}\sqrt{s\sigma_{ij}} = 4\sigma_{ij}\sqrt{(p_i \cdot p_j)^2 - m_i^2 m_j^2} = 4E_i E_j \sigma_{ij} v_{ij}.$$
(30.1)

See Chapter 20 for more details.

# 30.3 mssm/an\_11: Annihilation cross sections (1-loop)

#### 30.3.1 Annihilation cross sections at 1-loop – general

The annihilation cross sections at 1-loop that we have implemented in DarkSUSY are those to  $\gamma\gamma$ ,  $Z\gamma$  and gg. The derivation of these is described in the works [22, 23]. Let us mention that these one-loop expressions formally violate unitarity for diagrams with electroweak gauge-boson exchange because they do not take into account the nonperturbative effects related to multiple electroweak gauge boson exchange ('Sommerfeld effect') as described in [?, ?]. In practice, one can still trust the result for sub-TeV neutralinos – but should keep in mind that the implemented cross sections become unphysical in the limit  $m_W/m_{\chi} \rightarrow 0$ .

To see how these routines are called, see the file src\_models/mssm/an/dsandwdcosnn.f where the  $\gamma\gamma$ ,  $Z\gamma$  and gg contributions are added to the annihilation cross section at the end.

# 30.4 mssm/an\_ib: Internal bremsstrahlung

#### 30.4.1 Internal Bremsstrahlung (IB) – theory

Whenever WIMPs annihilate into pairs of charged particles  $X\bar{X}$ , this process will inevitably be accompanied by internal bremsstrahlung (IB), i.e. the emission of an additional photon in the final state (note that in contrast to ordinary, or external, bremsstrahlung no external electromagnetic field is required for the emission of the photon). In many cases, IB photons completely dominate the annihilation spectrum at the highest accessible energies. The resulting, characteristic sharp step at an energy corresponding to the dark matter particle's mass, often accompanied by a bump-like feature at slightly smaller energies, is a spectral signature that is hard to mimic by astrophysical sources and in that respect similar to monochromatic photons. In fact, being an  $\mathcal{O}(\alpha_{\rm em})$  correction to the tree-level annihilation rate, one would generically expect IB photons to be more copiously produced than the loop-suppressed [i.e.  $\mathcal{O}(\alpha_{\rm em}^2)$ ] monochromatic photons; this has been confirmed in [24] for a large part of the supersymmetric parameter space, not the least due to the appearance of efficient enhancement mechanisms.

#### **30.4.1.1** General considerations

For didactic purposes, one may follow [24] and distinguish between photons directly radiated off the external legs (*final state radiation*, FSR) and photons radiated from virtual charged particles (*virtual internal bremsstrahlung*, VIB). The IB photons are thus the total contribution from both FSR and VIB photons.

For relativistic charged final states, FSR diagrams are always dominated by photons emitted *collinearly* with X or  $\bar{X}$ . This is a purely kinematical effect and related to the fact that the propagator of the corresponding outgoing particle,

$$D(p) \propto \left( (k+p)^2 - m_X^2 \right)^{-1},$$
 (30.2)

diverges in this situation. Here, k and p denote the momenta of the photon and the outgoing particle, respectively. The resulting photon spectrum turns out to be of a universal form, almost independent of the underlying particle physics model (see, e.g., [159]):

$$\frac{dN_{\gamma,\text{FSR}}^{XX}}{dx} \approx \frac{\alpha Q_X^2}{\pi} \mathcal{F}_X(x) \log\left(\frac{s(1-x)}{m_X^2}\right) \,. \tag{30.3}$$

Here,  $Q_X$  and  $m_X$  are the electric charge and mass of X; the splitting function  $\mathcal{F}(x)$  depends only on the spin of the final state particles and takes the form

$$\mathcal{F}_{\text{fermion}}(x) = \frac{1 + (1 - x)^2}{x}$$
(30.4)

for fermions and

$$\mathcal{F}_{\text{boson}}(x) = \frac{1-x}{x} \tag{30.5}$$

for bosons. Due to the logarithmic enhancement that becomes apparent in Eq. (30.3), FSR photons are often the main source for IB. A prominent example where FSR in this universal form not only dominates IB but in fact the total gamma-ray spectrum from WIMP annihilations, is the case of Kaluza-Klein dark matter [160].

In general, one can single out two situations where photons emitted from virtual charged particles may give an even more important contribution to the total IB spectrum than FSR: i) the three-body final state  $X\bar{X}\gamma$  satisfies a symmetry of the initial state that cannot be satisfied by the two-body final state  $X\bar{X}$  or ii) X is a boson and the annihilation into  $X\bar{X}$  is dominated by t-channel diagrams, with the t-channel particle almost degenerate in mass with the annihilating WIMP. In contrast to FSR, the contribution from VIB photons can not be given in a model-independent way but is very sensitive to the underlying short-distance physics. For more details, see [24].

#### **30.4.1.2** IB from neutralino annihilations

For supersymmetric dark matter annhibitions, the relevant final states for IB are  $W^+W^-$ ,  $W^{\pm}H^{\mp}$ ,  $H^+H^-$  and  $f\bar{f}$ ; both of the situations just mentioned above can arise, and *VIB* contributions become important in considerable regions of the parameter space.

Let us first note that for neutralino annihilations, in contrast to the situation for, e.g., Kaluza-Klein dark matter, we cannot in general expect very large FSR contributions. This is because the lightest charged final states, for which the logarithmic enhancement shown in Eq. (30.3) would be most effective, are fermionic and therefore strongly helicity suppressed. Fermion final states containing an additional photon, however, are not subject to such a suppression [161]. In the limit of vanishing fermion mass, and assuming that both corresponding sfermions have the same mass, the photon multiplicity is given by [24]

$$\frac{\mathrm{d}N^{f^+f^-}}{\mathrm{d}x} = \alpha_{\rm em}Q_f^2 \frac{|\tilde{g}_R|^4 + |\tilde{g}_L|^4}{64\pi^2} \left(m_\chi^2 \langle \sigma v \rangle_{\chi\chi \to f\bar{f}}\right)^{-1}$$
(30.6)  
  $\times (1-x) \left\{ \frac{4x}{(1+\mu)(1+\mu-2x)} - \frac{2x}{(1+\mu-x)^2} - \frac{(1+\mu)(1+\mu-2x)}{(1+\mu-x)^3} \log \frac{1+\mu}{1+\mu-2x} \right\},$ 

where  $\mu \equiv m_{\tilde{f}_R}^2/m_{\chi}^2 = m_{\tilde{f}_L}^2/m_{\chi}^2$  and  $\tilde{g}_R P_L$  ( $\tilde{g}_L P_R$ ) denotes the coupling between neutralino, fermion and right-handed (left-handed) sfermion. In the above expression, a large factor  $m_{\chi}^2/m_f^2$  due to the lifted helicity suppression (from  $\langle \sigma v \rangle_{\chi\chi \to f\bar{f}} \propto m_f^2 m \chi^{-4}$ ) appears, and another enhancement at high photon energies for sfermions degenerate with the neutralino.

For large neutralino masses  $m_{\chi} \gg m_W$  and charginos almost degenerate with the neutralino,  $W^+W^-$  and  $W^{\pm}H^{\mp}$  final states are affected by the second of the mechanisms for VIB enhancement that were discussed in the previous subsection. Of these two channels, IB from  $W^+W^-$  nearly always dominates; for pure Higgsinos (or Winos), the resulting photon multiplicity in this limit is well approximated by [24, 162]:

$$\frac{\mathrm{d}N^{W^+W^-}}{\mathrm{d}x} \approx \frac{\alpha_{\mathrm{em}}}{\pi} \frac{4(1-x+x^2)^2}{(1-x+\epsilon/2)x} \left[ \log\left(2\frac{1-x+\epsilon/2}{\epsilon}\right) - 1/2 + x - x^3 \right], \tag{30.7}$$

where  $\epsilon \equiv m_W/m_{\chi}$ .

Charged Higgs pairs  $H^+H^-$ , finally, provide yet another interesting example where the two-body final state is not allowed due to symmetry restrictions; in this case, in the limit  $v \to 0$ , enforced by CP conservation. The annihilation into  $H^+H^-\gamma$ , on the other hand, is possible. However, since charged Higgs bosons in most models have considerably larger masses than gauge bosons, they are expected to give negligible IB contributions compared to the latter.

#### 30.4.1.3 The implementation in DarkSUSY

Let us now briefly describe how IB is implemented in DarkSUSY. The total gamma-ray spectrum from WIMP annihilations is given by

$$\frac{dN_{\gamma,\text{tot}}}{dx} = \sum_{f} B_f \left( \frac{dN_{\gamma,\text{sec}}^f}{dx} + \frac{dN_{\gamma,\text{IB}}^f}{dx} + \frac{dN_{\gamma,\text{line}}^f}{dx} \right), \qquad (30.8)$$

where  $B_f$  denotes the branching ratio into the annihilation channel f. The first term encodes the contribution from secondary photons, mainly produced through the decay of neutral pions, as described in Chapter 11. We recall here that these contributions are included by using the Monte Carlo code Pythia [10] to simulate the decay of a hypothetical particle with mass  $2m\chi$  and user-specified branching ratios  $B_f$ . In this way, also FSR associated to this decay is automatically included in  $dN_{\gamma,\text{sec}}^f/dx$  (the main contribution here comes from photons directly radiated off the external legs, but also photons radiated from other particles in the decay cascade are taken into account).

Of course, IB from the decay of such a hypothetical particle cannot in general be expected to show the same characteristics as IB from the actual annihilation of two WIMPs and for this reason an additional term  $dN_{\gamma,\text{IB}}^f/dx$  is included that accounts for the difference between the full

IB contribution and the FSR part already taken into account for by Pythia. For details regarding the implemented procedure of separating these two contributions in a consistent way, see [24]. The contributions  $dN_{\gamma,\text{IB}}^f/dx$ , in contrast to  $dN_{\gamma,\text{sec}}^f/dx$ , are generically highly model-dependent. At the moment, they are fully implemented only for neutralino annihilations and included by default. However, one may easily switch to a user-defined contribution, choose to neglect IB

default. However, one may easily switch to a user-defined contribution, choose to neglect IB completely or, for comparison, to only include the FSR part by replacing the call to dsibyield in mssm/an\_yield/dsanyield. For example, a version of dsibyield containing only the FSR part of the yield is provided in mssm/an\_ib/dsanyield\_fsr.

For the supersymmetric case, the full expressions for all relevant three-body final states are implemented, i.e. not just the approximations given in the last subsection, which only apply to the limits described there. For performance reasons, IB is only included when virtual *t*-channel particles are sufficiently degenerate in mass that large IB contributions to the total spectrum can be expected. With a call to **dslBset**, this default behaviour can be customized. Finally, radiative corrections to the annihilation into charged particles  $X\bar{X}$  of course also change the number of  $X\bar{X}$ pairs per annihilation and thus the corresponding yield of particles in the further decay of the annihilation products. At the moment, apart from the photon yield, only the IB positron yield is implemented in **DarkSUSY**. By default, only the annihilation into the dominating channel  $e^+e^-\gamma$  is taken into account in this case; again, this behaviour can be modified by a call to **dslBset**.

# 30.5 mssm/an\_sf: Annihilation cross sections (with sfermions)

#### 30.5.1 Annihilation cross sections with sfermions – general

In this directory, all the (co)annihilation cross sections involving one or more sfermions in the initial state are calculated. The code here is based upon the work described in [14]. All the cross sections are calculated with Form and converted to Fortran with a script **form2f** [158].

The main routines here are

Routine	Purpose
dsasdwdcossfsf	Calculates the invariant annihilation rate between two sfermions in the initial
	state.
dsasdwdcossfchi	Calculates the invariant annihilation rate between one sfermion and one fermion (neutralino or chargino) in the iniital state.

# 30.6 mssm/an\_stu: t, u and s diagrams for ff-annihilation

#### 30.6.1 Annihilation amplitudes for fermion-fermion annihilation

In this directory, all the helicity amplitudes needed for neutralino-neutralino, neutralino-chargino and chargino-chargino annihilation are calculated. The helicity amplitudes have been calculated with general expressions for vertices, masses etc. in **Reduce** and converted to Fortran files. The calculation of these are described in more detail in [13].

Each routine here adds the contribution to the helicity amplitudes from one particular diagram and the sum over contributed diagrams is done in the routines **an/dsandwdcosnn**, **an/dsandwdcoscn** and **an/dsandwdcoscc**. The naming convention for the routines here is the following: The first part of the routine name is **dsan** to indicate that they deal with annihilations in DarkSUSY. The next character tells which kind of process it is *s*-, *t*- or *u*-channel and the next two caracters tell which initial state particles we have (**f** for fermion), the next character is the kind of propagating particle (**f** for fermion, **s** for scalar and **v** for vector boson), and finally, the last two characters tell the kind of final state particles. So, to take an example, the routine **dsansffsvv** calculates the helicity amplitudes for annihilation of two fermions to two vector bosons via *s*-channel exchange of a scalar. There are also a few special cases (routines ending in  $\mathbf{ex}$  or  $\mathbf{in}$ ) for diagrams with clashing arrows.

### 30.7 mssm/dd: Direct detection

#### **30.7.1** Direct detection – theory

# NOTE: The description below is outdated. **DarkSUSY** now includes much better form factors.

If neutralinos are indeed the CDM needed on galaxy scales and larger, there should be a substantial flux of these particles in the Milky Way halo. Since the interaction strength is essentially given by the same weak couplings as, e.g., for neutrinos there is a non-negligible chance of detecting them in low-background counting experiments [163]. Due to the large parameter space of MSSM, even with the simplifying assumptions above, there is a rather wide span of predictions for the event rate in detectors of various types. It is interesting, however, that the models giving the largest rates are already starting to be probed by present direct detection experiments [21, 164].

The rate for direct detection of galactic neutralinos, integrated over deposited energy assuming no energy threshold, is

$$R = \sum_{i} N_{i} n_{\chi} \langle \sigma_{i\chi} v \rangle, \qquad (30.9)$$

where  $N_i$  is the number of nuclei of species *i* in the detector,  $n_{\chi}$  is the local galactic neutralino number density,  $\sigma_{i\chi}$  is the neutralino-nucleus elastic cross section, and the angular brackets denote an average over *v*, the neutralino speed relative to the detector as described in Chapter 16.

In DarkSUSY, the basic quantities computed are the neutralino-nucleon cross sections, which are free of complications related to nuclear structure, and various experimental details like energy threshold, efficiencies etc. However, as a crude estimate of the expected rates in realistic detectors, the total neutralino-nucleus scattering rates can be obtained for <sup>76</sup>Ge,  $Al_2O_3$  (sapphire) and NaI. Usually, it is the spin-independent interaction that gives the most important contribution in target materials such as Na, Cs, Ge, I, or Xe, due to the enhancement caused by the coherence of all nucleons in the target nucleus.

The neutralino-nucleus elastic cross section can be written as

$$\sigma_{i\chi} = \frac{1}{4\pi v^2} \int_0^{4m_{i\chi}^2 v^2} \mathrm{d}q^2 G_{i\chi}^2(q^2), \qquad (30.10)$$

where  $m_{i\chi}$  is the neutralino-nucleus reduced mass, q is the momentum transfer and  $G_{i\chi}(q^2)$  is the effective neutralino-nucleus vertex. We write

$$G_{i\chi}^2(q^2) = A_i^2 F_S^2(q^2) G_S^2 + 4\Lambda_i^2 F_A^2(q^2) G_A^2,$$
(30.11)

which shows the coherent enhancement factor  $A_i^2$  for the spin-independent cross section (often  $\Lambda_i^2$  is written as  $\lambda^2 J(J+1)$ ). We assume gaussian nuclear form factors [90]

$$F_S(q^2) = F_A(q^2) = \exp(-q^2 R_i^2 / 6\hbar^2), \qquad (30.12)$$

$$R_i = (0.3 + 0.89A_i^{1/3}) \text{fm}, \qquad (30.13)$$

which should provide us with a good approximation of the integrated detection rate [165, 166], in which we are only interested. (To obtain the differential rate with reasonable accuracy, better approximations are needed [167].)

#### CHAPTER 30. THE MINIMAL SUPERSYMMETRIC STANDARD MODEL

Using heavy-squark effective lagrangians [168, 169, 170, 74, 171], we get

$$G_S = \sum_{q=u,d,s,c,b,t} \langle \bar{q}q \rangle \left( \sum_{h=H_1,H_2} \frac{g_{h\chi\chi}g_{hqq}}{m_h^2} - \frac{1}{2} \sum_{k=1}^6 \frac{g_{L\tilde{q}_k\chi q}g_{R\tilde{q}_k\chi q}}{m_{\tilde{q}_k}^2} \right)$$
(30.14)

 $\quad \text{and} \quad$ 

$$G_A = \sum_{q=u,d,s} \Delta q \left( \frac{g_{Z\chi\chi}g_{Zqq}}{m_Z^2} + \frac{1}{8} \sum_{k=1}^6 \frac{g_{L\tilde{q}_k\chi q}^2 + g_{R\tilde{q}_k\chi q}^2}{m_{\tilde{q}_k}^2} \right).$$
(30.15)

The g's are elementary vertices involving the particles indicated by the indices, and they read

$$g_{h\chi\chi} = \begin{cases} (gZ_{\chi2} - g_yZ_{\chi1}) (-Z_{\chi3}\cos\alpha + Z_{\chi4}\sin\alpha), & \text{for } H_1, \\ (gZ_{\chi2} - g_yZ_{\chi1}) (Z_{\chi3}\sin\alpha + Z_{\chi4}\cos\alpha), & \text{for } H_2, \end{cases}$$
(30.16)

$$g_{hqq} = \begin{cases} -Y_q \cos \alpha / \sqrt{2}, & \text{for } H_1, \\ +Y_q \sin \alpha / \sqrt{2}, & \text{for } H_2, \end{cases}$$
(30.17)

$$g_{Z\chi\chi} = \frac{g}{2\cos\theta_W} \left( Z_{\chi3}^2 - Z_{\chi4}^2 \right)$$
(30.18)

$$g_{Zqq} = -\frac{g}{2\cos\theta_W}T_{3q}, \qquad (30.19)$$

$$g_{L\tilde{q}_k\chi q} = g_{LL}\Gamma^{kq}_{QL} + g_{RL}\Gamma^{kq}_{QR}, \qquad (30.20)$$

$$g_{R\tilde{q}_k\chi q} = g_{LR}\Gamma^{kq}_{QL} + g_{RR}\Gamma^{kq}_{QR}, \qquad (30.21)$$

with

$$g_{LL} = -\frac{1}{\sqrt{2}} \left( T_{3q} g Z_{\chi 2} + \frac{1}{3} g_y Z_{\chi 1} \right), \qquad (30.22)$$

$$g_{RR} = \sqrt{2}e_q g_y Z_{\chi 1}, \tag{30.23}$$

$$g_{LR} = g_{RL} = \begin{cases} -Y_q Z_{\chi 3}, & \text{for } q = u, c, t, \\ -Y_q Z_{\chi 4}, & \text{for } q = d, s, b, \end{cases}$$
(30.24)

and

$$Y_q = \begin{cases} m_q/v_2, & \text{for } q = u, c, t, \\ m_q/v_1, & \text{for } q = d, s, b. \end{cases}$$
(30.25)

Defining (N = n, p)

$$f_{Tq}^{N} \equiv \frac{\langle N | m_{q} \bar{q} q | N \rangle}{m_{N}}, \qquad (30.26)$$

we take in  $\mathsf{DarkSUSY}$  the numerical values [172]

$$\begin{aligned} f_{Tu}^{p} &= 0.023, \qquad f_{Td}^{p} = 0.034, \\ f_{Tc}^{p} &= 0.0595, \qquad f_{Ts}^{p} = 0.14, \\ f_{Tt}^{p} &= 0.0595, \qquad f_{Tb}^{p} = 0.0595 \end{aligned} \tag{30.27}$$

 $\quad \text{and} \quad$ 

$$\begin{aligned}
f_{Tu}^{n} &= 0.019, & f_{Td}^{n} = 0.041, \\
f_{Tc}^{n} &= 0.0592 & f_{Ts}^{n} = 0.14, \\
f_{Tt}^{n} &= 0.0592, & f_{Tb}^{n} = 0.0592.
\end{aligned} \tag{30.28}$$

For the quark contributions to the nucleon spin we take [173]

$$\Delta u = 0.77, \qquad \Delta d = -0.40, \qquad \Delta s = -0.12.$$
 (30.29)

However, the older set of data [174]

$$\Delta u = 0.77, \qquad \Delta d = -0.49, \qquad \Delta s = -0.15$$
 (30.30)

can optionally be used.

Moreover, we take for the  $\Lambda$  factors

$$\Lambda_{\rm Al}^2 = 0.087, \qquad \Lambda_{\rm Na}^2 = 0.041 \quad \text{and} \quad \Lambda_{\rm I}^2 = 0.007,$$
(30.31)

according to the odd-group model [175].

# 30.8 mssm/ge: General SUSY model setup: masses, vertices etc

#### 30.8.1 Supersymmetric model

We will here review the definition of the MSSM as given in [1].

#### 30.8.1.1 Parameters

In our notation, the superpotential and the soft supersymmetry-breaking scalar potential minimal supersymmetric standard model (MSSM) with R-parity conservation [130, 131, 176] read respectively

$$W = \epsilon_{ij} \left( -\hat{\mathbf{e}}_{R}^{*} \mathbf{h}_{E} \hat{\mathbf{l}}_{L}^{i} \hat{H}_{1}^{j} - \hat{\mathbf{d}}_{R}^{*} \mathbf{h}_{D} \hat{\mathbf{q}}_{L}^{i} \hat{H}_{1}^{j} + \hat{\mathbf{u}}_{R}^{*} \mathbf{h}_{U} \hat{\mathbf{q}}_{L}^{i} \hat{H}_{2}^{j} - \mu \hat{H}_{1}^{i} \hat{H}_{2}^{j} \right), \qquad (30.32)$$

$$V_{\text{soft}} = \epsilon_{ij} \left( -\tilde{\mathbf{e}}_{R}^{*} \mathbf{A}_{E} \mathbf{h}_{E} \tilde{\mathbf{l}}_{L}^{i} H_{1}^{j} - \tilde{\mathbf{d}}_{R}^{*} \mathbf{A}_{D} \mathbf{h}_{D} \tilde{\mathbf{q}}_{L}^{i} H_{1}^{j} + \tilde{\mathbf{u}}_{R}^{*} \mathbf{A}_{U} \mathbf{h}_{U} \tilde{\mathbf{q}}_{L}^{i} H_{2}^{j} - B \mu H_{1}^{i} H_{2}^{j} \right)$$

$$+ h.c.)$$

$$+ H_{1}^{i*} m_{1}^{2} H_{1}^{i} + H_{2}^{i*} m_{2}^{2} H_{2}^{i}$$

$$+ \tilde{\mathbf{q}}_{L}^{i*} \mathbf{M}_{Q}^{2} \tilde{\mathbf{q}}_{L}^{i} + \tilde{\mathbf{l}}_{L}^{i*} \mathbf{M}_{L}^{2} \tilde{\mathbf{l}}_{L}^{i} + \tilde{\mathbf{u}}_{R}^{*} \mathbf{M}_{U}^{2} \tilde{\mathbf{u}}_{R} + \tilde{\mathbf{d}}_{R}^{*} \mathbf{M}_{D}^{2} \tilde{\mathbf{d}}_{R} + \tilde{\mathbf{e}}_{R}^{*} \mathbf{M}_{E}^{2} \tilde{\mathbf{e}}_{R}. \qquad (30.33)$$

Here *i* and *j* are SU(2) indices ( $\epsilon_{12} = +1$ ), h's, A's and M's are  $3 \times 3$  matrices in generation space, and the other boldface letters are vectors in generation space.

The current version of DarkSUSY uses only a restricted set of parameters. Namely the number of free parameters (a grand total of 124 [177]) is reduced by setting the off-diagonal elements of the **A**'s and **M**'s to zero and imposing CP conservation (except in the CKM matrix).

#### 30.8.1.2 Mass spectrum

For easy reference, we now give the particle mass matrices, together with our convention for the mixing matrices.

Concerning the Higgs sector, we choose as independent parameters  $\tan \beta$  and the mass  $m_A$  of the CP-odd Higgs boson. The code provides five options for the calculation of the Higgs masses:

higloop=0: tree level formulas; higloop=1: the effective potential approach in [178, 179, 180] (correcting the sign of  $\mu$  in eq. (4) of [180]); higloop=2: the effective potential approach in [181] with addition of D-terms and correction of some signs and numerical factors; higloop=3: the analytical approximations to the RGE-improved effective potential in [182]; higloop=4: the pole mass calculation in [183]; higloop=5: FeynHiggs (requires FeynHiggs to be installed, default) [184]

The masses of the Higgs bosons are obtained from

$$\mathcal{M}_{H}^{2} = \begin{pmatrix} m_{Z}^{2}\cos^{2}\beta + m_{A}^{2}\sin^{2}\beta + \Delta_{11} & -\sin\beta\cos\beta(m_{Z}^{2} + m_{A}^{2}) + \Delta_{12} \\ -\sin\beta\cos\beta(m_{Z}^{2} + m_{A}^{2}) + \Delta_{21} & m_{Z}^{2}\sin^{2}\beta + m_{A}^{2}\cos^{2}\beta + \Delta_{22} \end{pmatrix}$$
(30.34)

$$m_{H^{\pm}}^2 = m_A^2 + m_W^2 + \Delta_{\pm}.$$
(30.35)

		Higgs bose	n	
Channel	$H_{1}^{0}$	$H_{2}^{0}$	$H_{3}^{0}$	$H^+$
i=	j=1	j=2	j=3	j=4
1	$c\bar{c}$	$c\bar{c}$	$c\bar{c}$	$u \overline{d}$
2	$bar{b}$	$bar{b}$	$bar{b}$	$u\bar{s}$
3	$tar{t}$	$t \overline{t}$	$t\bar{t}$	$u \overline{b}$
4	$\tau^+\tau^-$	$\tau^+\tau^-$	$\tau^+\tau^-$	$c \bar{d}$
5	$W^+W^-$	$W^+W^-$	_	$c\bar{s}$
6	$Z^{0}Z^{0}$	$Z^{0}Z^{0}$	_	$c\bar{b}$
7	_	$H_{1}^{0}H_{1}^{0}$	_	$t ar{d}$
8	$H_{2}^{0}H_{2}^{0}$	_	_	$t\bar{s}$
9	$H_{3}^{0}H_{3}^{0}$	$H_{3}^{0}H_{3}^{0}$	_	$tar{b}$
10	$H^+H^-$	$H^+H^-$	_	$\nu_e e^+$
11	_	_	$ZH_1^0$	$ u_{\mu}\mu^{+}$
12	_	_	$ZH_2^0$	$\nu_{\tau}\tau^+$
13	$ZH_3^0$	$ZH_3^0$	_	$W^{+}H_{1}^{0}$
14	$W^+H^-/W^-H^+$	$W^+H^-/W^-H^+$	$W^+H^-/W^-H^+$	$W^{+}H_{2}^{0}$
15	$\mu^+\mu^-$	$\mu^+\mu^-$	$\mu^+\mu^-$	$W^{+}H_{3}^{0}$
16	$s\bar{s}$	$s\bar{s}$	$s\bar{s}$	_
17	gg	gg	gg	_
18	$\gamma\gamma$	$\gamma\gamma$	$\gamma\gamma$	_
19	$Z^0\gamma$	$Z^0\gamma$	$Z^0\gamma$	_
20	${\widetilde f}{\widetilde f}'$	${ ilde f}{ ilde f}'$	$ ilde{f} ilde{f}'$	${\widetilde f}{\widetilde f}'$

Table 30.2: Higgs partial widths hdwidth(i,j). Index i refers to the decay channel and index j to the Higgs boson. All widths are given in GeV. Note that typically we have that  $m_{H_2} < m_{H_3} < m_{H^+} < m_{H_1}$  so many of these decay channels are not kinematically allowed, but included for completeness. If the HDECAY interface is used, the channels where  $m_{H_2} < m_{H_3} < m_{H^+} < m_{H_1}$  is not satisfied are not included. Channels 16–19 are only included if HDECAY is used.

The quantities  $\Delta_{ij}$  and  $\Delta_{\pm}$  are the one-loop radiative corrections, calculated acording to the value of higloop as described above. Diagonalization of  $\mathcal{M}_{H}^{2}$  gives the two CP-even Higgs boson masses,  $m_{H_{1,2}}$ , and their mixing angle  $\alpha$  ( $-\pi/2 < \alpha < 0$ ). For higloop=4, the pole masses are then obtained solving  $m_{H_{i}}^{2\text{pole}} = m_{H_{i}}^{2} + \prod_{ii}(m_{H_{i}}^{2\text{pole}}) - \prod_{ii}(0)$ , where  $\prod_{ii}(p^{2})$  is  $H_{i}H_{i}$  the self-energy. In this case,  $m_{H_{3}}$  is the pole mass and  $m_{A}$  is the running mass.

The Higgs widths are calculated at tree level, but with QCD corrections. The decays to supersymmetric particles are also included in the total width, so the sum of the partial widths in Table 30.2 does not necessarily sum up to the total width given in width(k). The loop corrections are also available via an interface to HDECAY.

The neutralinos  $\tilde{\chi}_i^0$  are linear combinations of the neutral gauginos  $\tilde{B}$ ,  $\tilde{W}_3$  and of the neutral higgsinos  $\tilde{H}_1^0$ ,  $\tilde{H}_2^0$ . In this basis, we write their mass matrix as

$$\mathcal{M}_{\tilde{\chi}^{0}_{1,2,3,4}} = \begin{pmatrix} M_{1} & 0 & -m_{Z}s_{W}c_{\beta} & +m_{Z}s_{W}s_{\beta} \\ 0 & M_{2} & +m_{Z}c_{W}c_{\beta} & -m_{Z}c_{W}s_{\beta} \\ -m_{Z}s_{W}c_{\beta} & +m_{Z}c_{W}c_{\beta} & \delta_{33} & -\mu \\ +m_{Z}s_{W}s_{\beta} & -m_{Z}c_{W}s_{\beta} & -\mu & \delta_{44} \end{pmatrix},$$
(30.36)

with  $c_W = \cos \theta_W$ ,  $s_W = \sin \theta_W$ ,  $c_\beta = \cos \beta$ , and  $s_\beta = \sin \beta$ . Here  $\delta_{33}$  and  $\delta_{44}$  are radiative corrections important when two higgsinos are close in mass. Their explicit expressions are from ref. [136]. To neglect these radiative corrections set neuloop=0 instead of neuloop=1 (default). The neutralino mass eigenstates are written as

$$\tilde{\chi}_i^0 = N_{i1}\tilde{B} + N_{i2}\tilde{W}^3 + N_{i3}\tilde{H}_1^0 + N_{i4}\tilde{H}_2^0.$$
(30.37)

The phases of  $N_{ij}$  are chosen so that the neutralino masses  $m_{\tilde{\chi}_i^0} \geq 0$ .

The charginos are linear combinations of the charged gauge bosons  $\tilde{W}^{\pm}$  and of the charged higgsinos  $\tilde{H}_1^-$ ,  $\tilde{H}_2^+$ . Their mass matrix,

$$\mathcal{M}_{\tilde{\chi}^{\pm}} = \begin{pmatrix} M_2 & \sqrt{2}m_W \sin\beta \\ \sqrt{2}m_W \cos\beta & \mu \end{pmatrix}, \qquad (30.38)$$

is diagonalized by the following linear combinations

$$\tilde{\chi}_i^- = U_{i1}W^- + U_{i2}H_1^-, \qquad (30.39)$$

$$\tilde{\chi}_{i}^{+} = V_{i1}\tilde{W}^{+} + V_{i2}\tilde{H}_{1}^{+}.$$
(30.40)

We choose  $\det(U) = 1$  and  $U^* \mathcal{M}_{\tilde{\chi}^{\pm}} V^{\dagger} = \operatorname{diag}(m_{\tilde{\chi}_1^{\pm}}, m_{\tilde{\chi}_2^{\pm}})$  with non-negative chargino masses  $m_{\tilde{\chi}^{\pm}} \geq 0$ .

 $m_{\tilde{\chi}_i^{\pm}} \ge 0.$ When discussing the squark mass matrix including mixing, it is convenient to choose a basis where the squarks are rotated in the same way as the corresponding quarks in the standard model. We follow the conventions of the particle data group [185] and put the mixing in the left-handed *d*-quark fields, so that the definition of the Cabibbo-Kobayashi-Maskawa matrix is  $\mathbf{K} = \mathbf{V}_1 \mathbf{V}_2^{\dagger}$ , where  $\mathbf{V}_1$  ( $\mathbf{V}_2$ ) rotates the interaction left-handed *u*-quark (*d*-quark) fields to mass eigenstates. For sleptons we choose an analogous basis, but due to the masslessness of neutrinos no analog of the CKM matrix appears.

We then obtain the general  $6 \times 6 \ \tilde{u}$ - and  $\tilde{d}$ -squark mass matrices:

$$\mathcal{M}_{\tilde{u}}^{2} = \begin{pmatrix} \mathbf{M}_{Q}^{2} + \mathbf{m}_{u}^{\dagger}\mathbf{m}_{u} + D_{LL}^{u}\mathbf{1} & \mathbf{m}_{u}^{\dagger}(\mathbf{A}_{U}^{\dagger} - \mu^{*}\cot\beta) \\ (\mathbf{A}_{U} - \mu\cot\beta)\mathbf{m}_{u} & \mathbf{M}_{U}^{2} + \mathbf{m}_{u}\mathbf{m}_{u}^{\dagger} + D_{RR}^{u}\mathbf{1} \end{pmatrix},$$
(30.41)

$$\mathcal{M}_{\tilde{d}}^{2} = \begin{pmatrix} \mathbf{K}^{\dagger} \mathbf{M}_{Q}^{2} \mathbf{K} + \mathbf{m}_{d} \mathbf{m}_{d}^{\dagger} + D_{LL}^{d} \mathbf{1} & \mathbf{m}_{d}^{\dagger} (\mathbf{A}_{D}^{\dagger} - \mu^{*} \tan \beta) \\ (\mathbf{A}_{D} - \mu \tan \beta) \mathbf{m}_{d} & \mathbf{M}_{D}^{2} + \mathbf{m}_{d}^{\dagger} \mathbf{m}_{d} + D_{RR}^{d} \mathbf{1} \end{pmatrix},$$
(30.42)

and the general sneutrino and charged slepton mass matrices

$$\mathcal{M}_{\tilde{\nu}}^2 = \mathbf{M}_L^2 + D_{LL}^{\nu} \mathbf{1} \tag{30.43}$$

$$\mathcal{M}_{\tilde{e}}^{2} = \begin{pmatrix} \mathbf{M}_{L}^{2} + \mathbf{m}_{e}\mathbf{m}_{e}^{\dagger} + D_{LL}^{e}\mathbf{1} & \mathbf{m}_{e}^{\dagger}(\mathbf{A}_{E}^{\dagger} - \mu^{*}\tan\beta) \\ (\mathbf{A}_{E} - \mu\tan\beta)\mathbf{m}_{e} & \mathbf{M}_{E}^{2} + \mathbf{m}_{e}^{\dagger}\mathbf{m}_{e} + D_{RR}^{e}\mathbf{1} \end{pmatrix}.$$
(30.44)

Here

$$D_{LL}^{f} = m_Z^2 \cos 2\beta (T_{3f} - e_f \sin^2 \theta_w), \qquad (30.45)$$

$$D_{RR}^f = m_Z^2 \cos 2\beta e_f \sin^2 \theta_w. \tag{30.46}$$

In the chosen basis,  $\mathbf{m}_u = \text{diag}(m_u, m_c, m_t)$ ,  $\mathbf{m}_d = \text{diag}(m_d, m_s, m_b)$  and  $\mathbf{m}_e = \text{diag}(m_e, m_\mu, m_\tau)$ .

The slepton and squark mass eigenstates  $\tilde{f}_k$  ( $\tilde{\nu}_k$  with k = 1, 2, 3 and  $\tilde{e}_k$ ,  $\tilde{u}_k$  and  $\tilde{d}_k$  with  $k = 1, \ldots, 6$ ) diagonalize the previous mass matrices and are related to the current sfermion eigenstates  $\tilde{\mathbf{f}}_L$  and  $\tilde{\mathbf{f}}_R$  via (a = 1, 2, 3)

$$\tilde{f}_{La} = \sum_{k=1}^{6} \tilde{f}_k \Gamma_{FL}^{*ka},$$
(30.47)

$$\tilde{f}_{Ra} = \sum_{k=1}^{6} \tilde{f}_k \Gamma_{FR}^{*ka}.$$
(30.48)

$\nu_e$	knue,knu(1)	$\gamma$	kgamma	$\tilde{\chi}_i^0$	$kn(i) \ i = 1 \dots 4$	$ \tilde{u}_1 $	ksu(1),ksqu(1)
e	ke,kl(1)	$W^{\pm}$	kw	$\tilde{\chi}_k^{\pm}$	kcha $(k)$ $k = 1, 2$	$\tilde{u}_2$	ksu(2),ksqu(4)
$ u_{\mu}$	knumu,knu(2)	$Z^0$	kz	$\tilde{g}$	kgluin	$  \tilde{d}_1  $	ksd(1),ksqd(1)
$\mu$	kmu,kl(2)	g	kgluon	$\tilde{\nu}_e$	ksnue,ksnu(1)	$\tilde{d}_2$	ksd(2),ksqd(4)
$\nu_{\tau}$	knutau,knu(3)			$\tilde{e}_1$	kse(1),ksl(1)	$\tilde{c}_1$	ksc(1), ksqu(2)
$\tau$	ktau,kl(3)			$\tilde{e}_2$	kse(2),ksl(4)	$\tilde{c}_2$	ksc(2), ksqu(5)
u	ku,kqu(1)	$H^0$	kh1	$\tilde{ u}_{\mu}$	ksnumu,ksnu(2)	$\tilde{s}_1$	kss(1), ksqd(2)
d	kd,kqd(1)	$h^0$	kh2	$\tilde{\mu}_1$	ksmu(1),ksl(2)	$\tilde{s}_2$	kss(2), ksqd(5)
c	kc,kqu(2)	$A^0$	kh3	$\tilde{\mu}_2$	ksmu(2),ksl(5)	$ \tilde{b}_1 $	ksb(1), ksqd(3)
s	ks,kqd(2)	$H^{\pm}$	khc	$\tilde{\nu}_{\tau}$	ksnuta,ksnu(3)	$\tilde{b}_2$	ksb(2),ksqd(6)
b	kb,kqd(3)	$G^0$	kgold0	$ ilde{ au}_1$	kstau(1),ksl(3)	$ \tilde{t}_1 $	kst(1), ksqu(3)
t	kt,kqu(3)	$G^{\pm}$	kgoldc	$ ilde{ au}_2$	kstau(2),ksl(6)	$\tilde{t}_2$	kst(2), ksqu(6)

Table 30.3: Particle codes (synonyms are separated by commas).

The squark and charged slepton mixing matrices  $\Gamma_{UL,R}$ ,  $\Gamma_{DL,R}$  and  $\Gamma_{EL,R}$  have dimension  $6 \times 3$ , while the sneutrino mixing matrix  $\Gamma_{\nu L}$  has dimension  $3 \times 3$ .

This version of DarkSUSY allows only for diagonal matrices  $\mathbf{A}_U$ ,  $\mathbf{A}_D$ ,  $\mathbf{A}_E$ ,  $\mathbf{M}_Q$ ,  $\mathbf{M}_U$ ,  $\mathbf{M}_D$ ,  $\mathbf{M}_E$ , and  $\mathbf{M}_L$ . This ansatz, while not being the most general one, implies the absence of tree-level flavor changing neutral currents in all sectors of the model. In this case, the squark mass matrices can be diagonalized analytically. For example, for the top squark one has, in terms of the top squark mixing angle  $\theta_{\tilde{t}}$ ,

$$\Gamma_{UL}^{\tilde{t}_1\tilde{t}} = \Gamma_{UR}^{\tilde{t}_2\tilde{t}} = \cos\theta_{\tilde{t}}, \qquad \Gamma_{UL}^{\tilde{t}_2\tilde{t}} = -\Gamma_{UR}^{\tilde{t}_1\tilde{t}} = \sin\theta_{\tilde{t}}.$$
(30.49)

Special values of the sfermion masses can be set with the parameters msquarks, and msleptons. If msquarks=msleptons=0, the sfermion masses are obtained with the diagonalization described above. If msquarks>0 (or msleptons>0), all squark masses are set to msquarks (or all slepton masses to msleptons). Finally, if msquarks<0 (or msleptons<0), the squark (or slepton) masses are set equal to the neutralino mass but never less than |msquarks| (or |msleptons|). This is to provide the lightest possible sfermions compatible with a neutralino LSP. In all of these cases, there is no mixing between sfermions.

The particle masses are available in an array mass(p), where p is the particle code from table ??. Similarly, particle decay width are available as width(p), but currently only the width of the Higgs bosons are calculated, the other particles having fictitious widths of 1 or 5 GeV (for the sole purpose of regularizing annihilation amplitudes close to poles).

#### 30.8.1.3 Three-particle vertices

We define three-particle vertices  $gl(i,j,k) = g_{ijk}^L$  and  $gr(i,j,k) = g_{ijk}^R$  as follows. We adopt the convention that the order of the particles in the indices is the order in which they appear in the corresponding lagrangian term, so the last particle is always entering. If there are charged particles in the vertex, they are both assumed positively charged, and the particle that exits the vertex is indexed before the particle that enters.

• Three scalar bosons:

$$\mathcal{L}_{\text{int}} = g_{\phi_i \phi_j \phi_k} m_W \phi_i \phi_j \phi_k \tag{30.50}$$

where  $\phi_i$  is a Higgs or a Goldstone boson. In this case, g|=gr=g. Available vertices are  $\phi_i\phi_j\phi_k = H_i^0H_j^0H_k^0$ ,  $H_i^0H^-H^+$ ,  $H_i^0A^0A^0$ ,  $H_i^0G^0G^0$ ,  $H_i^0G^-G^+$ ,  $H_i^0G^-H^+$ ,  $H_i^0G^-G^+$ ,  $A^0G^-H^+$ ,  $A^0G^0H_i^0$ , and permutations.

• Two scalar and one vector bosons:

$$\mathcal{L}_{\text{int}} = g_{V\phi_1\phi_2} V^{\mu} \phi_1 i \overleftarrow{\partial}_{\mu} \phi_2.$$
(30.51)

Available vertices are  $V\phi_1\phi_2 = Z^0H_i^0A^0$ ,  $Z^0H^-H^+$ ,  $\gamma H^-H^+$ ,  $W^-H^+A^0$ ,  $W^-H^+H_i^0$ , and permutations.

• One scalar and two vector bosons:

$$\mathcal{L}_{\rm int} = g_{\phi V_1 V_2} m_W g_{\mu\nu} \phi V_1^{\mu} V_2^{\nu} \tag{30.52}$$

Available vertices are  $\phi V_1 V_2 = H_i^0 W^- W^+$ ,  $H_i^0 Z^0 Z^0$ .

• Three vector bosons:

$$ig_{V_1V_2V_3}\left[(k_1 - k_3)_{\nu}g_{\mu\lambda} + (k_3 - k_2)_{\mu}g_{\lambda\nu} + (k_2 - k_1)g_{\mu\nu}\right]$$
(30.53)

 $\rightarrow$ 

with all momenta incoming and assigned as  $V_1^{\mu}(k_1)$ ,  $V_2^{\nu}(k_2)$  and  $V_3^{\lambda}(k_3)$ . Available vertices are  $Z^0W^-W^+$  and  $\gamma W^-W^+$ .

• One scalar boson and two Dirac fermions:

$$\mathcal{L}_{\text{int}} = \phi \overline{\psi}_1 (g^L_{\phi\psi_1\psi_2} P_L + g^R_{\phi\psi_1\psi_2} P_R) \psi_2 \tag{30.54}$$

Available vertices are  $\phi \psi_1 \psi_2 =$ 

• One vector boson and two Dirac fermions:

$$\mathcal{L}_{\text{int}} = V_{\mu} \overline{\psi}_1 \gamma^{\mu} (g^L_{V\psi_1\psi_2} P_L + g^R_{V\psi_1\psi_2} P_R) \psi_2$$
(30.55)

Available vertices are  $V\psi_1\psi_2 =$ 

• One scalar boson, one Dirac and one Majorana fermion:

$$\mathcal{L}_{\text{int}} = \phi \overline{\psi} (g^L_{\phi\psi\chi} P_L + g^R_{\phi\psi\chi} P_R) \chi$$
(30.56)

Available vertices are  $\phi \psi \chi =$ 

• One vector boson, one Dirac and one Majorana fermion:

$$\mathcal{L}_{\rm int} = V_{\mu} \overline{\psi} \gamma^{\mu} (g_{V\psi\chi}^L P_L + g_{V\psi\chi}^R P_R) \chi \tag{30.57}$$

Available vertices are  $V\psi\chi =$ 

• One scalar boson and two Majorana fermions:

$$\mathcal{L}_{\text{int}} = \tag{30.58}$$

Available vertices are...

• One vector boson and two Majorana fermions:

$$\mathcal{L}_{\text{int}} = \tag{30.59}$$

Explicit expressions for the coupling constants  $g_{ijk}$  can be obtained in [130, 131, 176], with radiative corrections to trilinear scalar couplings in [186]. We have rederived from the superpotential all vertices we have implemented.

Implemented vertices: those listed above plus  $Z^0 W^{\pm} W^{\mp}$ ,  $Z^0 H_i^0 H_i^0$ ,  $W^{\pm} H^{\mp} A^0$ ,  $W^{\pm} H^{\mp} H_i^0$ ,  $H_i^0 W^{\pm} W^{\mp}$ ,  $H_i^0 Z^0 Z^0$ ,  $Z^0 A^0 H$ ,  $H_i^0 A^0 A^0$ ,  $A^0 ff$ ,  $H_i^0 ff$ ,  $Z^0 ff$ ,  $Z^0 \tilde{\chi}^0 \tilde{\chi}^0$ ,  $H_i^0 \tilde{\chi}^0 \tilde{\chi}^0$ ,  $Z^0 \tilde{\chi}^0 \tilde{\chi}^0$ ,  $W^{\mp} \tilde{\chi}^0 \tilde{\chi}^{\pm}$ ,  $H^{\mp} \tilde{\chi}^0 \tilde{\chi}^{\pm}$ ,  $\tilde{q} \tilde{g} q$ ,  $\tilde{f} \tilde{\chi}^0 f$ ,  $H_i^0 \tilde{\chi}^{\pm} \tilde{\chi}^{\mp}$ ,  $A^0 \tilde{\chi}^{\pm} \tilde{\chi}^{\mp}$ ,  $W^{\pm} ff'$ ,  $H^{\pm} ff'$ ,  $\gamma W^{\pm} W^{\mp}$ ,  $\gamma H^{\pm} H^{\mp}$ ,  $Z^0 \tilde{\chi}^{\pm} \tilde{\chi}^{\mp}$ ,  $\gamma \tilde{\chi}^{\pm} \tilde{\chi}^{\mp}$ ,  $\gamma ff$ , GHH, GGH,  $G^{\mp} \tilde{\chi}^0 \tilde{\chi}^{\pm}$ .

#### **30.8.2** General supersymmetry – routines

Input parameters, options, results, etc. are contained in common blocks in the file dsmssm.h, which the user has to include. The input parameters are (a = 1, 2, 3)

 $\begin{array}{ll} \mathsf{ma}=m_A, & \mathsf{tanbe}=\tan\beta, & \mathsf{mu}=\mu, & \mathsf{m1}=M_1, \\ \mathsf{m2}=M_2, & \mathsf{m3}=M_3, & \mathsf{asofte}(a)=A_{Eaa}, & \mathsf{asoftu}(a)=A_{Uaa}, \\ \mathsf{asoftd}(a)=A_{Daa}, & \mathsf{mass2q}(a)=M_{Qaa}^2, & \mathsf{mass21}(a)=M_{Laa}^2, & \mathsf{mass2u}(a)=M_{Uaa}^2, \\ \mathsf{mass2d}(a)=M_{Daa}^2, & \mathsf{mass2e}(a)=M_{Eaa}^2. \end{array}$ The options are (see previous subsections for a description)

higloop choice of tree-level or radiatively corrected Higgs boson masses; neuloop choice of tree-level or radiatively corrected neutralino masses;

msquarks, msleptons choice of squark and slepton masses.

To initialize DarkSUSY for a new model, you should call

#### subroutine dsmodelsetup(unphys,warning)

Purpose:		To calculate the particle spectrum, widths and couplings.
Output:		
unphys	i	non-zero if the model is unphysical
warning	i	non-zero if (typically the Higgs) code has issued a warning.

which calculates couplings, masses and some basic cross sections.

The following subroutines specify the values of the model parameters, and read/write them to a file. The user should create his own versions by editing a copy of them. Please call them with a different name.

subroutine dsgive model(mu,m2,ma,tanbe,msq,atm,abm)

Purpose:		Set the MSSM parameters as specified by the arguments.
Inputs:		
mu	r8	The $\mu$ parameter in GeV.
m2	r8	The $M_2$ parameter in GeV.
ma	r8	The mass of the CP-odd Higgs boson, $m_A$ in GeV.
tanbe	r8	an eta.
msq	r8	Sets $\mathbf{M}^Q$ , etc. to a common mass scale $m_0$ in GeV.
atm	r8	Sets $A_t$ in units of $m_0$ (range: $-3 - 3$ ).
abm	r8	Sets $A_b$ in units of $m_0$ (range: $-3 - 3$ ).

There are also routines to setup more general MSSM models, the currently most general being dsgive\_model25.

The following subroutines are useful in the analysis.

#### subroutine wspctm(unit)

Davaan o o o .		White the particle mass spectrum and mining matrices to unit unit
Purpose:		write the particle mass spectrum and mixing matrices to unit unit.
Innuts		
inp aco.		
unit	i	Unit number to write to.

#### subroutine wvertx(unit)

Purpose:		Write all non-vanishing three-particle vertices to unit unit.
Inputs:		
unit	i	Unit number to write to.

#### subroutine wunph(unit)

*Purpose:* Write the reason for which the model is not physically acceptable (tachyons, etc.) to unit unit.

#### Inputs:

unit i Unit number to write to.

subroutine we	xcl(u	nit)
Purpose:		Write the reason(s) for which the model is experimentally excluded to unit unit.
Inputs: unit	i	Unit number to write to.
subroutine <b>dsv</b>	whwa	ar(unit)
Purpose:		Write the reason(s) for which the Higgs calculation issued warnings to unit unit.

#### Inputs: unit i Unit number to write to.

# 30.9 mssm/ge cmssm:

cMSSM interface (Isasugra)

#### 30.9.1 mSUGRA (ISASUGRA) interface to DarkSUSY

If Isasugra is available, DarkSUSY can use Isasugra to generate mSUGRA models. In src\_modles/mssm/ge\_cmssm, routines are available to transfer the mSUGRA parameters from DarkSUSY to Isasugra, call Isasugra and then transfer back the results to DarkSUSY. The philosophy of this interface is that whenever a user uses Isasugra, we should use all the results of Isasugra also in DarkSUSY. That means that instead of calculating the mass spectrum from the low-energy parameters obtained from Isasugra, we extract the masses and mixings from Isasugra.

# 30.10 mssm/ge\_slha: SUSY Les Houches Accord interface

#### 30.10.1 SUSY Les Houches Accord

DarkSUSY includes routines to read and write SUSY Les Houches Accord [134, 135] files (SLHA files). This is done with the help of SLHALIB by T. Hahn [187].

DarkSUSY will write SLHA2 files and expect to get SLHA2 as well. The implementation right now takes a middle path between dumping everything or just a minimal set of inputs to the SLHA2 file. This choice was made to make most other SLHA2-aware programs able to exchange SLHA2 files with DarkSUSY.

However, we have not made a careful testing with lots of other codes, so if you try this out, please let us know if there are some things that don't work or could work better.

### 30.11 mssm/ini: Initialization routines

#### **30.11.1** Initialization routines

Before DarkSUSY is used for some calculations, it needs to be initialized. This is done with a call to **dsinit**, which in turn calls the particle-specific initialization routine dsinit\_module that resides in each module. This means that the call to **dsinit** should be the first call in any program using DarkSUSY. Any calls the user makes to other routines, either to calculte things or select a different model (e.g. a different halo model) should come after the call to **dsinit**.

In the mssm module, the routine dsinit\_module that all standard parameters are defined, such as standard model parameters and particle codes. It also sets more particle-specific defaults for Dark-SUSY routines where this is available, via calls to routines such as dsddset mssm and dsanset mssm.

# 30.12 mssm/rd: Relic density

#### 30.12.1 Relic density of neutralinos

In the folder **mssm/rd**, we provide one of the two interface functions that are necessary for the **core** library to solve the Boltzmann equation for any cold dark matter particle and hence calculate its relic density:<sup>‡</sup> dsrdparticles determines which particles can coannihilate (based on their mass differences) and puts these particles into a common block for the annihilation rate routines (dsanwx); it also checks where we have resonances and thresholds and adds these to an array, which is passed to the relic density routines. The relic density routines then use this knowledge to make sure the tabulation of the cross section and the integrations are performed correctly at these difficult points.

For convenience, we include one further routine in this folder: dsrdwrate writes the invariant rate to a specified unit. This is mostly useful for debugging purposes.

#### **30.12.2** Internal degrees of freedom

In Section 20.1.2, we have reviewed the standard way of calculating the relic density. Here, we add some comments which are specific to the implementation of the effective invariant rate in the mssm module

If we look at Eqs. (20.34) and (20.39) we see that we have a freedom on how to treat particles degenerate in mass, e.g. a chargino can be treated either

- a) as two separate species  $\chi_i^+$  and  $\chi_i^-$ , each with internal degrees of freedom  $g_{\chi^+} = g_{\chi^-} = 2$ , or,
- b) as a single species  $\chi_i^{\pm}$  with  $g_{\chi_i^{\pm}} = 4$  internal degrees of freedom.

Of course the two views are equivalent, we just have to be careful including the  $g_i$ 's consistently whichever view we take. In a), we have the advantage that all the  $W_{ij}$  that enter into Eq. (20.34) enter as they are, i.e. without any correction factors for the degrees of freedom. On the other hand we get many terms in the sum that are identical and we need some book-keeping machinery to avoid calculating identical terms more than once. On the other hand, with option b), the sum over  $W_{ij}$  in Eq. (20.34) is much simpler only containing terms that are not identical (except for the trivial identity  $W_{ij} = W_{ji}$  which is easily taken care of). However, the individual  $W_{ij}$  will be some linear combinations of the more basic  $W_{ij}$  entering in option a), where the coefficients have to be calculated for each specific type of initial condition.

Below we will perform this calculation to show how the  $W_{ij}$  look like in option b) for different initial states. We will use a prime on the  $W_{ij}$  when they refer to these combined states to indicate the difference.

#### 30.12.2.1 Neutralino-chargino annihilation

The starting point is Eq. (20.34) which we will use to define the  $W_{ij}$  in option b) such that  $W_{\text{eff}}$  is the same as in option a). Eq. (20.39) is then guaranteed to be the same in both cases since the sum in the denominator is linear in  $g_i$ .

Now consider annihilation between  $\chi_i^0$  and  $\chi_c^+$  or  $\chi_c^-$ . The corresponding terms in Eq. (20.34)

 $<sup>\</sup>ddagger$  The other, dsanwx, returns the effective invariant rate and resides in mssm/an/.

does for option a) read

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W_{ij} = \frac{p_{ic}}{p_{11}} \frac{2 \cdot 2}{2^2} \left[ W_{\chi_i^0 \chi_c^+} + W_{\chi_i^0 \chi_c^-} + \underbrace{W_{\chi_c^+ \chi_i^0}}_{W_{\chi_i^0 \chi_c^+}} + \underbrace{W_{\chi_c^- \chi_i^0}}_{W_{\chi_i^0 \chi_c^-}} \right] \\ = 2 \frac{p_{ic}}{p_{11}} \left[ W_{\chi_i^0 \chi_c^+} + \underbrace{W_{\chi_i^0 \chi_c^-}}_{W_{\chi_i^0 \chi_c^+}} \right] = 4 \frac{p_{ic}}{p_{11}} W_{\chi_i^0 \chi_c^+}$$
(30.60)

For option b), we instead get

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W_{ij} = \frac{p_{ic}}{p_{11}} \frac{2 \cdot 4}{2^2} \left[ W'_{\chi_i^0 \chi_c^{\pm}} + \underbrace{W'_{\chi_c^{\pm} \chi_i^0}}_{W'_{\chi_i^0 \chi_c^{\pm}}} \right] = 4 \frac{p_{ic}}{p_{11}} W'_{\chi_i^0 \chi_c^{\pm}}$$
(30.61)

Comparing Eq. (30.61) and Eq. (30.60) we see that they are indentical if we make the identification

$$W'_{\chi_i^0 \chi_c^\pm} \equiv W_{\chi_i^0 \chi_c^+} \tag{30.62}$$

#### 30.12.2.2 Chargino-chargino annihilation

First consider the case where we include the terms in the sum for which we have annihilation between  $\chi_c^+$  or  $\chi_c^-$  and  $\chi_d^+$  or  $\chi_d^-$  with  $c \neq d$ . In option a), the corresponding terms in Eq. (20.34) reads

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_{i}g_{j}}{g_{1}^{2}} W_{ij}$$

$$= \frac{p_{cd}}{p_{11}} \frac{2 \cdot 2}{2^{2}} \left[ W_{\chi_{c}^{+}\chi_{d}^{+}} + W_{\chi_{c}^{+}\chi_{d}^{-}} + W_{\chi_{c}^{-}\chi_{d}^{+}} + W_{\chi_{c}^{-}\chi_{d}^{-}} + \frac{W_{\chi_{d}^{+}\chi_{c}^{-}}}{W_{\chi_{c}^{-}\chi_{d}^{-}}} + \frac{W_{\chi_{d}^{+}\chi_{c}^{-}}}{W_{\chi_{c}^{-}\chi_{d}^{-}}} + \frac{W_{\chi_{d}^{-}\chi_{c}^{-}}}{W_{\chi_{c}^{-}\chi_{d}^{-}}} \right]$$

$$= 2 \frac{p_{cd}}{p_{11}} \left[ W_{\chi_{c}^{+}\chi_{d}^{+}} + W_{\chi_{c}^{+}\chi_{d}^{-}} + \frac{W_{\chi_{c}^{-}\chi_{d}^{-}}}{W_{\chi_{c}^{+}\chi_{d}^{-}}} + \frac{W_{\chi_{c}^{-}\chi_{d}^{-}}}{W_{\chi_{c}^{+}\chi_{d}^{-}}} \right]$$

$$= 4 \frac{p_{cd}}{p_{11}} \left[ W_{\chi_{c}^{+}\chi_{d}^{+}} + W_{\chi_{c}^{+}\chi_{d}^{-}} \right]$$
(30.63)

In option b), the corresponding terms would instead read

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W'_{ij} = \frac{p_{cd}}{p_{11}} \frac{4 \cdot 4}{2^2} \left[ W'_{\chi_c^{\pm} \chi_d^{\pm}} + \underbrace{W'_{\chi_d^{\pm} \chi_c^{\pm}}}_{W'_{\chi_c^{\pm} \chi_d^{\pm}}} \right] = 8 \frac{p_{cd}}{p_{11}} W'_{\chi_c^{\pm} \chi_d^{\pm}}$$
(30.64)

Comparing Eq. (30.63) and Eq. (30.64) we see that they are identical if we make the following identification

$$W'_{\chi_c^{\pm}\chi_d^{\pm}} \equiv \frac{1}{2} \left[ W_{\chi_c^{+}\chi_d^{+}} + W_{\chi_c^{+}\chi_d^{-}} \right]$$
(30.65)

For clarity, let's also consider the case where c = d. In option a), the terms in  $W_{\text{eff}}$  are

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W_{ij} = \frac{p_{cc}}{p_{11}} \frac{2 \cdot 2}{2^2} \left[ W_{\chi_c^+ \chi_c^+} + W_{\chi_c^+ \chi_c^-} + \underbrace{W_{\chi_c^- \chi_c^+}}_{W_{\chi_c^+ \chi_c^-}} + \underbrace{W_{\chi_c^- \chi_c^-}}_{W_{\chi_c^+ \chi_c^+}} + \underbrace{W_{\chi_c^- \chi_c^-}}_{W_{\chi_c^+ \chi_c^+}} \right]$$

$$= 2 \frac{p_{cc}}{p_{11}} \left[ W_{\chi_c^+ \chi_c^+} + W_{\chi_c^+ \chi_c^-} \right]$$
(30.66)

In option b), the corresponding term would instead read

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W'_{ij} = \frac{p_{cc}}{p_{11}} \frac{4 \cdot 4}{2^2} W'_{\chi_c^{\pm} \chi_c^{\pm}} + = 4 \frac{p_{cc}}{p_{11}} W'_{\chi_c^{\pm} \chi_c^{\pm}}$$
(30.67)

Comparing Eq. (30.66) and Eq. (30.67) we see that they are identical if we make the following identification

$$W'_{\chi_c^{\pm}\chi_c^{\pm}} \equiv \frac{1}{2} \left[ W_{\chi_c^{+}\chi_c^{+}} + W_{\chi_c^{+}\chi_c^{-}} \right]$$
(30.68)

i.e. the same identification as in the case  $c \neq d$ .

#### 30.12.2.3 Neutralino-sfermion annihilation

For each sfermion we have in total four different states,  $\tilde{f}_1$ ,  $\tilde{f}_2$ ,  $\tilde{f}_1^*$  and  $\tilde{f}_2^*$ . Of these, the  $\tilde{f}_1$  and  $\tilde{f}_2$  in general have different masses and have to be treated separately. Considering only one mass eigenstate  $\tilde{f}_k$ , option a) then means that we treat  $\tilde{f}_k$  and  $\tilde{f}_k^*$  as two separate species with  $g_i = 1$  degree of freedom each, whereas option b) means that we treat them as one species  $\tilde{f}_k'$  with  $g_i = 2$  degrees of freedom. As before, the prime indicates that we mean both the particle and the antiparticle state.

Note, that for squarks we also have the number of colours  $N_c = 3$  to take into account. In option a) we should choose to treat even colour state differently, i.e.  $g_i = 1$ , whereas  $g_i = 6$  in case b). The expressions would be the same as above except that both the expression in a) and b) would be multiplied by the colour factor  $N_c = 3$ . The expression relating case a) and case b) is thus unaffected by this colour factor. Note however, that in option b) we take the average over the squark colours (or in this case calculate it only for one colour. See sections 30.12.2.6 and 30.12.2.7 below for more details.

For option a), Eq. (20.34) then reads

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W_{ij} = \frac{p_{ik}}{p_{11}} \frac{2 \cdot 1}{2^2} \left[ W_{\chi_i^0 \tilde{f}_k} + W_{\chi_i^0 \tilde{f}_k^*} + \underbrace{W_{\tilde{f}_k \chi_i^0}}_{W_{\chi_i^0 \tilde{f}_k}} + \underbrace{W_{\tilde{f}_k^* \chi_i^0}}_{W_{\chi_i^0 \tilde{f}_k^*}} \right]$$
$$= \frac{p_{ik}}{p_{11}} \left[ W_{\chi_i^0 \tilde{f}_k} + \underbrace{W_{\chi_i^0 \tilde{f}_k^*}}_{W_{\chi_i^0 \tilde{f}_k}} \right] = 2 \frac{p_{ik}}{p_{11}} W_{\chi_i^0 \tilde{f}_k}$$
(30.69)

whereas for option b), Eq. (20.34) reads

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W'_{ij} = \frac{p_{ik}}{p_{11}} \frac{2 \cdot 2}{2^2} \left[ W'_{\chi_i^0 \tilde{f'}_k} + \underbrace{W'_{\tilde{f'}_k \chi_i^0}}_{W'_{\chi_i^0 \tilde{f'}_k}} \right] = 2 \frac{p_{ik}}{p_{11}} W'_{\chi_i^0 \tilde{f'}_k}$$
(30.70)

Comparing Eq. (30.70) and Eq. (30.69) we see that they are indentical if we make the identification

$$W'_{\chi_{i}^{0}\tilde{f}'_{k}} \equiv W_{\chi_{i}^{0}\tilde{f}_{k}} \tag{30.71}$$
For clarity, for squarks the corresponding expression would be

$$W'_{\chi^0_i \tilde{q'}_k} \equiv \frac{1}{3} \sum_{a=1}^3 W_{\chi^0_i \tilde{q}^a_k}$$
(30.72)

where a is a colour index.

### 30.12.2.4 Chargino-sfermion annihilation

In option a) the chargino has  $g_i = 2$  and the sfermion has  $g_i = 1$  degrees of freedom, whereas in option b), the chargino has  $g_i = 4$  and the sfermion has  $g_i = 2$  degrees of freedom

For option a), Eq. (20.34) then reads

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_{i}g_{j}}{g_{1}^{2}} W_{ij}$$

$$= \frac{p_{ck}}{p_{11}} \frac{2 \cdot 1}{2^{2}} \left[ W_{\chi_{c}^{+}\tilde{f}_{k}} + W_{\chi_{c}^{+}\tilde{f}_{k}^{*}} + W_{\chi_{c}^{-}\tilde{f}_{k}} + W_{\chi_{c}^{-}\tilde{f}_{k}} + W_{\chi_{c}^{-}\tilde{f}_{k}} + W_{\chi_{c}^{+}\tilde{f}_{k}^{*}} + W_{\tilde{f}_{k}\chi_{c}^{-}} + W_{\tilde{f}_{k}\chi_{c}^{-}} \right]$$

$$= \frac{p_{ck}}{p_{11}} \left[ W_{\chi_{c}^{+}\tilde{f}_{k}} + W_{\chi_{c}^{+}\tilde{f}_{k}^{*}} + \frac{W_{\chi_{c}^{-}\tilde{f}_{k}}}{W_{\chi_{c}^{-}\tilde{f}_{k}}} + \frac{W_{\chi_{c}^{-}\tilde{f}_{k}}}{W_{\chi_{c}^{-}\tilde{f}_{k}^{*}}} \right] = 2 \frac{p_{ck}}{p_{11}} \left[ W_{\chi_{c}^{+}\tilde{f}_{k}} + W_{\chi_{c}^{+}\tilde{f}_{k}^{*}} + \frac{W_{\chi_{c}^{-}\tilde{f}_{k}}}{W_{\chi_{c}^{+}\tilde{f}_{k}^{*}}} \right] = (30.73)$$

In option b), Eq. (20.34) reads

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W'_{ij} = \frac{p_{ck}}{p_{11}} \frac{4 \cdot 2}{2^2} \left[ W'_{\chi_c^{\pm} \tilde{f}'_k} + \underbrace{W'_{\tilde{f}'_k \chi_c^{\pm}}}_{W'_{\chi_c^{\pm} \tilde{f}'_k}} \right] = 4 \frac{p_{ck}}{p_{11}} W'_{\chi_c^{\pm} \tilde{f}'_k}$$
(30.74)

Comparing Eq. (30.74) and Eq. (30.73) we see that they are indentical if we make the identification

$$W'_{\chi_c^{\pm}\tilde{f}'_k} \equiv \frac{1}{2} \left[ W_{\chi_c^{+}\tilde{f}_k} + W_{\chi_c^{+}\tilde{f}_k^{*}} \right]$$
(30.75)

For clarity, for squarks the corresponding expression would be

$$W'_{\chi^{\pm}_{c}\tilde{q'}_{k}} \equiv \frac{1}{2} \frac{1}{3} \sum_{a=1}^{3} \left[ W_{\chi^{+}_{c}\tilde{q}^{a}_{k}} + W_{\chi^{+}_{c}\tilde{q}^{a*}_{k}} \right]$$
(30.76)

where a is a colour index.

### 30.12.2.5 Sfermion-sfermion annihilation

First consider the case where we have annihilation between sfmerions of different types, i.e. annihilation between  $\tilde{f}_k$  or  $\tilde{f}_k^*$  and  $\tilde{f}_l$  or  $\tilde{f}_l^*$ .

For option a), Eq. (20.34) then reads

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W_{ij}$$

$$= \frac{p_{kl}}{p_{11}} \frac{1 \cdot 1}{2^2} \left[ W_{\tilde{f}_k \tilde{f}_l} + W_{\tilde{f}_k \tilde{f}_l^*} + W_{\tilde{f}_k^* \tilde{f}_l} + W_{\tilde{f}_k^* \tilde{f}_l^*} \right]$$

$$= \frac{1}{2} \frac{p_{kl}}{p_{11}} \left[ W_{\tilde{f}_k \tilde{f}_l} + W_{\tilde{f}_k \tilde{f}_l^*} + W_{\tilde{f}_k^* \tilde{f}_l^*} + W_{\tilde{f}_k^* \tilde{f}_l^*} + W_{\tilde{f}_k^* \tilde{f}_l^*} + W_{\tilde{f}_k^* \tilde{f}_l^*} \right]$$

$$(30.77)$$

In option b) we would get

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W'_{ij}$$
  
=  $\frac{p_{kl}}{p_{11}} \frac{2 \cdot 2}{2^2} \left[ W'_{\tilde{f}'_k \tilde{f}'_l} + \underbrace{W'_{\tilde{f}'_l \tilde{f}'_k}}_{W'_{\tilde{f}'_k \tilde{f}'_l}} \right] = 2 \frac{p_{kl}}{p_{11}} \left[ W'_{\tilde{f}'_k \tilde{f}'_l} \right]$  (30.78)

Comparing Eq. (30.78) and Eq. (30.77) we see that they are indentical if we make the identification

$$W'_{\tilde{f'}_k \tilde{f'}_l} \equiv \frac{1}{2} \left[ W_{\tilde{f}_k \tilde{f}_l} + W_{\tilde{f}_k \tilde{f}_l^*} \right]$$
(30.79)

It is easy to show that this relation holds true even if k = l.

### 30.12.2.6 Squark-squark annihilation

Even though we treated sfermion-sfermion annihilation in the previous subsection, squarks have colour which can complicate things, so let's for clarity consider squarks separately.

Let's denote the squarks  $\tilde{q}_k^a$  where *a* is now a colour index. In option a) we will let each colour be a seprate species, which means that  $g_i = 1$  in this case. In option b) we will instead have  $g_i = 6$ . In option a) we would have

$$\begin{split} W_{\text{eff}} &= \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W_{ij} \\ &= \frac{p_{kl}}{p_{11}} \frac{1 \cdot 1}{2^2} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b*}} + W_{\tilde{q}_k^{a*} \tilde{q}_l^b} + W_{\tilde{q}_k^{a*} \tilde{q}_l^b} + W_{\tilde{q}_k^{a*} \tilde{q}_l^{b*}} \right] \\ &+ \underbrace{W_{\tilde{q}_l^a \tilde{q}_k^b}}_{W_{\tilde{q}_k^a \tilde{q}_l^b}} + \underbrace{W_{\tilde{q}_l^a \tilde{q}_k^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} + \underbrace{W_{\tilde{q}_l^{a*} \tilde{q}_k^b}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{11}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{f}_l^b} + W_{\tilde{q}_k^a \tilde{f}_l^{b*}} + \underbrace{W_{\tilde{q}_k^{a*} \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{11}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{f}_l^b} + W_{\tilde{q}_k^a \tilde{f}_l^{b*}} + \underbrace{W_{\tilde{q}_k^{a*} \tilde{q}_l^b}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{11}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{f}_l^b} + W_{\tilde{q}_k^a \tilde{f}_l^{b*}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{11}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b*}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{11}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b*}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{11}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b*}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{11}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b*}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{kl}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b*}} + \underbrace{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}}_{W_{\tilde{q}_k^a \tilde{q}_l^{b*}}} \right] \\ &= \frac{1}{2} \frac{p_{kl}}{p_{kl}} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b$$

In option b) we would get

$$W_{\text{eff}} = \sum_{ij} \frac{p_{ij}}{p_{11}} \frac{g_i g_j}{g_1^2} W'_{ij}$$
  
=  $\frac{p_{kl}}{p_{11}} \frac{6 \cdot 6}{2^2} \left[ W'_{\tilde{q'}_k \tilde{q'}_l} + \underbrace{W'_{\tilde{q'}_l \tilde{q'}_k}}_{W'_{\tilde{q'}_k \tilde{q'}_l}} \right] = 18 \frac{p_{kl}}{p_{11}} \left[ W'_{\tilde{q'}_k \tilde{q'}_l} \right]$  (30.81)

Comparing Eq. (30.81) and Eq. (30.80) we see that they are indentical if we make the identification

$$W'_{\tilde{q'}_k\tilde{q'}_l} \equiv \frac{1}{2} \frac{1}{9} \sum_{a,b=1}^3 \left[ W_{\tilde{q}_k^a \tilde{q}_l^b} + W_{\tilde{q}_k^a \tilde{q}_l^{b*}} \right]$$
(30.82)

i.e. we get the same relation as for other sfermions, the only difference being that we in option b) should also take the average over the colour states.

### 30.12.2.7 Sfermion-squark annihilation

For clarity, if we have annihilation between a non-coloured sfermion and a squark, we would in the same way as in the previous subsection get

$$W'_{\tilde{f}'_{k}\tilde{q}'_{l}} \equiv \frac{1}{2} \frac{1}{3} \sum_{b=1}^{3} \left[ W_{\tilde{f}_{k}\tilde{q}_{l}^{b}} + W_{\tilde{f}_{k}\tilde{q}_{l}^{b*}} \right]$$
(30.83)

#### 30.12.2.8 Summary of degrees of freedom

We have found above the following relations between option b) and option a),

$$\begin{array}{lll} W_{\chi_{i}^{0}\chi_{j}^{\pm}}^{\prime} &\equiv & W_{\chi_{i}^{0}\chi_{j}^{+}} = W_{\chi_{i}^{0}\chi_{j}^{-}} &, \quad \forall \ i = 1, \dots, 4, \ j = 1, 2 \\ W_{\chi_{i}^{\pm}\chi_{j}^{\pm}}^{\prime} &\equiv & \frac{1}{2} \left[ W_{\chi_{i}^{+}\chi_{j}^{+}} + W_{\chi_{i}^{+}\chi_{j}^{-}} \right] = \frac{1}{2} \left[ W_{\chi_{i}^{-}\chi_{j}^{-}} + W_{\chi_{i}^{-}\chi_{j}^{+}} \right] &, \quad \forall \ i = 1, 2, \ j = 1, 2 \\ W_{\chi_{i}^{0}\tilde{f}'_{k}}^{\prime} &\equiv & W_{\chi_{i}^{0}\tilde{f}_{k}} &, \quad \forall \ i = 1, \dots, 4, \ k = 1, 2 \\ W_{\chi_{c}^{\pm}\tilde{f}'_{k}}^{\prime} &\equiv & \frac{1}{2} \left[ W_{\chi_{c}^{+}\tilde{f}_{k}} + W_{\chi_{c}^{+}\tilde{f}_{k}^{+}} \right] &, \quad \forall \ c = 1, 2, \ k = 1, 2 \\ W_{\tilde{f}'_{k}\tilde{f}'_{l}}^{\prime} &\equiv & \frac{1}{2} \left[ W_{\tilde{f}_{k}\tilde{f}_{l}} + W_{\tilde{f}_{k}\tilde{f}_{k}^{+}} \right] &, \quad \forall \ k = 1, 2, \ l = 1, 2 \\ W_{\tilde{q}'_{k}\tilde{q}'_{l}}^{\prime} &\equiv & \frac{1}{2} \frac{1}{9} \sum_{a,b=1}^{3} \left[ W_{\tilde{q}_{k}^{a}\tilde{q}_{l}^{b}} + W_{\tilde{q}_{k}^{a}\tilde{q}_{l}^{b*}} \right] &, \quad \forall \ k = 1, 2, \ l = 1, 2 \end{array}$$

$$(30.84)$$

We don't list all the possible cases with squarks explicitly, the principle being that we in option b) should take the *average* over the squark colour states (see the squark-squark entry in the list above).

We will choose option b) and the code (dsandwdcoscn, dsandwdcoscn, dsasdwdcossfsf and dsasdwdcossfchi) should thus return W' as defined above. Note again that squarks are assumed to have  $g_i = 6$  degrees of freedom in this convention and the summing over colours should also be taken into account in the code.

### **30.13** Relic density – more details on routines

### **30.13.1** Global parameters

When the relic density has been calculated, the integer variable copart in dsandwcom.h is set to indicate which coannihilating particles that have been included in the calculation. In Table 30.4, the meaning of this variable is shown for the case of supersymmetric particles.

copart			PAW va	PAW variables	
Bit set	Octal value	Decimal value	cop1 bit	cop2 bit	Particle
0	1	1	0	_	$\tilde{\chi}_1^0$
1	2	2	1	_	$ ilde{\chi}^0_2$
2	4	4	2	_	$ ilde{\chi}^0_3$
3	10	8	3	_	$ ilde{\chi}_4^0$
4	20	16	4	_	$\tilde{\chi}_1^{\pm}$
5	40	32	5	_	$\tilde{\chi}_2^{\pm}$
6	100	64	6	_	$\tilde{e}_1$
7	200	128	7	_	$ ilde{\mu}_1$
8	400	256	8	_	$\tilde{ au}_1$
9	1000	512	9	_	$\tilde{e}_2$
10	2000	1024	10	_	$ ilde{\mu}_2$
11	4000	2048	11	_	$ ilde{ au}_2$
12	10000	4096	12	_	$\tilde{\nu}_e$
13	20000	8192	13	_	$\tilde{ u}_{\mu}$
14	40000	16384	14	_	$\tilde{\nu}_{ au}$
15	100000	32768	_	0	$\tilde{u}_1$
16	200000	65536	—	1	$\tilde{c}_1$
17	400000	131072	_	2	${ ilde t}_1$
18	1000000	262144	_	3	$\tilde{u}_2$
19	2000000	524288	_	4	$\tilde{c}_2$
20	4000000	1048576	_	5	$\tilde{t}_2$
21	10000000	2097152	_	6	$\tilde{d}_1$
22	20000000	4197304	_	7	$\tilde{s}_1$
23	40000000	8388608	_	8	$\tilde{b}_1$
24	100000000	16777216	_	9	$\overline{\tilde{d}_2}$
25	200 000 000	33554432	_	10	$\tilde{s}_2$
26	400 000 000	67108864	_	11	$\tilde{b}_2$
		0. 200 00 1			· 4

Table 30.4: The bits of copart are set to indicate which initial states that are included in the coannihilation calculation. In the output file \*.omegaco, the value of copart is written in octal format. In PAW cop1 and cop2 are available. Check if a bit is set with btest(cop1,bit).

# Chapter 31 Silveira-Zee (Scalar Singlet)

The module silveira\_zee implements the scalar singlet DM model of Silveira and Zee [4], also referred to as 'scalar phantoms' (which is the original name in [4]) and 'singlet Higgs DM'. This model adds a gauge-singlet real scalar field S to the standard model, with imposed  $Z_2$  symmetry  $S \to -S$ . Its Lagrangian is

$$\mathcal{L}_{\rm SZ} = \mathcal{L}_{\rm SM} + \frac{1}{2} \partial_{\mu} S \partial^{\mu} S - \frac{1}{2} \mu^2 S^2 - \frac{1}{2} \lambda S^2 H^{\dagger} H, \qquad (31.1)$$

where H is the Standard Model Higgs doublet. After electroweak symmetry breaking, the S boson acquires a tree-level mass

$$m_S = \sqrt{\mu^2 + \frac{1}{2}\lambda v_0^2} \,, \tag{31.2}$$

where  $v_0 = (\sqrt{2}G_F)^{-1/2} = 246.2$  GeV is the Higgs vacuum expectation value. The module uses the S mass  $m_S$  and the S-Higgs coupling constant  $\lambda$  as model parameters. These parameters are set with a call to dsgivemodel\_silveira\_zee, followed as usual by a call to dsmodelsetup to initialize the model for use with DarkSUSY.

# Chapter 32 Self-Interacting Dark Matter

The vdSIDM module implements a simplified model setup where DM is confined to a dark sector without (direct) interactions with the standard model. Besides the DM particle, which is assumed to be non-relativistic, there is also a dark radiation (DR) particle which in general will contribute to the relativistic degrees of freedom in the universe – a quantity which is conventionally stated in terms of the contribution from a single neutrino species,  $\Delta N_{\rm eff}$ , and which is tightly constrained by both BBN and CMB. Finally, the module implements a mediator particle. In principle, the code structure is set up such that arbitrary spin combinations of these three particles can be implemented, but presently only scalar and vector mediators (for Dirac DM and DR) are available. If the mass of the mediator particle is much lighter than the DM mass, the resulting Yukawa potential between DM particles results in a self-scattering rate that is strongly velocity-dependent – as indicated in the name of the module.

# 32.1 vdSIDM/an: Annihilation routines

For the spin combinations that are currently implemented in the vdSIDM module, DM annihilates via the t- and u-channel to a pair of mediators. In the limit of small CMS energies  $\sqrt{s} \rightarrow 2m_{\chi}$ , this gives the following contributions to the invariant rate:

$$W_{\chi\chi\to VV} \xrightarrow{p_{\rm CM}\ll m_{\chi}} 4\pi \alpha_{\chi}^2 \left(1 - \frac{m_V^2}{m_{\chi}^2}\right)^{3/2} \left(1 - \frac{m_V^2}{2m_{\chi}^2}\right)^{-2}$$
(32.1)

$$W_{\chi\chi\to\phi\phi} \xrightarrow{p_{\rm CM}\ll m_{\chi}} 6\pi \alpha_{\chi}^2 p_{\rm CM}^2 \left(1 - \frac{m_{\phi}^2}{m_{\chi}^2}\right)^{1/2} \left(1 - \frac{m_{\phi}^2}{2m_{\chi}^2}\right)^{-4} \left(1 - \frac{8}{9}\frac{m_{\phi}^2}{m_{\chi}^2} + \frac{2}{9}\frac{m_{\phi}^4}{m_{\chi}^4}\right), (32.2)$$

where  $\alpha_{\chi} \equiv g_{\chi}/(4\pi)$  and  $p_{\rm CM}$  is the initial CMS momentum of (each of) the DM particles. Furthermore, DM can annihilate to a pair of DR particles, by a mediator exchange in the *s*-channel. This gives

$$W_{\chi\chi\to V^*\to\tilde{\gamma}\tilde{\gamma}} = 8\pi\alpha_{\chi}s^2 \left(1 + \frac{2m_{\chi}^2}{s}\right)\frac{\Gamma_{V\to\tilde{\gamma}\tilde{\gamma}}}{m_V}\left|D_V\right|^2$$
(32.3)

$$W_{\chi\chi\to\phi^*\to\tilde{\gamma}\tilde{\gamma}} = 8\pi\alpha_{\chi}s^2\left(1-\frac{4m_{\chi}^2}{s}\right)\frac{\Gamma_{\phi\to\tilde{\gamma}\tilde{\gamma}}}{m_{\phi}}\left|D_{\phi}\right|^2, \qquad (32.4)$$

where  $D_{V,\phi}$  is defined as

$$|D_H(s)|^2 = \frac{1}{(s - m_H^2)^2 + m_H^2 \Gamma_H^2},$$
(32.5)

i.e. as the inverse of the denominator of the respective propagator.

The interface function dsanwx adds the full tree-level expressions for these processes, and then multiplies the leading-order expressions in the  $v \to 0$  limit by an enhancement factor S(v) due to the Sommerfeld effect. The factor S(v) itself is provided by the auxiliary function dsansommerfeld residing in src\_models/common/aux, which implements the analytic expressions from Ref. [188, 121] (resulting from approximating the Yukawa potential by a Hulthén potential). The Sommerfeld factor depends on whether the process is *s*-wave dominated (as in the vector mediator case) or *p*-wave dominated (as in the scalar mediator case), which hence is one of the input parameters for dsansommerfeld.

## 32.2 vdSIDM/cr: Cosmic rays

The cosmic-ray source functions in vdSIDM simply return zero, as there are currently no couplings to SM particles implemented. In a future version of the code, the user will be able to add partial widths of the mediator particles to visible channels. The cosmic-ray source functions dscrsource and dscrsource line will then be correspondingly updated.

# 32.3 vdSIDM/ini: Model setup

Currently, the module has two concrete example models fully implemented, where DM is a massive Dirac fermion  $\psi_{\chi}$ , DR is a massless fermion  $\psi_{\tilde{\gamma}}$ , and the mediator is either a vector V or a scalar  $\phi$ . The interaction parts of the respective Lagrangians are thus given by

$$\Delta \mathcal{L}_{\text{vector}} = g_{\chi} \bar{\psi}_{\chi} \sqrt[4]{\psi} \psi_{\chi} + g_{\tilde{\gamma}} \bar{\psi}_{\tilde{\gamma}} \sqrt[4]{\psi} \psi_{\tilde{\gamma}}$$
(32.6)

and

$$\Delta \mathcal{L}_{\text{scalar}} = g_{\chi} \bar{\psi}_{\chi} \psi_{\chi} \phi + g_{\tilde{\gamma}} \bar{\psi}_{\tilde{\gamma}} \psi_{\tilde{\gamma}} \phi \,. \tag{32.7}$$

Calling dsgivemodel\_vdSIDM\_vector or dsgivemodel\_vdSIDM\_scalar sets the DM mass, mediator mass and couplings for these models after which, as usual, the model is set up by a call to dsmodelsetup. This routine calculates for example the total decay width of the mediator, for which we add the contributions from decay to DR and (for heavy mediators, if kinematically allowed) decay into DM particles.

## 32.4 vdSIDM/kd: Kinetic decoupling

DM particles are kept in local thermal equilibrium via scattering with the DR particles. This is dominated by *t*-channel exchange of mediator particles, for which we implement the momentumaveraged scattering amplitude returned from Ref. [55]:

$$\left\langle \left| \mathcal{M} \right|^2 \right\rangle_t^{\text{vector}} = \frac{256}{3} g_\chi^2 g_{\tilde{\gamma}}^2 \left( \frac{m_\chi}{m_V} \right)^4 \left( \frac{\omega}{m_\chi} \right)^2, \qquad (32.8)$$

$$\left\langle \left|\mathcal{M}\right|^{2}\right\rangle_{t}^{\text{scalar}} = \frac{512}{3}g_{\chi}^{2}g_{\tilde{\gamma}}^{2}\left(\frac{m_{\chi}}{m_{\phi}}\right)^{4}\left(\frac{\omega}{m_{\chi}}\right)^{2}, \qquad (32.9)$$

where  $\omega$  is the energy of the DR particle. We also add the amplitudes for DM scattering directly off heat bath vector mediators,  $\left\langle |\mathcal{M}|^2 \right\rangle_t = 64g_{\chi}^4/3$ , and scalar mediators,  $\left\langle |\mathcal{M}|^2 \right\rangle_t = 16g_{\chi}^4/3$  [55]. The interface function dskdm2 returns the sum of these two contributions.

## 32.5 vdSIDM/rd: Relic density

In src\_models/vdSIDM/rd we collect a set of routines to handle the temperature of the dark sector in this module. First of all, the interface function dsrdxi returns the temperature ratio  $\xi = T_{\tilde{\gamma}}/T$ , where T is the photon temperature. This is calculated by assuming that the two sectors were in thermal equilubrium ( $\xi = 1$ ) at very early times when the DM particles were still relativistic, e.g. due to some portal that is effective only at very high energies and hence not relevant for the low-energy Lagrangian. After decoupling of the two sectors at a temperature  $T_{dc} \gg m_{\chi}$ , entropy would be separately conserved, which implies that the temperature evolves as

$$\xi(T) = \frac{\left[g_*^{\rm SM}(T)/g_*^{\rm DS}(T)\right]^{\frac{1}{3}}}{\left[g_*^{\rm SM}(T_{\rm dc})/g_*^{\rm DS}(T_{\rm dc})\right]^{\frac{1}{3}}},$$
(32.10)

where  $g_*^{\text{SM,DS}}$  are the entropy degrees of freedom in the visible and dark sector, respectively. Of course, dsrdxi could be replaced in the usual way by any user-supplied function in order to implement a different scenario for how the temperature of the dark sector evolves with time. Care must be taken, however, to ensure that such an implementation is consistent with the model assumptions. A constant value of  $\xi$  for example, as is often assumed for illustration, is *not* consistent with the assumption that there is no interaction between dark and visible sector.

Any additional relativistic energy density, as provided by the DR particles, is conventionally stated in terms of

$$\Delta N_{\text{eff}} \equiv \frac{\rho_{\text{DS}}}{\rho_{1\nu}} = \frac{4}{7} g_*^{\text{DS}} \left(\frac{T_{\tilde{\gamma}}}{T_{\nu}}\right)^4 \,, \tag{32.11}$$

where  $\rho_{1\nu}$  is the energy density contributed by one massless neutrino species, and  $T_{\nu}$  is the neutrino temperature (which differs from the photon temperature after  $e^{\pm}$  annihilation). For convenience, DarkSUSY provides a function dsrddeltaneff to compute  $\Delta N_{\text{eff}}$  as a function of (photon) temperature.

# 32.6 vdSIDM/si: Dark matter self-interactions

The interface function dssisigtm returns the transfer cross section  $\sigma_T$  for a Yukawa potential, by implementing the explicit expressions for the various scattering regimes discussed in detail in Section 25.1.

# Bibliography

- P. Gondolo, J. Edsjö, P. Ullio, L. Bergström, M. Schelke and E. A. Baltz, *DarkSUSY:* Computing supersymmetric dark matter properties numerically, JCAP 0407 (2004) 008, [astro-ph/0406204].
- [2] T. Bringmann, J. Edsjö, P. Gondolo, P. Ullio and L. Bergström, DarkSUSY 6: An Advanced Tool to Compute Dark Matter Properties Numerically, .
- [3] H. Zhao, Analytical models for galactic nuclei, Mon. Not. Roy. Astron. Soc. 278 (1996) 488-496, [astro-ph/9509122].
- [4] V. Silveira and A. Zee, SCALAR PHANTOMS, Phys. Lett. B161 (1985) 136–140.
- [5] J. McDonald, Gauge singlet scalars as cold dark matter, Phys. Rev. D50 (1994) 3637–3649, [hep-ph/0702143].
- [6] C. P. Burgess, M. Pospelov and T. ter Veldhuis, The Minimal model of nonbaryonic dark matter: A Singlet scalar, Nucl. Phys. B619 (2001) 709–728, [hep-ph/0011335].
- [7] J. M. Cline, K. Kainulainen, P. Scott and C. Weniger, Update on scalar singlet dark matter, Phys. Rev. D88 (2013) 055025, [1306.4710].
- [8] J. Einasto, On the Construction of a Composite Model for the Galaxy and on the Determination of the System of Galactic Parameters, Trudy Astrofizicheskogo Instituta Alma-Ata 5 (1965) 87–100.
- [9] J. F. Navarro, C. S. Frenk and S. D. M. White, The Structure of cold dark matter halos, Astrophys. J. 462 (1996) 563-575, [astro-ph/9508025].
- [10] T. Sjöstrand, S. Mrenna and P. Z. Skands, PYTHIA 6.4 Physics and Manual, JHEP 05 (2006) 026, [hep-ph/0603175].
- [11] M. Cirelli, G. Corcella, A. Hektor, G. Hutsi, M. Kadastik, P. Panci et al., PPPC 4 DM ID: A Poor Particle Physicist Cookbook for Dark Matter Indirect Detection, JCAP 1103 (2011) 051, [1012.4515].
- [12] P. Gondolo and G. Gelmini, Cosmic abundances of stable particles: Improved analysis, Nucl. Phys. B360 (1991) 145–179.
- [13] J. Edsjö and P. Gondolo, Neutralino relic density including coannihilations, Phys. Rev. D56 (1997) 1879–1894, [hep-ph/9704361].
- [14] J. Edsjö, M. Schelke, P. Ullio and P. Gondolo, Accurate relic densities with neutralino, chargino and sfermion coannihilations in mSUGRA, JCAP 0304 (2003) 001, [hep-ph/0301106].

- [15] T. Bringmann, Particle Models and the Small-Scale Structure of Dark Matter, New J. Phys. 11 (2009) 105027, [0903.0189].
- [16] L. Bergström, J. Edsjö and P. Ullio, Possible indications of a clumpy dark matter halo, Phys. Rev. D58 (1998) 083507, [astro-ph/9804050].
- [17] L. Bergström, J. Edsjö and P. Gondolo, Indirect detection of dark matter in km size neutrino telescopes, Phys. Rev. D58 (1998) 103519, [hep-ph/9806293].
- [18] E. A. Baltz and J. Edsjo, Positron propagation and fluxes from neutralino annihilation in the halo, Phys. Rev. D59 (1998) 023511, [astro-ph/9808243].
- [19] L. Bergstrom, J. Edsjo and P. Ullio, Cosmic anti-protons as a probe for supersymmetric dark matter?, Astrophys. J. 526 (1999) 215-235, [astro-ph/9902012].
- [20] T. Bringmann, A. J. Galea and P. Walia, Leading QCD Corrections for Indirect Dark Matter Searches: a Fresh Look, Phys. Rev. D93 (2016) 043529, [1510.02473].
- [21] L. Bergström and P. Gondolo, Limits on direct detection of neutralino dark matter from b —> s gamma decays, Astropart. Phys. 5 (1996) 263–278, [hep-ph/9510252].
- [22] L. Bergström and P. Ullio, Full one loop calculation of neutralino annihilation into two photons, Nucl. Phys. B504 (1997) 27–44, [hep-ph/9706232].
- [23] P. Ullio and L. Bergström, Neutralino annihilation into a photon and a Z boson, Phys. Rev. D57 (1998) 1962–1971, [hep-ph/9707333].
- [24] T. Bringmann, L. Bergström and J. Edsjö, New Gamma-Ray Contributions to Supersymmetric Dark Matter Annihilation, JHEP 01 (2008) 049, [0710.3169].
- [25] L. Bergström, T. Bringmann and J. Edsjö, New Positron Spectral Features from Supersymmetric Dark Matter - a Way to Explain the PAMELA Data?, Phys. Rev. D78 (2008) 103520, [0808.3725].
- [26] T. Bringmann and F. Calore, Significant Enhancement of Neutralino Dark Matter Annihilation from Electroweak Bremsstrahlung, Phys. Rev. Lett. 112 (2014) 071301, [1308.1089].
- [27] T. Bringmann, F. Calore, A. Galea and M. Garny, Electroweak and Higgs Boson Internal Bremsstrahlung: General considerations for Majorana dark matter annihilation and application to MSSM neutralinos, JHEP 09 (2017) 041, [1705.03466].
- [28] M. S. Turner, Probing the Structure of the Galactic Halo with gamma Rays Produced by WIMP Annihilations, Phys. Rev. D34 (1986) 1921.
- [29] J. R. Ipser and P. Sikivie, Estimates of the Density of Dark Matter Near the Center of the Galaxy, Phys. Rev. D35 (1987) 3695.
- [30] K. Freese and J. Silk, Halo gamma-rays from cold dark matter annihilation, Phys. Rev. D40 (1989) 3828–3833.
- [31] V. Berezinsky, A. Bottino and G. Mignola, High-energy gamma radiation from the galactic center due to neutralino annihilation, Phys. Lett. B325 (1994) 136-142, [hep-ph/9402215].
- [32] G. Lake, Detectability of gamma-rays from clumps of dark matter, Nature **346** (1990) 39–40.
- [33] J. Silk and A. Stebbins, Clumpy cold dark matter, Astrophys. J. 411 (1993) 439-449.

- [34] C. Calcaneo-Roldan and B. Moore, The Surface brightness of dark matter: Unique signatures of neutralino annihilation in the galactic halo, Phys. Rev. D62 (2000) 123005, [astro-ph/0010056].
- [35] L. Bergström, J. Edsjö, P. Gondolo and P. Ullio, Clumpy neutralino dark matter, Phys. Rev. D59 (1999) 043506, [astro-ph/9806072].
- [36] L. Bergström, J. Edsjö and C. Gunnarsson, Neutralino gamma-ray signals from accreting halo dark matter, Phys. Rev. D63 (2001) 083515, [astro-ph/0012346].
- [37] E. A. Baltz, C. Briot, P. Salati, R. Taillet and J. Silk, Detection of neutralino annihilation photons from external galaxies, Phys. Rev. D61 (2000) 023514, [astro-ph/9909112].
- [38] D. B. Cline and Y.-T. Gao, Low-energy cosmic γ-rays from cosmological photino annihilation, Astron. Astrophys. 231 (1990) L23–L26.
- [39] Y.-T. Gao, F. W. Stecker and D. B. Cline, The Lightest supersymmetric particle and the extragalactic gamma-ray background, Astron. Astrophys. 249 (1991) 1–4.
- [40] L. Bergström, J. Edsjö and P. Ullio, Spectral gamma-ray signatures of cosmological dark matter annihilation, Phys. Rev. Lett. 87 (2001) 251301, [astro-ph/0105048].
- [41] G. Belanger, K. Kannike, A. Pukhov and M. Raidal, Z<sub>3</sub> Scalar Singlet Dark Matter, JCAP 1301 (2013) 022, [1211.1014].
- [42] G. BÃľlanger, K. Kannike, A. Pukhov and M. Raidal, Minimal semi-annihilating Z<sub>N</sub> scalar dark matter, JCAP 1406 (2014) 021, [1403.4960].
- [43] P. Ko and Y. Tang, Self-interacting scalar dark matter with local Z<sub>3</sub> symmetry, JCAP 1405 (2014) 047, [1402.6449].
- [44] S.-M. Choi and H. M. Lee, SIMP dark matter with gauged Z<sub>3</sub> symmetry, JHEP 09 (2015) 063, [1505.00960].
- [45] C. Arina, T. Bringmann, J. Silk and M. Vollmann, Enhanced Line Signals from Annihilating Kaluza-Klein Dark Matter, Phys. Rev. D90 (2014) 083506, [1409.0007].
- [46] A. L. Fitzpatrick, W. Haxton, E. Katz, N. Lubbers and Y. Xu, The Effective Field Theory of Dark Matter Direct Detection, JCAP 1302 (2013) 004, [1203.3542].
- [47] P. Gondolo and S. Scopel, , in prep. (2018).
- [48] A. V. Maccio', A. A. Dutton and F. C. v. d. Bosch, Concentration, Spin and Shape of Dark Matter Haloes as a Function of the Cosmological Model: WMAP1, WMAP3 and WMAP5 results, Mon. Not. Roy. Astron. Soc. 391 (2008) 1940–1954, [0805.1926].
- [49] J. Schaye et al., The EAGLE project: Simulating the evolution and assembly of galaxies and their environments, Mon. Not. Roy. Astron. Soc. 446 (2015) 521–554, [1407.7040].
- [50] M. Schaller, C. S. Frenk, R. G. Bower, T. Theuns, A. Jenkins, J. Schaye et al., Baryon effects on the internal structure of ΛCDM haloes in the EAGLE simulations, Mon. Not. Roy. Astron. Soc. 451 (2015) 1247–1267, [1409.8617].
- [51] L. Wang, A. A. Dutton, G. S. Stinson, A. V. MacciAš, C. Penzo, X. Kang et al., NIHAO project âĂŞ I. Reproducing the inefficiency of galaxy formation across cosmic time with a large sample of cosmological hydrodynamical simulations, Mon. Not. Roy. Astron. Soc. 454 (2015) 83–94, [1503.04818].

- [52] E. Tollet et al., NIHAO âÅŞ IV: core creation and destruction in dark matter density profiles across cosmic time, Mon. Not. Roy. Astron. Soc. 456 (2016) 3542–3552, [1507.03590].
- [53] T. Bringmann and S. Hofmann, Thermal decoupling of WIMPs from first principles, JCAP 0704 (2007) 016, [hep-ph/0612238].
- [54] T. Binder, T. Bringmann, M. Gustafsson and A. Hryczuk, Early kinetic decoupling of dark matter: when the standard way of calculating the thermal relic density fails, Phys. Rev. D96 (2017) 115010, [1706.07433].
- [55] T. Bringmann, H. T. Ihle, J. Kersten and P. Walia, Suppressing structure formation at dwarf galaxy scales and below: late kinetic decoupling as a compelling alternative to warm dark matter, Phys. Rev. D94 (2016) 103529, [1603.04884].
- [56] J. Kasahara, Neutralino Dark Matter: The Mass of the smallest Halo and the golden Region, Ph.D. thesis, University of Utah, 2009.
- [57] P. Gondolo, J. Hisano and K. Kadota, The Effect of quark interactions on dark matter kinetic decoupling and the mass of the smallest dark halos, Phys. Rev. D86 (2012) 083523, [1205.1914].
- [58] A. M. Green, S. Hofmann and D. J. Schwarz, *The First wimpy halos*, *JCAP* 0508 (2005) 003, [astro-ph/0503387].
- [59] A. Loeb and M. Zaldarriaga, The Small-scale power spectrum of cold dark matter, Phys. Rev. D71 (2005) 103520, [astro-ph/0504112].
- [60] E. Bertschinger, The Effects of Cold Dark Matter Decoupling and Pair Annihilation on Cosmological Perturbations, Phys. Rev. D74 (2006) 063509, [astro-ph/0607319].
- [61] J. Diemand, B. Moore and J. Stadel, Earth-mass dark-matter haloes as the first structures in the early Universe, Nature 433 (2005) 389–391, [astro-ph/0501589].
- [62] K. Griest and D. Seckel, Three exceptions in the calculation of relic abundances, Phys. Rev. D43 (1991) 3191–3203.
- [63] M. Srednicki, R. Watkins and K. A. Olive, Calculations of Relic Densities in the Early Universe, Nucl. Phys. B310 (1988) 693.
- [64] M. Drees, F. Hajkarim and E. R. Schmitz, The Effects of QCD Equation of State on the Relic Density of WIMP Dark Matter, JCAP 1506 (2015) 025, [1503.03513].
- [65] L. M. Krauss.
- [66] W. H. Press and D. N. Spergel, Capture by the sun of a galactic population of weakly interacting massive particles, Astrophys. J. 296 (1985) 679–684.
- [67] J. Silk, K. A. Olive and M. Srednicki, The Photino, the Sun and High-Energy Neutrinos, Phys. Rev. Lett. 55 (1985) 257–259.
- [68] L. M. Krauss, K. Freese, W. Press and D. Spergel, Cold dark matter candidates and the solar neutrino problem, Astrophys. J. 299 (1985) 1001.
- [69] L. M. Krauss, M. Srednicki and F. Wilczek, Solar System Constraints and Signatures for Dark Matter Candidates, Phys. Rev. D33 (1986) 2079–2083.
- [70] T. K. Gaisser, G. Steigman and S. Tilav, Limits on Cold Dark Matter Candidates from Deep Underground Detectors, Phys. Rev. D34 (1986) 2206.

- [71] K. Griest and D. Seckel, Cosmic Asymmetry, Neutrinos and the Sun, Nucl. Phys. B283 (1987) 681–705.
- [72] J. S. Hagelin, K. W. Ng and K. A. Olive, A High-energy Neutrino Signature From Supersymmetric Relics, Phys. Lett. B180 (1986) 375–380.
- [73] K. Freese, Can Scalar Neutrinos Or Massive Dirac Neutrinos Be the Missing Mass?, Phys. Lett. 167B (1986) 295–300.
- [74] M. Kamionkowski, Energetic neutrinos from heavy neutralino annihilation in the sun, Phys. Rev. D44 (1991) 3021–3042.
- [75] F. Halzen, T. Stelzer and M. Kamionkowski, Signatures of dark matter in underground detectors, Phys. Rev. D45 (1992) 4439–4442.
- [76] A. Bottino, V. de Alfaro, N. Fornengo, G. Mignola and M. Pignone, Indirect search for neutralinos at neutrino telescopes, Phys. Lett. B265 (1991) 57–63.
- [77] A. Bottino, N. Fornengo, G. Mignola and L. Moscoso, Signals of neutralino dark matter from earth and sun, Astropart. Phys. 3 (1995) 65–76, [hep-ph/9408391].
- [78] R. Gandhi, J. L. Lopez, D. V. Nanopoulos, K.-j. Yuan and A. Zichichi, Scrutinizing supergravity models through neutrino telescopes, Phys. Rev. D49 (1994) 3691–3703, [astro-ph/9309048].
- [79] L. Bergström, J. Edsjö and P. Gondolo, Indirect neutralino detection rates in neutrino telescopes, Phys. Rev. D55 (1997) 1765–1770, [hep-ph/9607237].
- [80] F. Halzen, Status of neutrino astronomy: The Quest for kilometer scale instruments, Comments Nucl. Part. Phys. 22 (1997) 155–170, [astro-ph/9701029].
- [81] S. Ritz and D. Seckel, Detailed Neutrino Spectra From Cold Dark Matter Annihilations in the Sun, Nucl. Phys. B304 (1988) 877–908.
- [82] G. F. Giudice and E. Roulet, Energetic Neutrinos From Supersymmetric Dark Matter, Nucl. Phys. B316 (1989) 429–442.
- [83] M. Drees, G. Jungman, M. Kamionkowski and M. M. Nojiri, Neutralino annihilation into gluons, Phys. Rev. D49 (1994) 636–647, [hep-ph/9306325].
- [84] G. Jungman and M. Kamionkowski, Neutrinos from particle decay in the sun and earth, Phys. Rev. D51 (1995) 328–340, [hep-ph/9407351].
- [85] V. Berezinsky, A. Bottino, J. R. Ellis, N. Fornengo, G. Mignola and S. Scopel, Searching for relic neutralinos using neutrino telescopes, Astropart. Phys. 5 (1996) 333–352, [hep-ph/9603342].
- [86] M. Kamionkowski, K. Griest, G. Jungman and B. Sadoulet, Model independent comparison of direct versus indirect detection of supersymmetric dark matter, Phys. Rev. Lett. 74 (1995) 5174–5177, [hep-ph/9412213].
- [87] J. Edsjö and P. Gondolo, WIMP mass determination with neutrino telescopes, Phys. Lett. B357 (1995) 595–601, [hep-ph/9504283].
- [88] L. Bergström, J. Edsjö and M. Kamionkowski, Astrophysical neutrino detection with angular and energy resolution, Astropart. Phys. 7 (1997) 147–160, [astro-ph/9702037].
- [89] G. Jungman, M. Kamionkowski and K. Griest, Supersymmetric dark matter, Phys. Rept. 267 (1996) 195–373, [hep-ph/9506380].

- [90] A. Gould, Resonant Enhancements in WIMP Capture by the Earth, Astrophys. J. 321 (1987) 571.
- [91] J. N. Bahcall, M. H. Pinsonneault and S. Basu, Solar models: Current epoch and time dependences, neutrinos, and helioseismological properties, Astrophys. J. 555 (2001) 990-1012, [astro-ph/0010346].
- [92] W. F. McDonough, Compositional Model for the Earth's Core, Treatise on Geochemistry 2 (Dec., 2003) 568.
- [93] The Earth: its properties, composition, and structure. Encyclopædia Britannica, Inc., 1999.
- [94] A. Gould, Gravitational diffusion of solar system WIMPs, Astrophys. J. 368 (Feb., 1991) 610–615.
- [95] P. Farinella, C. Froeschlé, C. Froeschlé, R. Gonczi, G. Hahn, A. Morbidelli et al., Asteroids falling into the Sun, Nature 371 (Sept., 1994) 314–317.
- [96] A. Gould and S. M. Khairul Alam, Can heavy WIMPs be captured by the earth?, Astrophys. J. 549 (2001) 72-75, [astro-ph/9911288].
- [97] J. Lundberg and J. Edsjö, WIMP diffusion in the solar system including solar depletion and its effect on earth capture rates, Phys. Rev. D69 (2004) 123505, [astro-ph/0401113].
- [98] J. Edsjö, "WimpSim Neutrino Monte Carlo." http://www.fysik.su.se/~edsjo/wimpsim/, 2007.
- [99] J. Edsjö, Aspects of neutrino detection of neutralino dark matter, Ph.D. thesis, Uppsala U., 1997. hep-ph/9704384.
- [100] J. Edsjö, "NuSigma Neutrino Interaction Monte Carlo." http://www.fysik.su.se/~edsjo/wimpsim/, 2007.
- [101] PARTICLE DATA GROUP collaboration, D. E. Groom et al., Review of particle physics. Particle Data Group, Eur. Phys. J. C15 (2000) 1–878.
- [102] M. Blennow, J. Edsjö and T. Ohlsson, Neutrinos from WIMP annihilations using a full three-flavor Monte Carlo, JCAP 0801 (2008) 021, [0709.3898].
- [103] D. N. Spergel and P. J. Steinhardt, Observational evidence for selfinteracting cold dark matter, Phys. Rev. Lett. 84 (2000) 3760-3763, [astro-ph/9909386].
- [104] A. Loeb and N. Weiner, Cores in Dwarf Galaxies from Dark Matter with a Yukawa Potential, Phys. Rev. Lett. 106 (2011) 171302, [1011.6374].
- [105] M. Vogelsberger, J. Zavala and A. Loeb, Subhaloes in Self-Interacting Galactic Dark Matter Haloes, Mon. Not. Roy. Astron. Soc. 423 (2012) 3740, [1201.5892].
- [106] A. H. G. Peter, M. Rocha, J. S. Bullock and M. Kaplinghat, Cosmological Simulations with Self-Interacting Dark Matter II: Halo Shapes vs. Observations, Mon. Not. Roy. Astron. Soc. 430 (2013) 105, [1208.3026].
- [107] J. Zavala, M. Vogelsberger and M. G. Walker, Constraining Self-Interacting Dark Matter with the Milky Way's dwarf spheroidals, Mon. Not. Roy. Astron. Soc. 431 (2013) L20–L24, [1211.6426].
- [108] L. G. van den Aarssen, T. Bringmann and C. Pfrommer, Is dark matter with long-range interactions a solution to all small-scale problems of ΛCDM cosmology?, Phys. Rev. Lett. 109 (2012) 231301, [1205.5809].

- [109] O. D. Elbert, J. S. Bullock, S. Garrison-Kimmel, M. Rocha, J. OâĹŹÂśorbe and A. H. G. Peter, Core formation in dwarf haloes with self-interacting dark matter: no fine-tuning necessary, Mon. Not. Roy. Astron. Soc. 453 (2015) 29–37, [1412.1477].
- [110] A. Kamada, M. Kaplinghat, A. B. Pace and H.-B. Yu, How the Self-Interacting Dark Matter Model Explains the Diverse Galactic Rotation Curves, Phys. Rev. Lett. 119 (2017) 111102, [1611.02716].
- [111] A. Robertson, R. Massey, V. Eke, S. Tulin, H.-B. Yu, Y. BahâĹŽÂI et al., The diverse density profiles of galaxy clusters with self-interacting dark matter plus baryons, Mon. Not. Roy. Astron. Soc. 476 (2018) L20, [1711.09096].
- [112] J. S. Bullock and M. Boylan-Kolchin, Small-Scale Challenges to the ΛCDM Paradigm, Ann. Rev. Astron. Astrophys. 55 (2017) 343–387, [1707.04256].
- [113] R. A. Flores and J. R. Primack, Observational and theoretical constraints on singular dark matter halos, Astrophys. J. 427 (1994) L1-4, [astro-ph/9402004].
- [114] B. Moore, Evidence against dissipationless dark matter from observations of galaxy haloes, Nature 370 (1994) 629.
- [115] M. Boylan-Kolchin, J. S. Bullock and M. Kaplinghat, Too big to fail? The puzzling darkness of massive Milky Way subhaloes, Mon. Not. Roy. Astron. Soc. 415 (2011) L40, [1103.0007].
- [116] M. Boylan-Kolchin, J. S. Bullock and M. Kaplinghat, The Milky Way's bright satellites as an apparent failure of LCDM, Mon. Not. Roy. Astron. Soc. 422 (2012) 1203–1218, [1111.2048].
- [117] K. A. Oman et al., The unexpected diversity of dwarf galaxy rotation curves, Mon. Not. Roy. Astron. Soc. 452 (2015) 3650–3665, [1504.01437].
- [118] K. A. Oman, J. F. Navarro, L. V. Sales, A. Fattahi, C. S. Frenk, T. Sawala et al., Missing dark matter in dwarf galaxies?, Mon. Not. Roy. Astron. Soc. 460 (2016) 3610–3623, [1601.01026].
- [119] B. Moore, S. Ghigna, F. Governato, G. Lake, T. R. Quinn, J. Stadel et al., Dark matter substructure within galactic halos, Astrophys. J. 524 (1999) L19–L22, [astro-ph/9907411].
- [120] A. A. Klypin, A. V. Kravtsov, O. Valenzuela and F. Prada, Where are the missing Galactic satellites?, Astrophys. J. 522 (1999) 82–92, [astro-ph/9901240].
- [121] S. Tulin, H.-B. Yu and K. M. Zurek, Beyond Collisionless Dark Matter: Particle Physics Dynamics for Dark Matter Halo Structure, Phys. Rev. D87 (2013) 115007, [1302.3898].
- [122] F. Kahlhoefer, K. Schmidt-Hoberg and S. Wild, Dark matter self-interactions from a general spin-0 mediator, JCAP 1708 (2017) 003, [1704.02149].
- [123] S. Tulin and H.-B. Yu, Dark Matter Self-interactions and Small Scale Structure, Phys. Rept. 730 (2018) 1–57, [1705.02358].
- [124] J. L. Feng, M. Kaplinghat and H.-B. Yu, Halo Shape and Relic Density Exclusions of Sommerfeld-Enhanced Dark Matter Explanations of Cosmic Ray Excesses, Phys. Rev. Lett. 104 (2010) 151301, [0911.0422].
- [125] F.-Y. Cyr-Racine, K. Sigurdson, J. Zavala, T. Bringmann, M. Vogelsberger and C. Pfrommer, ETHOSâĂŽÃĎÃőan effective theory of structure formation: From dark particle physics to the matter distribution of the Universe, Phys. Rev. D93 (2016) 123527, [1512.05344].

- [126] S. A. Khrapak, A. V. Ivlev, G. E. Morfill and S. K. Zhdanov, Scattering in the Attractive Yukawa Potential in the Limit of Strong Interaction, Phys. Rev. Lett. 90 (2003) 225002.
- [127] S. A. Khrapak, A. V. Ivlev and G. E. Morfill, Momentum transfer in complex plasmas, Phys. Rev. E 70 (Nov, 2004) 056405.
- [128] Y. L. Luke, The Special Functions and Their Approximations, vol. II of Mathematics in Science and Engineering, Volume 53-II, Editor: Richard Bellman. 1969.
- [129] K. G. Chetyrkin, J. H. Kuhn and M. Steinhauser, RunDec: A Mathematica package for running and decoupling of the strong coupling and quark masses, Comput. Phys. Commun. 133 (2000) 43-65, [hep-ph/0004189].
- [130] H. E. Haber and G. L. Kane, The Search for Supersymmetry: Probing Physics Beyond the Standard Model, Phys. Rept. 117 (1985) 75–263.
- [131] J. F. Gunion and H. E. Haber, Higgs Bosons in Supersymmetric Models. 1., Nucl. Phys. B272 (1986) 1.
- [132] F. E. Paige, S. D. Protopopescu, H. Baer and X. Tata, ISAJET 7.69: A Monte Carlo event generator for pp, anti-p p, and e+e- reactions, hep-ph/0312045.
- [133] http://www.phy.bnl.gov/~isajet.
- [134] P. Z. Skands et al., SUSY Les Houches accord: Interfacing SUSY spectrum calculators, decay packages, and event generators, JHEP 07 (2004) 036, [hep-ph/0311123].
- [135] B. C. Allanach et al., SUSY Les Houches Accord 2, Comput. Phys. Commun. 180 (2009) 8–25, [0801.0045].
- [136] M. Drees, M. M. Nojiri, D. P. Roy and Y. Yamada, Light Higgsino dark matter, Phys. Rev. D56 (1997) 276–290, [hep-ph/9701219].
- [137] D. Pierce and A. Papadopoulos, Radiative corrections to neutralino and chargino masses in the minimal supersymmetric model, Phys. Rev. D50 (1994) 565–570, [hep-ph/9312248].
- [138] A. B. Lahanas, K. Tamvakis and N. D. Tracas, One loop corrections to the neutralino sector and radiative electroweak breaking in the MSSM, Phys. Lett. B324 (1994) 387–396, [hep-ph/9312251].
- [139] S. Heinemeyer, W. Hollik and G. Weiglein, FeynHiggs: A Program for the calculation of the masses of the neutral CP even Higgs bosons in the MSSM, Comput. Phys. Commun. 124 (2000) 76–89, [hep-ph/9812320].
- [140] http://www.feynhiggs.de.
- [141] S. Heinemeyer, W. Hollik and G. Weiglein, QCD corrections to the masses of the neutral CP even Higgs bosons in the MSSM, Phys. Rev. D58 (1998) 091701, [hep-ph/9803277].
- [142] S. Heinemeyer, W. Hollik and G. Weiglein, The Masses of the neutral CP even Higgs bosons in the MSSM: Accurate analysis at the two loop level, Eur. Phys. J. C9 (1999) 343–366, [hep-ph/9812472].
- [143] S. Heinemeyer, W. Hollik and G. Weiglein, The Mass of the lightest MSSM Higgs boson: A Compact analytical expression at the two loop level, Phys. Lett. B455 (1999) 179–191, [hep-ph/9903404].
- [144] T. Bringmann et al., DarkBit: A GAMBIT module for computing dark matter observables and likelihoods, Eur. Phys. J. C77 (2017) 831, [1705.07920].

- [145] GAMBIT collaboration, C. BalÃązs et al., ColliderBit: a GAMBIT module for the calculation of high-energy collider observables and likelihoods, Eur. Phys. J. C77 (2017) 795, [1705.07919].
- [146] P. Bechtle, O. Brein, S. Heinemeyer, G. Weiglein and K. E. Williams, *HiggsBounds:* Confronting Arbitrary Higgs Sectors with Exclusion Bounds from LEP and the Tevatron, Comput. Phys. Commun. 181 (2010) 138–167, [0811.4169].
- [147] ATLAS collaboration, G. Aad et al., Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using  $\sqrt{s} = 8$  TeV proton-proton collision data, JHEP **09** (2014) 176, [1405.7875].
- [148] F. Mahmoudi, SuperIso: A Program for calculating the isospin asymmetry of B —> K\* gamma in the MSSM, Comput. Phys. Commun. 178 (2008) 745–754, [0710.2067].
- [149] THE GAMBIT FLAVOUR WORKGROUP collaboration, F. U. Bernlochner et al., FlavBit: A GAMBIT module for computing flavour observables and likelihoods, Eur. Phys. J. C77 (2017) 786, [1705.07933].
- [150] BABAR collaboration, J. P. Lees et al., Exclusive Measurements of  $b \rightarrow s\gamma$  Transition Rate and Photon Energy Spectrum, Phys. Rev. D86 (2012) 052012, [1207.2520].
- [151] BABAR collaboration, J. P. Lees et al., Precision Measurement of the  $B \to X_s \gamma$  Photon Energy Spectrum, Branching Fraction, and Direct CP Asymmetry  $A_{CP}(B \to X_{s+d}\gamma)$ , Phys. Rev. Lett. **109** (2012) 191801, [1207.2690].
- [152] BELLE collaboration, A. Abdesselam et al., Measurement of the inclusive B → X<sub>s+d</sub>γ branching fraction, photon energy spectrum and HQE parameters, in Proceedings, 38th International Conference on High Energy Physics (ICHEP 2016): Chicago, IL, USA, August 3-10, 2016, 2016, 1608.02344, https://inspirehep.net/record/1479946/files/arXiv:1608.02344.pdf.
- [153] LHCB collaboration, R. Aaij et al., Measurement of the  $B_s^0 \to \mu^+\mu^-$  branching fraction and effective lifetime and search for  $B^0 \to \mu^+\mu^-$  decays, Phys. Rev. Lett. **118** (2017) 191801, [1703.05747].
- [154] T. Moroi, The Muon anomalous magnetic dipole moment in the minimal supersymmetric standard model, Phys. Rev. D53 (1996) 6565–6575, [hep-ph/9512396].
- [155] MUON G-2 collaboration, G. W. Bennett et al., Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL, Phys. Rev. D73 (2006) 072003, [hep-ex/0602035].
- [156] GAMBIT collaboration, P. Athron et al., SpecBit, DecayBit and PrecisionBit: GAMBIT modules for computing mass spectra, particle decay rates and precision observables, 1705.07936.
- [157] A. C. Hearn, *Reduce 3.5*, 1993.
- [158] J. Edsjö, "form2f, perl script to convert form output to fortran."
- [159] A. Birkedal, K. T. Matchev, M. Perelstein and A. Spray, Robust gamma ray signature of WIMP dark matter, hep-ph/0507194.
- [160] L. Bergström, T. Bringmann, M. Eriksson and M. Gustafsson, Gamma rays from Kaluza-Klein dark matter, Phys. Rev. Lett. 94 (2005) 131301, [astro-ph/0410359].
- [161] L. Bergström, Radiative Processes in Dark Matter Photino Annihilation, Phys. Lett. B225 (1989) 372–380.

- [162] L. Bergström, T. Bringmann, M. Eriksson and M. Gustafsson, Gamma rays from heavy neutralino dark matter, Phys. Rev. Lett. 95 (2005) 241301, [hep-ph/0507229].
- [163] M. W. Goodman and E. Witten, Detectability of Certain Dark Matter Candidates, Phys. Rev. D31 (1985) 3059.
- [164] A. Bottino, F. Donato, G. Mignola, S. Scopel, P. Belli and A. Incicchitti, Exploring the supersymmetric parameter space by direct search for WIMPs, Phys. Lett. B402 (1997) 113-121, [hep-ph/9612451].
- [165] J. R. Ellis and R. A. Flores, Realistic Predictions for the Detection of Supersymmetric Dark Matter, Nucl. Phys. B307 (1988) 883–908.
- [166] J. R. Ellis and R. A. Flores, Elastic supersymmetric relic nucleus scattering revisited, Phys. Lett. B263 (1991) 259–266.
- [167] J. Engel, Nuclear form-factors for the scattering of weakly interacting massive particles, Phys. Lett. B264 (1991) 114–119.
- [168] K. Griest, Cross-Sections, Relic Abundance and Detection Rates for Neutralino Dark Matter, Phys. Rev. D38 (1988) 2357.
- [169] R. Barbieri, M. Frigeni and G. F. Giudice, Dark Matter Neutralinos in Supergravity Theories, Nucl. Phys. B313 (1989) 725–735.
- [170] G. B. Gelmini, P. Gondolo and E. Roulet, Neutralino dark matter searches, Nucl. Phys. B351 (1991) 623–644.
- [171] A. Bottino, V. de Alfaro, N. Fornengo, S. Mignola and S. Scopel, On the neutralino as dark matter candidate. 2. Direct detection., Astropart. Phys. 2 (1994) 77–90, [hep-ph/9309219].
- [172] J. Gasser, H. Leutwyler and M. E. Sainio, Sigma term update, Phys. Lett. B253 (1991) 252–259.
- [173] SPIN MUON collaboration, D. Adams et al., A New measurement of the spin dependent structure function g1(x) of the deuteron, Phys. Lett. B357 (1995) 248–254.
- [174] R. L. Jaffe and A. Manohar, The G(1) Problem: Fact and Fantasy on the Spin of the Proton, Nucl. Phys. B337 (1990) 509–546.
- [175] J. Engel and P. Vogel, Spin dependent cross-sections of weakly interacting massive particles on nuclei, Phys. Rev. D40 (1989) 3132–3135.
- [176] H. E. Haber and D. Wyler, RADIATIVE NEUTRALINO DECAY, Nucl. Phys. B323 (1989) 267–310.
- [177] S. Dimopoulos and S. D. Thomas, Dynamical relaxation of the supersymmetric CP violating phases, Nucl. Phys. B465 (1996) 23–33, [hep-ph/9510220].
- [178] J. R. Ellis, G. Ridolfi and F. Zwirner, Radiative corrections to the masses of supersymmetric Higgs bosons, Phys. Lett. B257 (1991) 83–91.
- [179] J. R. Ellis, G. Ridolfi and F. Zwirner, On radiative corrections to supersymmetric Higgs boson masses and their implications for LEP searches, Phys. Lett. B262 (1991) 477–484.
- [180] A. Brignole, J. R. Ellis, G. Ridolfi and F. Zwirner, The Supersymmetric charged Higgs boson mass and LEP phenomenology, Phys. Lett. B271 (1991) 123–132.

- [181] M. Drees and M. M. Nojiri, One loop corrections to the Higgs sector in minimal supergravity models, Phys. Rev. D45 (1992) 2482–2492.
- [182] M. Carena, J. R. Espinosa, M. Quiros and C. E. M. Wagner, Analytical expressions for radiatively corrected Higgs masses and couplings in the MSSM, Phys. Lett. B355 (1995) 209–221, [hep-ph/9504316].
- [183] M. Carena, M. Quiros and C. E. M. Wagner, Effective potential methods and the Higgs mass spectrum in the MSSM, Nucl. Phys. B461 (1996) 407–436, [hep-ph/9508343].
- [184] T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak and G. Weiglein, *FeynHiggs 2.7, Nucl. Phys. Proc. Suppl.* 205-206 (2010) 152–157, [1007.0956].
- [185] PARTICLE DATA GROUP collaboration, C. Patrignani et al., Review of Particle Physics, Chin. Phys. C40 (2016) 100001.
- [186] H. E. Haber, Perspectives in Higgs Physics II. World Scientific, 1997.
- [187] T. Hahn, SUSY Les Houches Accord 2 I/O made easy, Comput. Phys. Commun. 180 (2009) 1681–1693, [hep-ph/0605049].
- [188] S. Cassel, Sommerfeld factor for arbitrary partial wave processes, J. Phys. G37 (2010) 105009, [0903.5307].